

Section 4: Ryan White Part A Regional Summary Data

Section 4 presents summary data for Ryan White Part A Cleveland and Columbus Planning Regions. Data presented include HIV/AIDS surveillance data on reported persons living with a diagnosis of HIV infection, reported diagnoses of HIV infection and reported cases of AIDS.

The planning regions include the following counties:

Ryan White Part A Cleveland Counties

Ashtabula
Cuyahoga
Geauga
Lake
Lorain
Medina

Ryan White Part A Columbus Counties

Delaware
Fairfield
Franklin
Licking
Madison
Morrow
Pickaway
Union

HIV/AIDS Data by Ryan White Part A Cleveland and Columbus Regions

The following tables present HIV/AIDS data for Ryan White Part A Cleveland and Columbus Regions. There are eight data tables for each Ryan White Part:

- Reported persons living with a diagnosis of HIV infection as of December 31, 2014, by current disease status and selected characteristics
- Reported persons living with a diagnosis of HIV infection as of December 31, 2014, by current disease status and transmission/exposure category
- Reported persons living with a diagnosis of HIV infection by year and selected characteristics, 2010-2014
- Reported persons living with a diagnosis of HIV infection by year and transmission/exposure category, 2010-2014
- Reported diagnoses of HIV infection by year of diagnosis and cumulative diagnoses by selected characteristics, 2010-2014
- Reported diagnoses of HIV infection by year of diagnosis and cumulative diagnoses by transmission/exposure category, 2010-2014
- Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, 2010-2014
- Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission/exposure category, 2010-2014

When viewing these tables, it is important to take into account the limitations of the data. These data represent confidential reports of HIV infection and AIDS diagnoses; not all persons with HIV infection. The HIV diagnosis data represent the earliest date of diagnosis reported to the HIV/AIDS Surveillance Program. The earliest date reported may not be the earliest date an individual became aware of their HIV infection. Individuals may have previously tested anonymously or were diagnosed out-of-state prior to being confidentially tested and reported to Ohio.

Ryan White Part A Cleveland Region

Table 127. Reported persons living with a diagnosis of HIV infection as of December 31, 2014, by current disease status and selected characteristics, Cleveland Part A

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	379.3	3,964	78%	2,000	78%	1,964	78%
Females	100.4	1,122	22%	576	22%	546	22%
Age at end of year							
<13	4.2	14	<1%	11	<1%	3	<1%
13-14	*	2	<1%	-	-	2	<1%
15-19	30.7	43	1%	37	1%	6	<1%
20-24	171.3	235	5%	192	7%	43	2%
25-29	258.3	339	7%	255	10%	84	3%
30-34	295.4	387	8%	264	10%	123	5%
35-39	331.5	407	8%	245	10%	162	6%
40-44	438.1	586	12%	301	12%	285	11%
45-49	560.8	815	16%	364	14%	451	18%
50-54	576.4	949	19%	385	15%	564	22%
55-64	336.1	1,040	20%	412	16%	628	25%
65+	74.8	269	5%	110	4%	159	6%
Race/Ethnicity							
White, not Hispanic	111.4	1,726	34%	853	33%	873	35%
Black/African American, not Hispanic	644.8	2,652	52%	1,362	53%	1,290	51%
Hispanic/Latino	448.2	508	10%	228	9%	280	11%
Asian/Pacific Islander	40.0	19	<1%	13	1%	6	<1%
American Indian/Alaska Native	144.7	5	<1%	3	<1%	2	<1%
Unknown	*	176	3%	117	5%	59	2%
Race/Sex							
White, not Hispanic Males	198.8	1,502	30%	733	28%	769	31%
White, not Hispanic Females	28.2	224	4%	120	5%	104	4%
Black/African American, not Hispanic Males	1,033.3	1,961	39%	1,006	39%	955	38%
Black/African American, not Hispanic Females	311.9	691	14%	356	14%	335	13%
Hispanic/Latino Males	606.9	346	7%	158	6%	188	7%
Hispanic/Latina Females	287.6	162	3%	70	3%	92	4%
Asian/Pacific Islander Males	70.2	16	<1%	11	<1%	5	<1%
Asian/Pacific Islander Females	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Males	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Females	*	2	<1%	1	<1%	1	<1%
Unknown	*	176	3%	117	5%	59	2%
Total	235.2	5,086		2,576		2,510	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Cleveland Region (continued)

Table 128. Reported persons living with a diagnosis of HIV infection as of December 31, 2014, by current disease status and transmission/exposure category, Cleveland Part A

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2,641	67%	1,312	66%	1,329	68%
Injection drug use (IDU)	198	5%	70	4%	128	7%
Male-to-male sexual contact and IDU	191	5%	60	3%	131	7%
Heterosexual contact	275	7%	124	6%	151	8%
Other/unknown	631	16%	415	21%	216	11%
Subtotal	3,936	100%	1,981	100%	1,955	100%
Female adult or adolescent						
Injection drug use	137	13%	48	9%	89	17%
Heterosexual contact	713	66%	343	62%	370	70%
Other/unknown	235	22%	164	30%	71	13%
Subtotal	1,085	100%	555	100%	530	100%
Child (<13 yrs at diagnosis)						
Perinatal	53	82%	29	73%	24	96%
Other/unknown	12	18%	11	28%	1	4%
Subtotal	65	100%	40	100%	25	100%
Total	5,086		2,576		2,510	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	2,505	49%	1,267	49%	1,238	49%
Injection drug use (IDU) only	162	3%	63	2%	99	4%
Heterosexual contact only	988	19%	467	18%	521	21%
MSM & IDU	146	3%	55	2%	91	4%
IDU & Heterosexual contact	173	3%	55	2%	118	5%
MSM & Heterosexual contact	136	3%	45	2%	91	4%
MSM & IDU & Heterosexual contact	45	1%	5	<1%	40	2%
Perinatal exposure	56	1%	31	1%	25	1%
Other/unknown	875	17%	588	23%	287	11%
Total	5,086		2,576		2,510	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Cleveland Region (continued)

Table 129. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Cleveland Part A, 2010-2014

Characteristic	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	3,195	77%	3,360	77%	3,553	77%	3,766	77%	3,964	78%
Females	966	23%	1,007	23%	1,051	23%	1,097	23%	1,122	22%
Age at end of year										
<13	19	<1%	15	<1%	14	<1%	14	<1%	14	<1%
13-14	9	<1%	10	<1%	5	<1%	2	<1%	2	<1%
15-19	43	1%	36	1%	37	1%	41	1%	43	1%
20-24	167	4%	179	4%	205	4%	225	5%	235	5%
25-29	246	6%	259	6%	277	6%	311	6%	339	7%
30-34	298	7%	318	7%	354	8%	373	8%	387	8%
35-39	433	10%	402	9%	389	8%	399	8%	407	8%
40-44	695	17%	688	16%	641	14%	589	12%	586	12%
45-49	863	21%	852	20%	835	18%	858	18%	815	16%
50-54	659	16%	743	17%	846	18%	907	19%	949	19%
55-64	607	15%	716	16%	819	18%	918	19%	1,040	20%
65+	122	3%	149	3%	182	4%	226	5%	269	5%
Race/Ethnicity										
White, not Hispanic	1,485	36%	1,544	35%	1,603	35%	1,679	35%	1,726	34%
Black/African American, not Hispanic	2,099	50%	2,223	51%	2,357	51%	2,499	51%	2,652	52%
Hispanic/Latino	423	10%	438	10%	467	10%	494	10%	508	10%
Asian/Pacific Islander	16	<1%	16	<1%	17	<1%	17	<1%	19	<1%
American Indian/Alaska Native	5	<1%	5	<1%	5	<1%	5	<1%	5	<1%
Unknown	133	3%	141	3%	155	3%	169	3%	176	3%
Race/Sex										
White, not Hispanic Males	1,294	31%	1,343	31%	1,396	30%	1,462	30%	1,502	30%
White, not Hispanic Females	191	5%	201	5%	207	4%	217	4%	224	4%
Black/African American, not Hispanic Males	1,511	36%	1,609	37%	1,712	37%	1,825	38%	1,961	39%
Black/African American, not Hispanic Females	588	14%	614	14%	645	14%	674	14%	691	14%
Hispanic/Latino Males	274	7%	285	7%	310	7%	333	7%	346	7%
Hispanic/Latina Females	149	4%	153	4%	157	3%	161	3%	162	3%
Asian/Pacific Islander Males	13	<1%	13	<1%	14	<1%	14	<1%	16	<1%
Asian/Pacific Islander Females	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	3	<1%	3	<1%
American Indian/Alaska Native Females	2	<1%	2	<1%	2	<1%	2	<1%	2	<1%
Unknown	133	3%	141	3%	155	3%	169	3%	176	3%
Total	4,161		4,367		4,604		4,863		5,086	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Cleveland Region (continued)

Table 130. Reported persons living with a diagnosis of HIV infection by year and transmission/exposure category, Cleveland Part A, 2010-2014

Transmission Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	2,115	67%	2,236	67%	2,372	67%	2,510	67%	2,641	67%
Injection drug use (IDU)	188	6%	190	6%	193	5%	197	5%	198	5%
Male-to-male sexual contact and IDU	175	6%	176	5%	186	5%	189	5%	191	5%
Heterosexual contact	241	8%	246	7%	253	7%	263	7%	275	7%
Other/unknown	449	14%	485	15%	522	15%	580	16%	631	16%
Subtotal	3,168	100%	3,333	100%	3,526	100%	3,739	100%	3,936	100%
Female adult or adolescent										
Injection drug use	126	14%	129	13%	131	13%	134	13%	137	13%
Heterosexual contact	633	68%	655	67%	673	66%	698	66%	713	66%
Other/unknown	172	18%	188	19%	212	21%	228	22%	235	22%
Subtotal	931	100%	972	100%	1,016	100%	1,060	100%	1,085	100%
Child (<13 yrs at diagnosis)										
Perinatal	53	85%	53	85%	53	85%	53	83%	53	82%
Other/unknown	9	15%	9	15%	9	15%	11	17%	12	18%
Subtotal	62	100%	62	100%	62	100%	64	100%	65	100%
Total	4,161		4,367		4,604		4,863		5,086	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Exposure Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	1,988	48%	2,108	48%	2,241	49%	2,375	49%	2,505	49%
Injection drug use (IDU) only	148	4%	150	3%	154	3%	160	3%	162	3%
Heterosexual contact only	874	21%	901	21%	926	20%	961	20%	988	19%
MSM & IDU	134	3%	135	3%	141	3%	144	3%	146	3%
IDU & Heterosexual contact	166	4%	169	4%	170	4%	171	4%	173	3%
MSM & Heterosexual contact	127	3%	128	3%	131	3%	135	3%	136	3%
MSM & IDU & Heterosexual contact	41	1%	41	1%	45	1%	45	1%	45	1%
Perinatal exposure	55	1%	55	1%	55	1%	56	1%	56	1%
Other/unknown	628	15%	680	16%	741	16%	816	17%	875	17%
Total	4,161		4,367		4,604		4,863		5,086	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Cleveland Region (continued)

Table 131. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Cleveland Part A, 2010-2014

Characteristic	Diagnosis of HIV Infection by Year														
	2010			2011			2012			2013			2014		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	15.2	159	82%	17.1	179	79%	18.8	196	82%	21.0	219	83%	19.5	204	88%
Females	3.2	36	18%	4.2	47	21%	3.9	44	18%	4.0	45	17%	2.4	27	12%
Age at diagnosis (yr)															
<13	*	-	-	*	-	-	*	1	<1%	*	1	<1%	*	1	<1%
13-14	*	1	1%	*	-	-	*	-	-	*	1	<1%	*	-	-
15-19	13.6	19	10%	10.0	14	6%	11.4	16	7%	12.1	17	6%	11.4	16	7%
20-24	27.0	37	19%	29.9	41	18%	35.7	49	20%	38.6	53	20%	45.2	62	27%
25-29	19.8	26	13%	23.6	31	14%	25.9	34	14%	31.2	41	16%	27.4	36	16%
30-34	19.8	26	13%	22.1	29	13%	25.2	33	14%	28.2	37	14%	23.7	31	13%
35-39	17.1	21	11%	12.2	15	7%	18.7	23	10%	19.5	24	9%	22.8	28	12%
40-44	12.0	16	8%	21.7	29	13%	14.2	19	8%	17.9	24	9%	14.2	19	8%
45-49	14.5	21	11%	19.3	28	12%	13.8	20	8%	21.3	31	12%	11.7	17	7%
50-54	8.5	14	7%	12.1	20	9%	15.2	25	10%	10.3	17	6%	4.9	8	3%
55-64	4.5	14	7%	5.5	17	8%	5.5	17	7%	5.2	16	6%	3.6	11	5%
65+	*	-	-	*	2	1%	*	3	1%	*	2	1%	*	2	1%
Race/Ethnicity															
White, not Hispanic	3.8	59	30%	4.3	66	29%	4.3	66	28%	5.2	81	31%	3.1	48	21%
Black/African American, not Hispanic	27.0	111	57%	33.3	137	61%	32.3	133	55%	34.8	143	54%	38.4	158	68%
Hispanic/Latino	15.0	17	9%	13.2	15	7%	22.9	26	11%	23.8	27	10%	15.0	17	7%
Asian/Pacific Islander	*	1	1%	*	1	<1%	*	1	<1%	*	-	-	*	2	1%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	7	4%	*	7	3%	*	14	6%	*	13	5%	*	6	3%
Race/Sex															
White, not Hispanic Males	7.0	53	27%	7.3	55	24%	7.9	60	25%	9.4	71	27%	5.3	40	17%
White, not Hispanic Females	0.8	6	3%	1.4	11	5%	0.8	6	3%	1.3	10	4%	1.0	8	3%
Black/African American, not Hispanic Males	45.3	86	44%	55.9	106	47%	53.2	101	42%	61.1	116	44%	73.8	140	61%
Black/African American, not Hispanic Females	11.3	25	13%	14.0	31	14%	14.4	32	13%	12.2	27	10%	8.1	18	8%
Hispanic/Latino Males	21.1	12	6%	19.3	11	5%	40.3	23	10%	38.6	22	8%	28.1	16	7%
Hispanic/Latina Females	8.9	5	3%	*	4	2%	*	3	1%	8.9	5	2%	*	1	<1%
Asian/Pacific Islander Males	*	1	1%	*	1	<1%	*	1	<1%	*	-	-	*	2	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	7	4%	*	7	3%	*	14	6%	*	13	5%	*	6	3%
Total	9.0	195		10.4	226		11.1	240		12.2	264		10.7	231	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Cleveland Region (continued)

Table 132. Reported diagnoses of HIV infection by year of diagnosis and transmission/exposure category, Cleveland Part A, 2010-2014

Transmission Category ^a	Diagnosis of HIV Infection by Year									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	110	70%	132	74%	134	68%	140	64%	135	67%
Injection drug use (IDU)	1	1%	3	2%	4	2%	3	1%	1	<1%
Male-to-male sexual contact and IDU	4	3%	2	1%	10	5%	2	1%	2	1%
Heterosexual contact	3	2%	5	3%	5	3%	11	5%	12	6%
Other/unknown	40	25%	37	21%	43	22%	63	29%	53	26%
Subtotal	158	100%	179	100%	196	100%	219	100%	203	100%
Female adult or adolescent										
Injection drug use	4	11%	4	9%	2	5%	3	7%	3	11%
Heterosexual contact	17	47%	25	53%	19	45%	24	56%	17	63%
Other/unknown	15	42%	18	38%	21	50%	16	37%	7	26%
Subtotal	36	100%	47	100%	42	100%	43	100%	27	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	2	100%	2	100%	1	100%
Subtotal	1	100%	-	-	2	100%	2	100%	1	100%
Total	195		226		240		264		231	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Exposure Category ^a	Diagnosis of HIV Infection by Year									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	110	56%	132	58%	131	55%	137	52%	134	58%
Injection drug use (IDU) only	1	1%	3	1%	5	2%	5	2%	2	1%
Heterosexual contact only	20	10%	30	13%	24	10%	35	13%	29	13%
MSM & IDU	3	2%	2	1%	6	3%	2	1%	2	1%
IDU & Heterosexual contact	4	2%	4	2%	1	<1%	1	<1%	2	1%
MSM & Heterosexual contact	-	-	-	-	3	1%	3	1%	1	<1%
MSM & IDU & Heterosexual contact	1	1%	-	-	4	2%	-	-	-	-
Perinatal exposure	1	1%	-	-	1	<1%	1	<1%	-	-
Other/unknown	55	28%	55	24%	65	27%	80	30%	61	26%
Total	195		226		240		264		231	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Cleveland Region (continued)

Table 133. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Cleveland Part A, 2010-2014

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	74	74%	75	77%	61	70%	88	85%	47	81%	4,013	82%
Females	26	26%	23	23%	26	30%	16	15%	11	19%	881	18%
Age at AIDS diagnosis (yr)												
<13	2	2%	1	1%	1	1%	-	-	-	-	59	1%
13-14	1	1%	-	-	-	-	-	-	-	-	7	<1%
15-19	5	5%	4	4%	5	6%	3	3%	2	3%	103	2%
20-24	16	16%	13	13%	12	14%	13	13%	9	16%	413	8%
25-29	13	13%	15	15%	4	5%	19	18%	9	16%	826	17%
30-34	20	20%	10	10%	8	9%	13	13%	8	14%	969	20%
35-39	11	11%	6	6%	14	16%	11	11%	5	9%	940	19%
40-44	9	9%	12	12%	11	13%	13	13%	8	14%	632	13%
45-49	12	12%	21	21%	10	11%	13	13%	4	7%	444	9%
50-54	6	6%	8	8%	12	14%	7	7%	5	9%	231	5%
55-64	5	5%	7	7%	8	9%	10	10%	6	10%	214	4%
65+	-	-	1	1%	2	2%	2	2%	2	3%	56	1%
Race/Ethnicity												
White, not Hispanic	29	29%	27	28%	20	23%	35	34%	15	26%	1,946	40%
Black/African American, not Hispanic	61	61%	65	66%	55	63%	48	46%	36	62%	2,464	50%
Hispanic/Latino	9	9%	5	5%	11	13%	14	13%	5	9%	408	8%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	8	<1%
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	6	<1%
Unknown	1	1%	1	1%	1	1%	7	7%	2	3%	62	1%
Race/Sex												
White, not Hispanic Males	24	24%	23	23%	16	18%	28	27%	13	22%	1,776	36%
White, not Hispanic Females	5	5%	4	4%	4	5%	7	7%	2	3%	170	3%
Black/African American, not Hispanic Males	44	44%	47	48%	35	40%	42	40%	27	47%	1,893	39%
Black/African American, not Hispanic Females	17	17%	18	18%	20	23%	6	6%	9	16%	571	12%
Hispanic/Latino Males	5	5%	4	4%	9	10%	13	13%	5	9%	289	6%
Hispanic/Latina Females	4	4%	1	1%	2	2%	1	1%	-	-	119	2%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	7	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	3	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	3	<1%
Unknown	1	1%	1	1%	1	1%	7	7%	2	3%	62	1%
Total	100		98		87		104		58		4,894	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Cleveland Region (continued)

Table 134. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission/exposure category, Cleveland Part A, 2010-2014

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	50	68%	46	62%	36	59%	57	65%	29	62%	2,706	68%
Injection drug use (IDU)	2	3%	1	1%	3	5%	4	5%	1	2%	388	10%
Male-to-male sexual contact and IDU	3	4%	3	4%	4	7%	-	-	2	4%	293	7%
Heterosexual contact	3	4%	1	1%	2	3%	8	9%	5	11%	221	6%
Other/unknown	16	22%	23	31%	16	26%	19	22%	10	21%	381	10%
Subtotal	74	100%	74	100%	61	100%	88	100%	47	100%	3,989	100%
Female adult or adolescent												
Injection drug use	5	21%	1	4%	2	8%	3	19%	1	9%	198	23%
Heterosexual contact	11	46%	14	61%	12	48%	9	56%	3	27%	546	65%
Other/unknown	8	33%	8	35%	11	44%	4	25%	7	64%	102	12%
Subtotal	24	100%	23	100%	25	100%	16	100%	11	100%	846	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	1	100%	1	100%	-	-	-	-	51	86%
Other/unknown	-	-	-	-	-	-	-	-	-	-	8	14%
Subtotal	2	100%	1	100%	1	100%	-	-	-	-	59	100%
Total	100		98		87		104		58		4,894	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	47	47%	43	44%	36	41%	57	55%	29	50%	2,554	52%
Injection drug use (IDU) only	5	5%	1	1%	4	5%	3	3%	1	2%	359	7%
Heterosexual contact only	14	14%	15	15%	14	16%	17	16%	8	14%	767	16%
MSM & IDU	3	3%	2	2%	2	2%	-	-	2	3%	241	5%
IDU & Heterosexual contact	2	2%	1	1%	1	1%	4	4%	1	2%	227	5%
MSM & Heterosexual contact	3	3%	3	3%	-	-	-	-	-	-	149	3%
MSM & IDU & Heterosexual contact	-	-	1	1%	2	2%	-	-	-	-	52	1%
Perinatal exposure	2	2%	1	1%	1	1%	-	-	-	-	51	1%
Other/unknown	24	24%	31	32%	27	31%	23	22%	17	29%	494	10%
Total	100		98		87		104		58		4,894	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Columbus Region

Table 135. Reported persons living with a diagnosis of HIV infection as of December 31, 2014, by current disease status and selected characteristics, Columbus Part A

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	443.2	4,283	81%	2,255	80%	2,028	82%
Females	98.6	985	19%	548	20%	437	18%
Age at end of year							
<13	5.2	18	<1%	9	<1%	9	<1%
13-14	*	4	<1%	2	<1%	2	<1%
15-19	14.6	19	<1%	16	1%	3	<1%
20-24	188.4	255	5%	196	7%	59	2%
25-29	301.1	473	9%	310	11%	163	7%
30-34	339.8	491	9%	324	12%	167	7%
35-39	398.3	528	10%	280	10%	248	10%
40-44	489.1	660	13%	347	12%	313	13%
45-49	682.3	889	17%	437	16%	452	18%
50-54	622.8	853	16%	367	13%	486	20%
55-64	369.6	862	16%	409	15%	453	18%
65+	92.1	216	4%	106	4%	110	4%
Race/Ethnicity							
White, not Hispanic	185.9	2,731	52%	1,481	53%	1,250	51%
Black/African American, not Hispanic	684.9	2,035	39%	1,076	38%	959	39%
Hispanic/Latino	321.8	247	5%	123	4%	124	5%
Asian/Pacific Islander	37.3	27	1%	14	<1%	13	1%
American Indian/Alaska Native	152.2	6	<1%	5	<1%	1	<1%
Unknown	*	222	4%	104	4%	118	5%
Race/Sex							
White, not Hispanic Males	339.8	2,459	47%	1,322	47%	1,137	46%
White, not Hispanic Females	36.5	272	5%	159	6%	113	5%
Black/African American, not Hispanic Males	994.4	1,422	27%	745	27%	677	27%
Black/African American, not Hispanic Females	397.7	613	12%	331	12%	282	11%
Hispanic/Latino Males	469.3	188	4%	90	3%	98	4%
Hispanic/Latina Females	160.8	59	1%	33	1%	26	1%
Asian/Pacific Islander Males	62.0	22	<1%	11	<1%	11	<1%
Asian/Pacific Islander Females	13.5	5	<1%	3	<1%	2	<1%
American Indian/Alaska Native Males	*	4	<1%	4	<1%	-	-
American Indian/Alaska Native Females	*	2	<1%	1	<1%	1	<1%
Unknown	*	222	4%	104	4%	118	5%
Total	268.1	5,268		2,803		2,465	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Columbus Region (continued)

Table 136. Reported persons living with a diagnosis of HIV infection as of December 31, 2014, by current disease status and transmission/exposure category, Columbus Part A

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	3,016	71%	1,599	71%	1,417	70%
Injection drug use (IDU)	103	2%	27	1%	76	4%
Male-to-male sexual contact and IDU	146	3%	65	3%	81	4%
Heterosexual contact	232	5%	79	4%	153	8%
Other/unknown	765	18%	474	21%	291	14%
Subtotal	4,262	100%	2,244	100%	2,018	100%
Female adult or adolescent						
Injection drug use	89	9%	43	8%	46	11%
Heterosexual contact	625	66%	319	60%	306	73%
Other/unknown	237	25%	168	32%	69	16%
Subtotal	951	100%	530	100%	421	100%
Child (<13 yrs at diagnosis)						
Perinatal	46	84%	25	86%	21	81%
Other/unknown	9	16%	4	14%	5	19%
Subtotal	55	100%	29	100%	26	100%
Total	5,268		2,803		2,465	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	2,864	54%	1,527	54%	1,337	54%
Injection drug use (IDU) only	94	2%	29	1%	65	3%
Heterosexual contact only	857	16%	398	14%	459	19%
MSM & IDU	117	2%	50	2%	67	3%
IDU & Heterosexual contact	98	2%	41	1%	57	2%
MSM & Heterosexual contact	152	3%	72	3%	80	3%
MSM & IDU & Heterosexual contact	29	1%	15	1%	14	1%
Perinatal exposure	48	1%	26	1%	22	1%
Other/unknown	1,009	19%	645	23%	364	15%
Total	5,268		2,803		2,465	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Columbus Region (continued)

Table 137. Reported persons living with a diagnosis of HIV infection by year and selected characteristics, Columbus Part A, 2010-2014

Characteristic	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Sex										
Males	3,428	81%	3,659	81%	3,882	81%	4,085	81%	4,283	81%
Females	807	19%	855	19%	904	19%	950	19%	985	19%
Age at end of year										
<13	16	<1%	16	<1%	20	<1%	20	<1%	18	<1%
13-14	4	<1%	5	<1%	1	<1%	1	<1%	4	<1%
15-19	37	1%	28	1%	31	1%	23	<1%	19	<1%
20-24	238	6%	248	5%	264	6%	279	6%	255	5%
25-29	326	8%	366	8%	396	8%	429	9%	473	9%
30-34	395	9%	453	10%	485	10%	477	9%	491	9%
35-39	509	12%	483	11%	485	10%	497	10%	528	10%
40-44	747	18%	735	16%	703	15%	678	13%	660	13%
45-49	820	19%	867	19%	886	19%	922	18%	889	17%
50-54	571	13%	632	14%	700	15%	765	15%	853	16%
55-64	478	11%	571	13%	683	14%	774	15%	862	16%
65+	94	2%	110	2%	132	3%	170	3%	216	4%
Race/Ethnicity										
White, not Hispanic	2,280	54%	2,408	53%	2,527	53%	2,633	52%	2,731	52%
Black/African American, not Hispanic	1,580	37%	1,694	38%	1,820	38%	1,928	38%	2,035	39%
Hispanic/Latino	188	4%	207	5%	214	4%	230	5%	247	5%
Asian/Pacific Islander	16	<1%	18	<1%	21	<1%	23	<1%	27	1%
American Indian/Alaska Native	4	<1%	4	<1%	4	<1%	5	<1%	6	<1%
Unknown	167	4%	183	4%	200	4%	216	4%	222	4%
Race/Sex										
White, not Hispanic Males	2,049	48%	2,164	48%	2,272	47%	2,369	47%	2,459	47%
White, not Hispanic Females	231	5%	244	5%	255	5%	264	5%	272	5%
Black/African American, not Hispanic Males	1,079	25%	1,163	26%	1,256	26%	1,336	27%	1,422	27%
Black/African American, not Hispanic Females	501	12%	531	12%	564	12%	592	12%	613	12%
Hispanic/Latino Males	142	3%	157	3%	164	3%	174	3%	188	4%
Hispanic/Latina Females	46	1%	50	1%	50	1%	56	1%	59	1%
Asian/Pacific Islander Males	15	<1%	16	<1%	18	<1%	19	<1%	22	<1%
Asian/Pacific Islander Females	1	<1%	2	<1%	3	<1%	4	<1%	5	<1%
American Indian/Alaska Native Males	3	<1%	3	<1%	3	<1%	4	<1%	4	<1%
American Indian/Alaska Native Females	1	<1%	1	<1%	1	<1%	1	<1%	2	<1%
Unknown	167	4%	183	4%	200	4%	216	4%	222	4%
Total	4,235		4,514		4,786		5,035		5,268	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Columbus Region (continued)

Table 138. Reported persons living with a diagnosis of HIV infection by year and transmission/exposure category, Columbus Part A, 2010-2014

Transmission Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	2,390	70%	2,557	70%	2,720	70%	2,868	71%	3,016	71%
Injection drug use (IDU)	96	3%	99	3%	102	3%	102	3%	103	2%
Male-to-male sexual contact and IDU	124	4%	128	4%	131	3%	138	3%	146	3%
Heterosexual contact	210	6%	214	6%	221	6%	228	6%	232	5%
Other/unknown	588	17%	641	18%	687	18%	728	18%	765	18%
Subtotal	3,408	100%	3,639	100%	3,861	100%	4,064	100%	4,262	100%
Female adult or adolescent										
Injection drug use	80	10%	80	10%	82	9%	87	9%	89	9%
Heterosexual contact	519	67%	550	67%	578	66%	601	66%	625	66%
Other/unknown	180	23%	196	24%	212	24%	229	25%	237	25%
Subtotal	779	100%	826	100%	872	100%	917	100%	951	100%
Child (<13 yrs at diagnosis)										
Perinatal	39	81%	40	82%	44	83%	45	83%	46	84%
Other/unknown	9	19%	9	18%	9	17%	9	17%	9	16%
Subtotal	48	100%	49	100%	53	100%	54	100%	55	100%
Total	4,235		4,514		4,786		5,035		5,268	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Exposure Category ^a	Living with a Diagnosis of HIV Infection									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	2,269	54%	2,435	54%	2,583	54%	2,722	54%	2,864	54%
Injection drug use (IDU) only	88	2%	90	2%	93	2%	93	2%	94	2%
Heterosexual contact only	729	17%	764	17%	799	17%	829	16%	857	16%
MSM & IDU	102	2%	106	2%	108	2%	112	2%	117	2%
IDU & Heterosexual contact	88	2%	89	2%	91	2%	96	2%	98	2%
MSM & Heterosexual contact	121	3%	122	3%	137	3%	146	3%	152	3%
MSM & IDU & Heterosexual contact	22	1%	22	<1%	23	<1%	26	1%	29	1%
Perinatal exposure	41	1%	42	1%	46	1%	47	1%	48	1%
Other/unknown	775	18%	844	19%	906	19%	964	19%	1,009	19%
Total	4,235		4,514		4,786		5,035		5,268	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Columbus Region (continued)

Table 139. Reported diagnoses of HIV infection by year of diagnosis and selected characteristics, Columbus Part A, 2010-2014

Characteristic	Diagnosis of HIV Infection by Year														
	2010			2011			2012			2013			2014		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	25.1	243	86%	24.3	235	82%	23.9	231	83%	21.4	207	83%	21.1	204	87%
Females	4.0	40	14%	5.0	50	18%	4.9	49	18%	4.3	43	17%	3.1	31	13%
Age at diagnosis (yr)															
<13	*	1	<1%	*	1	<1%	*	4	1%	*	1	<1%	*	1	<1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	10.0	13	5%	13.1	17	6%	12.3	16	6%	11.6	15	6%	5.4	7	3%
20-24	44.3	60	21%	40.6	55	19%	42.9	58	21%	46.5	63	25%	37.7	51	22%
25-29	26.7	42	15%	28.0	44	15%	29.9	47	17%	21.6	34	14%	31.2	49	21%
30-34	27.7	40	14%	32.5	47	16%	27.7	40	14%	15.9	23	9%	27.7	40	17%
35-39	28.7	38	13%	30.2	40	14%	20.4	27	10%	16.6	22	9%	20.4	27	11%
40-44	26.7	36	13%	22.2	30	11%	20.7	28	10%	21.5	29	12%	13.3	18	8%
45-49	19.2	25	9%	16.1	21	7%	19.2	25	9%	16.9	22	9%	19.2	25	11%
50-54	8.8	12	4%	11.0	15	5%	11.0	15	5%	16.8	23	9%	5.1	7	3%
55-64	6.4	15	5%	5.1	12	4%	7.7	18	6%	6.9	16	6%	3.4	8	3%
65+	*	1	<1%	*	3	1%	*	2	1%	*	2	1%	*	2	1%
Race/Ethnicity															
White, not Hispanic	9.5	139	49%	9.1	133	47%	8.6	127	45%	7.4	109	44%	6.6	97	41%
Black/African American, not Hispanic	39.0	116	41%	38.7	115	40%	41.1	122	44%	35.7	106	42%	36.7	109	46%
Hispanic/Latino	18.2	14	5%	28.7	22	8%	10.4	8	3%	18.2	14	6%	23.4	18	8%
Asian/Pacific Islander	*	2	1%	*	2	1%	*	4	1%	*	2	1%	*	4	2%
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	<1%	*	1	<1%
Unknown	*	12	4%	*	13	5%	*	19	7%	*	18	7%	*	6	3%
Race/Sex															
White, not Hispanic Males	17.5	127	45%	16.6	120	42%	16.4	119	43%	13.8	100	40%	12.4	90	38%
White, not Hispanic Females	1.6	12	4%	1.7	13	5%	1.1	8	3%	1.2	9	4%	0.9	7	3%
Black/African American, not Hispanic Males	63.6	91	32%	58.0	83	29%	60.8	87	31%	56.6	81	32%	63.6	91	39%
Black/African American, not Hispanic Females	16.2	25	9%	20.8	32	11%	22.7	35	13%	16.2	25	10%	11.7	18	8%
Hispanic/Latino Males	32.5	13	5%	44.9	18	6%	20.0	8	3%	22.5	9	4%	37.4	15	6%
Hispanic/Latina Females	*	1	<1%	*	4	1%	*	-	-	13.6	5	2%	*	3	1%
Asian/Pacific Islander Males	*	1	<1%	*	1	<1%	*	3	1%	*	1	<1%	*	3	1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	1	<1%	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	<1%
Unknown	*	12	4%	*	13	5%	*	19	7%	*	18	7%	*	6	3%
Total	14.4	283		14.5	285		14.2	280		12.7	250		12.0	235	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Columbus Region (continued)

Table 140. Reported diagnoses of HIV infection by year of diagnosis and transmission/exposure category, Columbus Part A, 2010-2014

Transmission Category ^a	Diagnosis of HIV Infection by Year									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	176	73%	165	70%	168	73%	148	71%	151	74%
Injection drug use (IDU)	-	-	3	1%	2	1%	-	-	1	<1%
Male-to-male sexual contact and IDU	8	3%	6	3%	2	1%	7	3%	8	4%
Heterosexual contact	10	4%	4	2%	4	2%	7	3%	5	2%
Other/unknown	48	20%	57	24%	54	23%	45	22%	39	19%
Subtotal	242	100%	235	100%	230	100%	207	100%	204	100%
Female adult or adolescent										
Injection drug use	3	8%	-	-	1	2%	5	12%	2	7%
Heterosexual contact	28	70%	32	65%	28	61%	22	52%	22	73%
Other/unknown	9	23%	17	35%	17	37%	15	36%	6	20%
Subtotal	40	100%	49	100%	46	100%	42	100%	30	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	1	100%	4	100%	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	4	100%	1	100%	1	100%
Total	283		285		280		250		235	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Exposure Category ^a	Diagnosis of HIV Infection by Year									
	2010		2011		2012		2013		2014	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	171	60%	163	57%	154	55%	137	55%	145	62%
Injection drug use (IDU) only	-	-	2	1%	3	1%	1	<1%	1	<1%
Heterosexual contact only	38	13%	36	13%	32	11%	29	12%	27	11%
MSM & IDU	8	3%	6	2%	2	1%	3	1%	5	2%
IDU & Heterosexual contact	3	1%	1	<1%	-	-	4	2%	2	1%
MSM & Heterosexual contact	5	2%	2	1%	14	5%	11	4%	6	3%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	4	2%	3	1%
Perinatal exposure	1	<1%	1	<1%	4	1%	1	<1%	1	<1%
Other/unknown	57	20%	74	26%	71	25%	60	24%	45	19%
Total	283		285		280		250		235	

Notes: Diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and a later AIDS diagnosis, and concurrent diagnoses of HIV and AIDS. Diagnoses of HIV infection by year (2010-2014) represent all reported cases diagnosed in each year.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Columbus Region (continued)

Table 141. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by selected characteristics, Columbus Part A, 2010-2014

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	139	86%	130	81%	120	81%	118	83%	92	84%	3,667	87%
Females	23	14%	31	19%	28	19%	24	17%	17	16%	555	13%
Age at AIDS diagnosis (yr)												
<13	1	1%	1	1%	1	1%	-	-	-	-	28	1%
13-14	-	-	-	-	-	-	-	-	-	-	6	<1%
15-19	10	6%	8	5%	4	3%	3	2%	4	4%	82	2%
20-24	21	13%	22	14%	21	14%	33	23%	19	17%	421	10%
25-29	17	10%	16	10%	25	17%	18	13%	18	17%	715	17%
30-34	31	19%	26	16%	23	16%	17	12%	24	22%	940	22%
35-39	24	15%	39	24%	24	16%	20	14%	7	6%	750	18%
40-44	28	17%	23	14%	16	11%	14	10%	11	10%	546	13%
45-49	14	9%	13	8%	16	11%	13	9%	13	12%	361	9%
50-54	8	5%	6	4%	7	5%	14	10%	6	6%	206	5%
55-64	7	4%	7	4%	10	7%	9	6%	5	5%	142	3%
65+	1	1%	-	-	1	1%	1	1%	2	2%	25	1%
Race/Ethnicity												
White, not Hispanic	80	49%	75	47%	77	52%	63	44%	44	40%	2,639	63%
Black/African American, not Hispanic	73	45%	65	40%	50	34%	64	45%	52	48%	1,335	32%
Hispanic/Latino	4	2%	10	6%	8	5%	4	3%	6	6%	126	3%
Asian/Pacific Islander	-	-	1	1%	-	-	-	-	3	3%	16	<1%
American Indian/Alaska Native	-	-	1	1%	-	-	-	-	-	-	2	<1%
Unknown	5	3%	9	6%	13	9%	11	8%	4	4%	104	2%
Race/Sex												
White, not Hispanic Males	78	48%	68	42%	68	46%	57	40%	40	37%	2,461	58%
White, not Hispanic Females	2	1%	7	4%	9	6%	6	4%	4	4%	178	4%
Black/African American, not Hispanic Males	53	33%	46	29%	34	23%	48	34%	40	37%	996	24%
Black/African American, not Hispanic Females	20	12%	19	12%	16	11%	16	11%	12	11%	339	8%
Hispanic/Latino Males	4	2%	7	4%	8	5%	3	2%	6	6%	107	3%
Hispanic/Latina Females	-	-	3	2%	-	-	1	1%	-	-	19	<1%
Asian/Pacific Islander Males	-	-	1	1%	-	-	-	-	2	2%	13	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	1%	3	<1%
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	1	1%	-	-	-	-	-	-	2	<1%
Unknown	5	3%	9	6%	13	9%	11	8%	4	4%	104	2%
Total	162		161		148		142		109		4,222	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Ryan White Part A Columbus Region (continued)

Table 142. Reported cases of AIDS by year of AIDS diagnosis and cumulative AIDS diagnoses by transmission/exposure category, Columbus Part A, 2010-2014

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	86	62%	90	69%	78	65%	76	64%	64	70%	2,642	72%
Injection drug use (IDU)	-	-	2	2%	2	2%	-	-	-	-	183	5%
Male-to-male sexual contact and IDU	6	4%	4	3%	2	2%	2	2%	-	-	138	4%
Heterosexual contact	9	7%	5	4%	3	3%	8	7%	7	8%	192	5%
Other/unknown	37	27%	29	22%	35	29%	32	27%	21	23%	498	14%
Subtotal	138	100%	130	100%	120	100%	118	100%	92	100%	3,653	100%
Female adult or adolescent												
Injection drug use	2	9%	-	-	1	4%	1	4%	1	6%	75	14%
Heterosexual contact	17	74%	21	70%	22	81%	13	54%	12	71%	361	67%
Other/unknown	4	17%	9	30%	4	15%	10	42%	4	24%	101	19%
Subtotal	23	100%	30	100%	27	100%	24	100%	17	100%	537	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	1	100%	1	100%	-	-	-	-	26	81%
Other/unknown	-	-	-	-	-	-	-	-	-	-	6	19%
Subtotal	1	100%	1	100%	1	100%	-	-	-	-	32	100%
Total	162		161		148		142		109		4,222	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Exposure Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2010		2011		2012		2013		2014		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sex (MSM) only	82	51%	88	55%	74	50%	74	52%	62	57%	2,556	61%
Injection drug use (IDU) only	-	-	2	1%	3	2%	-	-	-	-	180	4%
Heterosexual contact only	26	16%	26	16%	25	17%	21	15%	19	17%	553	13%
MSM & IDU	5	3%	4	2%	2	1%	2	1%	-	-	118	3%
IDU & Heterosexual contact	2	1%	-	-	-	-	1	1%	1	1%	78	2%
MSM & Heterosexual contact	4	2%	2	1%	4	3%	2	1%	2	2%	86	2%
MSM & IDU & Heterosexual contact	1	1%	-	-	-	-	-	-	-	-	20	<1%
Perinatal exposure	1	1%	1	1%	1	1%	-	-	-	-	27	1%
Other/unknown	41	25%	38	24%	39	26%	42	30%	25	23%	604	14%
Total	162		161		148		142		109		4,222	

Note: AIDS diagnoses by year (2010-2014) represent all reported AIDS cases diagnosed in each year; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2014.

Dash (-) indicates no cases were reported for the given category.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.