

Persons Living with Diagnosed HIV Infection Reported in Delaware County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Delaware County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	102.7	109	81%	61	82%	48	79%
Females	24.2	26	19%	13	18%	13	21%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	1	1%	1	1%	-	-
15-19	*	-	-	-	-	-	-
20-24	*	2	1%	1	1%	1	2%
25-29	87.6	8	6%	7	9%	1	2%
30-34	60.1	7	5%	5	7%	2	3%
35-39	50.9	8	6%	5	7%	3	5%
40-44	97.3	16	12%	7	9%	9	15%
45-49	89.6	15	11%	6	8%	9	15%
50-54	169.4	26	19%	19	26%	7	11%
55-64	127.2	33	24%	13	18%	20	33%
65+	60.5	19	14%	10	14%	9	15%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	4	3%	4	5%	-	-
Black/African-American	259.5	22	16%	9	12%	13	21%
Hispanic/Latinx	*	3	2%	3	4%	-	-
White	58.8	104	77%	57	77%	47	77%
Multi-Race	*	2	1%	1	1%	1	2%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	3	2%	3	4%	-	-
Asian/Pacific Islander Females	*	1	1%	1	1%	-	-
Black/African-American Males	223.5	10	7%	5	7%	5	8%
Black/African-American Females	299.8	12	9%	4	5%	8	13%
Hispanic/Latino Males	*	2	1%	2	3%	-	-
Hispanic/Latina Females	*	1	1%	1	1%	-	-
White Males	105.2	92	68%	50	68%	42	69%
White Females	13.4	12	9%	7	9%	5	8%
Multi-Race Males	*	2	1%	1	1%	1	2%
Multi-Race Females	*	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-
Total	63.2	135		74		61	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^bHispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Delaware County

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	68	63%	37	62%	31	65%
Injection drug use (IDU)	1	1%	-	-	1	2%
Male-to-male sexual contact and IDU	7	6%	2	3%	5	10%
Heterosexual contact	2	2%	2	3%	-	-
Other/unknown	30	28%	19	32%	11	23%
Subtotal	108	100%	60	100%	48	100%
Female adult or adolescent						
Injection drug use	3	12%	2	15%	1	8%
Heterosexual contact	22	85%	10	77%	12	92%
Other/unknown	1	4%	1	8%	-	-
Subtotal	26	100%	13	100%	13	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-
Total	135		74		61	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Delaware County

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	64	47%	37	50%	27	44%
Injection drug use (IDU) only	2	1%	2	3%	-	-
Heterosexual contact only	24	18%	12	16%	12	20%
Male-to-male sexual contact & IDU	7	5%	2	3%	5	8%
IDU & Heterosexual contact	2	1%	-	-	2	3%
Male-to-male sexual contact & Heterosexual contact	4	3%	-	-	4	7%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	1	1%	1	1%	-	-
Other/unknown	31	23%	20	27%	11	18%
Total	135		74		61	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Delaware County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
13-14	*	-	-	*	-	-	*	1	5%	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
20-24	*	-	-	*	1	25%	*	-	-	*	-	-	*	1	1%	*	-	-	-	-
25-29	*	-	-	*	-	-	*	2	9%	*	-	-	83.8	6	6%	*	-	-	-	-
30-34	*	-	-	*	-	-	*	2	9%	*	-	-	55.1	5	5%	*	-	-	-	-
35-39	*	-	-	*	1	25%	*	2	9%	*	-	-	41.2	5	5%	*	-	-	-	-
40-44	*	-	-	*	-	-	875.9	6	27%	*	1	33%	61.2	8	8%	*	1	50%	-	-
45-49	*	-	-	*	-	-	*	2	9%	*	1	33%	87.5	12	12%	*	-	-	-	-
50-54	*	-	-	*	1	25%	*	3	14%	*	1	33%	157.8	21	20%	*	-	-	-	-
55-64	*	-	-	*	-	-	*	3	14%	*	-	-	124.1	29	28%	*	1	50%	-	-
65+	*	-	-	*	1	25%	*	1	5%	*	-	-	58.7	17	16%	*	-	-	-	-
Total	*	-	-	*	4		259.5	22		*	3		58.8	104		*	2		-	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Delaware County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	2	22%	2	100%	63	68%	1	50%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	1%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	1	33%	1	11%	-	-	4	4%	1	50%	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	2	2%	-	-	-	-
Other/unknown	-	-	2	67%	6	67%	-	-	22	24%	-	-	-	-
Subtotal	-	-	3	100%	9	100%	2	100%	92	100%	2	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	-	-	-	-	3	25%	-	-	-	-
Heterosexual contact	-	-	1	100%	12	100%	1	100%	8	67%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	8%	-	-	-	-
Subtotal	-	-	1	100%	12	100%	1	100%	12	100%	-	-	-	-
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	1	100%	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-	-	-	-	-
Total	-	-	4		22		3		104		2		-	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Delaware County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	2	9%	2	67%	59	57%	1	50%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	2	2%	-	-	-	-
Heterosexual contact only	-	-	1	25%	12	55%	1	33%	10	10%	-	-	-	-
Male-to-male sexual contact & IDU	-	-	1	25%	1	5%	-	-	4	4%	1	50%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	2	2%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	4	4%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	1	5%	-	-	-	-	-	-	-	-
Other/unknown	-	-	2	50%	6	27%	-	-	23	22%	-	-	-	-
Total	-	-	4		22		3		104		2		-	

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Delaware County, 2016-2020

Living with Diagnosed HIV Infection

Characteristic	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	124.4	121	86%	115.8	115	83%	109.3	111	85%	116.4	121	85%	102.7	109	81%
Females	20.2	20	14%	23.7	24	17%	18.4	19	15%	20.9	22	15%	24.2	26	19%
Age at end of year															
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	1	1%	*	1	1%
15-19	*	1	1%	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	2	1%	*	3	2%	*	2	2%	*	3	2%	*	2	1%
25-29	*	4	3%	*	4	3%	*	2	2%	57.0	5	3%	87.6	8	6%
30-34	*	3	2%	*	3	2%	*	4	3%	51.6	6	4%	60.1	7	5%
35-39	124.4	18	13%	100.5	15	11%	32.8	5	4%	64.2	10	7%	50.9	8	6%
40-44	84.5	13	9%	78.0	12	9%	108.8	17	13%	81.1	13	9%	97.3	16	12%
45-49	160.9	26	18%	151.1	25	18%	136.8	23	18%	131.1	22	15%	89.6	15	11%
50-54	245.5	35	25%	211.0	30	22%	187.9	27	21%	178.0	26	18%	169.4	26	19%
55-64	109.8	26	18%	140.6	34	24%	141.3	35	27%	157.2	40	28%	127.2	33	24%
65+	52.8	13	9%	49.1	13	9%	53.5	15	12%	57.5	17	12%	60.5	19	14%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	1	1%	*	2	1%	*	2	2%	*	2	1%	*	4	3%
Black/African-American	257.5	17	12%	284.8	20	14%	216.0	16	12%	280.0	22	15%	259.5	22	16%
Hispanic/Latinx	*	4	3%	*	4	3%	*	3	2%	*	2	1%	*	3	2%
White	69.3	117	83%	64.3	110	79%	62.4	108	83%	65.6	115	80%	58.8	104	77%
Multi-Race	*	2	1%	*	3	2%	*	1	1%	*	2	1%	*	2	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	3	2%
Asian/Pacific Islander Females	*	-	-	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Black/African-American Males	234.1	8	6%	219.2	8	6%	180.3	7	5%	288.9	12	8%	223.5	10	7%
Black/African-American Females	282.6	9	6%	355.7	12	9%	255.2	9	7%	270.1	10	7%	299.8	12	9%
Hispanic/Latino Males	*	2	1%	*	2	1%	*	1	1%	*	1	1%	*	2	1%
Hispanic/Latina Females	*	2	1%	*	2	1%	*	2	2%	*	1	1%	*	1	1%
White Males	129.6	108	77%	119.7	101	73%	118.2	101	78%	121.2	105	73%	105.2	92	68%
White Females	10.5	9	6%	10.4	9	6%	8.0	7	5%	11.3	10	7%	13.4	12	9%
Multi-Race Males	*	2	1%	*	3	2%	*	1	1%	*	2	1%	*	2	1%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	71.8	141		69.3	139		63.5	130		68.4	143		63.2	135	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Delaware County, 2016-2020

Transmission Category ^a	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	85	70%	79	69%	75	68%	84	70%	68	63%
Injection drug use (IDU)	1	1%	-	-	1	1%	-	-	1	1%
Male-to-male sexual contact and IDU	6	5%	9	8%	9	8%	8	7%	7	6%
Heterosexual contact	3	2%	2	2%	1	1%	2	2%	2	2%
Other/unknown	26	21%	25	22%	25	23%	26	22%	30	28%
Subtotal	121	100%	115	100%	111	100%	120	100%	108	100%
Female adult or adolescent										
Injection drug use	1	5%	-	-	-	-	2	9%	3	12%
Heterosexual contact	18	90%	24	100%	19	100%	20	91%	22	85%
Other/unknown	1	5%	-	-	-	-	-	-	1	4%
Subtotal	20	100%	24	100%	19	100%	22	100%	26	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	1	100%	1	100%
Total	141		139		130		143		135	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Delaware County, 2016-2020

Exposure Category ^b	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	79	56%	75	54%	72	55%	80	56%	64	47%
Injection drug use (IDU) only	1	1%	-	-	-	-	1	1%	2	1%
Heterosexual contact only	21	15%	26	19%	20	15%	22	15%	24	18%
Male-to-male sexual contact & IDU	6	4%	9	6%	8	6%	7	5%	7	5%
IDU & Heterosexual contact	1	1%	-	-	1	1%	1	1%	2	1%
Male-to-male sexual contact & Heterosexual contact	6	4%	4	3%	3	2%	4	3%	4	3%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	1	1%	1	1%	-	-
Perinatal exposure	-	-	-	-	-	-	1	1%	1	1%
Other/unknown	27	19%	25	18%	25	19%	26	18%	31	23%
Total	141		139		130		143		135	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.