



Department of
Health

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 9

Counties: Clark, Darke, Greene, Miami, Montgomery, Preble

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2025

Reported new diagnoses of HIV infection in 2024 by disease status and selected characteristics, HIV Prevention Planning Region 9

Characteristic	2024 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	15.4	79	74%	59	73%	18	78%	2	67%
Females	5.2	28	26%	22	27%	5	22%	1	33%
Age at diagnosis (years)									
0-9	*	-	-	-	-	-	-	-	-
10-14	*	-	-	-	-	-	-	-	-
15-19	11.7	8	7%	8	10%	-	-	-	-
20-24	17.7	12	11%	12	15%	-	-	-	-
25-29	23.7	16	15%	14	17%	2	9%	-	-
30-34	31.2	22	21%	19	23%	3	13%	-	-
35-39	13.5	9	8%	4	5%	5	22%	-	-
40-44	23.4	15	14%	9	11%	5	22%	1	33%
45-49	19.1	11	10%	7	9%	4	17%	-	-
50-54	9.8	6	6%	3	4%	1	4%	2	67%
55-64	5.3	7	7%	4	5%	3	13%	-	-
65+	*	1	1%	1	1%	-	-	-	-
Race/Ethnicity ^b									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	1	1%	1	1%	-	-	-	-
Black/African American	43.1	63	59%	49	60%	11	48%	3	100%
Hispanic/Latino	19.5	8	7%	7	9%	1	4%	-	-
White	4.2	34	32%	24	30%	10	43%	-	-
Multi-Race	*	1	1%	-	-	1	4%	-	-
Race/Ethnicity ^b and Sex									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	1	1%	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African American Males	63.5	44	41%	34	42%	8	35%	2	67%
Black/African American Females	24.7	19	18%	15	19%	3	13%	1	33%
Hispanic/Latino Males	23.6	5	5%	5	6%	-	-	-	-
Hispanic/Latina Females	*	3	3%	2	2%	1	4%	-	-
White Males	7.1	28	26%	19	23%	9	39%	-	-
White Females	1.5	6	6%	5	6%	1	4%	-	-
Multi-Race Males	*	1	1%	-	-	1	4%	-	-
Multi-Race Females	*	-	-	-	-	-	-	-	-
Total	10.2	107		81		23		3	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by disease status and transmission category, HIV Prevention Planning Region 9

Transmission Category ^a	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	43	54%	34	58%	9	50%	-	-
Injection drug use (IDU)	5	6%	4	7%	1	6%	-	-
Male-to-male sexual contact and IDU	4	5%	2	3%	2	11%	-	-
Heterosexual contact	6	8%	5	8%	1	6%	-	-
Other/unknown	21	27%	14	24%	5	28%	2	100%
Subtotal	79	100%	59	100%	18	100%	2	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	28	100%	22	100%	5	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	28	100%	22	100%	5	100%	1	100%
Child (<13 years at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	107		81		23		3	

Reported new diagnoses of HIV infection in 2024 by disease status and exposure category, HIV Prevention Planning Region 9

Exposure Category ^b	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	32	30%	27	33%	5	22%	-	-
Injection drug use (IDU) only	1	1%	1	1%	-	-	-	-
Heterosexual contact only	54	50%	40	49%	11	48%	3	100%
Male-to-male sexual contact & IDU	3	3%	2	2%	1	4%	-	-
IDU & Heterosexual contact	4	4%	3	4%	1	4%	-	-
Male-to-male sexual contact & Heterosexual contact	11	10%	7	9%	4	17%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	1	1%	-	-	1	4%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	1	1%	1	1%	-	-	-	-
Total	107		81		23		3	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 9

2024 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
0-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	53.1	6	10%	*	2	25%	*	-	-	*	-	-
20-24	*	-	-	*	-	-	84.0	9	14%	*	1	13%	*	2	6%	*	-	-
25-29	*	-	-	*	1	100%	86.6	9	14%	*	3	38%	*	3	9%	*	-	-
30-34	*	-	-	*	-	-	103.4	12	19%	*	1	13%	17.6	9	26%	*	-	-
35-39	*	-	-	*	-	-	52.4	5	8%	*	1	13%	*	3	9%	*	-	-
40-44	*	-	-	*	-	-	71.9	6	10%	*	-	-	16.1	8	24%	*	1	100%
45-49	*	-	-	*	-	-	93.9	7	11%	*	-	-	*	4	12%	*	-	-
50-54	*	-	-	*	-	-	*	4	6%	*	-	-	*	2	6%	*	-	-
55-64	*	-	-	*	-	-	*	4	6%	*	-	-	*	3	9%	*	-	-
65+	*	-	-	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	1	-	43.1	63	-	19.5	8	-	4.2	34	-	*	1	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 9

Transmission Category ^c	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	1	100%	24	55%	4	80%	13	46%	1	100%
Injection drug use (IDU)	-	-	-	-	1	2%	-	-	4	14%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	2%	-	-	3	11%	-	-
Heterosexual contact	-	-	-	-	4	9%	-	-	2	7%	-	-
Other/unknown	-	-	-	-	14	32%	1	20%	6	21%	-	-
Subtotal	-	-	1	100%	44	100%	5	100%	28	100%	1	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	19	100%	3	100%	6	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	19	100%	3	100%	6	100%	-	-
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-		1		63		8		34		1	

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 9

2024 diagnosis of HIV infection													
Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	
Male-to-male sexual contact only	-	-	1	100%	17	27%	4	50%	10	29%	-	-	
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	1	3%	-	-	
Heterosexual contact only	-	-	-	-	37	59%	4	50%	13	38%	-	-	
Male-to-male sexual contact & IDU	-	-	-	-	1	2%	-	-	2	6%	-	-	
IDU & Heterosexual contact	-	-	-	-	1	2%	-	-	3	9%	-	-	
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	7	11%	-	-	3	9%	1	100%	
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	3%	-	-	
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-	
Other/unknown	-	-	-	-	-	-	-	-	1	3%	-	-	
Total	-	-	1	-	63	-	8	-	34	-	1	-	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
Asterisk (*) indicates rate not calculated. for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 9

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2020			2021			2022			2023			2024				
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex																	
Males	10.5	53	87%	13.9	71	81%	12.6	64	75%	14.7	75	68%	15.4	79	74%	2,979	81%
Females	1.5	8	13%	3.2	17	19%	4.0	21	25%	6.6	35	32%	5.2	28	26%	682	19%
Age at diagnosis (years)																	
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	32	1%
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	11	<1%
15-19	*	2	3%	*	4	5%	9.0	6	7%	10.3	7	6%	11.7	8	7%	174	5%
20-24	11.9	8	13%	26.5	18	20%	21.5	15	18%	16.4	11	10%	17.7	12	11%	504	14%
25-29	20.2	14	23%	22.4	15	17%	24.0	16	19%	28.7	19	17%	23.7	16	15%	639	17%
30-34	18.1	12	20%	14.8	10	11%	17.5	12	14%	33.2	23	21%	31.2	22	21%	603	16%
35-39	17.6	11	18%	10.9	7	8%	15.6	10	12%	23.1	15	14%	13.5	9	8%	538	15%
40-44	*	3	5%	14.9	9	10%	16.2	10	12%	22.2	14	13%	23.4	15	14%	441	12%
45-49	8.5	5	8%	*	4	5%	*	4	5%	16.0	9	8%	19.1	11	10%	319	9%
50-54	*	2	3%	10.9	7	8%	*	3	4%	12.8	8	7%	9.8	6	6%	192	5%
55-64	*	4	7%	6.4	9	10%	4.4	6	7%	*	3	3%	5.3	7	7%	151	4%
65+	*	-	-	2.6	5	6%	*	3	4%	*	1	1%	*	1	1%	57	2%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander	*	-	-	*	-	-	*	2	2%	*	-	-	*	1	1%	17	<1%
Black/African American	18.5	26	43%	28.8	41	47%	28.8	41	48%	32.7	47	43%	43.1	63	59%	1,535	42%
Hispanic/Latino	*	4	7%	*	1	1%	17.4	6	7%	20.8	8	7%	19.5	8	7%	91	2%
White	3.3	27	44%	5.3	43	49%	4.3	35	41%	6.5	52	47%	4.2	34	32%	1,903	52%
Multi-Race	*	4	7%	*	3	3%	*	1	1%	*	3	3%	*	1	1%	110	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	<1%
Race/Ethnicity ^b and Sex																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	1	1%	*	-	-	*	1	1%	15	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	2	<1%
Black/African American Males	34.8	23	38%	55.1	37	42%	47.6	32	38%	44.1	30	27%	63.5	44	41%	1,144	31%
Black/African American Females	*	3	5%	*	4	5%	12.0	9	11%	22.4	17	15%	24.7	19	18%	391	11%
Hispanic/Latino Males	*	4	7%	*	-	-	33.9	6	7%	30.3	6	5%	23.6	5	5%	80	2%
Hispanic/Latina Females	*	-	-	*	1	1%	*	-	-	*	2	2%	*	3	3%	11	<1%
White Males	5.5	22	36%	7.7	31	35%	6.0	24	28%	9.6	38	35%	7.1	28	26%	1,648	45%
White Females	1.2	5	8%	2.9	12	14%	2.7	11	13%	3.4	14	13%	1.5	6	6%	255	7%
Multi-Race Males	*	4	7%	*	3	3%	*	1	1%	*	1	1%	*	1	1%	89	2%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	2	2%	*	-	-	21	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	<1%
Total	5.9	61		8.4	88		8.2	85		10.6	110		10.2	107		3,661	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 9

Diagnoses of HIV Infection by Year											
Transmission Category ^a	2020		2021		2022		2023		2024		Cumulative Diagnoses
	No.	%	No.	%	No.	%	No.	%	No.	%	No. %
Male adult or adolescent											
Male-to-male sexual contact	27	51%	44	62%	36	56%	36	48%	43	54%	2,047 69%
Injection drug use (IDU)	-	-	3	4%	1	2%	3	4%	5	6%	108 4%
Male-to-male sexual contact and IDU	4	8%	6	8%	1	2%	3	4%	4	5%	176 6%
Heterosexual contact	2	4%	3	4%	5	8%	2	3%	6	8%	146 5%
Other/unknown	20	38%	15	21%	21	33%	31	41%	21	27%	480 16%
Subtotal	53	100%	71	100%	64	100%	75	100%	79	100%	2,957 100%
Female adult or adolescent											
Injection drug use	1	13%	5	29%	1	5%	6	17%	-	-	99 15%
Heterosexual contact	4	50%	10	59%	13	62%	25	71%	28	100%	514 77%
Other/unknown	3	38%	2	12%	7	33%	4	11%	-	-	53 8%
Subtotal	8	100%	17	100%	21	100%	35	100%	28	100%	666 100%
Child (<13 years at diagnosis)											
Perinatal	-	-	-	-	-	-	-	-	-	-	29 76%
Other/unknown	-	-	-	-	-	-	-	-	-	-	9 24%
Subtotal	-	-	-	-	-	-	-	-	-	-	38 100%
Total	61		88		85		110		107		3,661

Trends in reported diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 9

Diagnoses of HIV Infection by Year											
Exposure Category ^b	2020		2021		2022		2023		2024		Cumulative Diagnoses
	No.	%	No.	%	No.	%	No.	%	No.	%	No. %
Male-to-male sexual contact only	21	34%	37	42%	30	35%	31	28%	32	30%	1,532 42%
Injection drug use (IDU) only	-	-	-	-	-	-	3	3%	1	1%	40 1%
Heterosexual contact only	9	15%	17	19%	25	29%	39	35%	54	50%	871 24%
Male-to-male sexual contact & IDU	2	3%	6	7%	1	1%	2	2%	3	3%	97 3%
IDU & Heterosexual contact	1	2%	8	9%	2	2%	6	5%	4	4%	166 5%
Male-to-male sexual contact & Heterosexual contact	6	10%	7	8%	6	7%	5	5%	11	10%	515 14%
Male-to-male sexual contact & IDU & Heterosexual contact	2	3%	-	-	-	-	1	1%	1	1%	79 2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	32 1%
Other/unknown	20	33%	13	15%	21	25%	23	21%	1	1%	329 9%
Total	61		88		85		110		107		3,661

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 9

Characteristic	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex						
Males	42	81%	20	87%	22	76%
Females	10	19%	3	13%	7	24%
Age at AIDS diagnosis (years)						
0-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	1	2%	-	-	1	3%
20-24	2	4%	1	4%	1	3%
25-29	7	13%	5	22%	2	7%
30-34	5	10%	2	9%	3	10%
35-39	10	19%	4	17%	6	21%
40-44	9	17%	4	17%	5	17%
45-49	8	15%	3	13%	5	17%
50-54	3	6%	-	-	3	10%
55-64	4	8%	1	4%	3	10%
65+	3	6%	3	13%	-	-
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African American	27	52%	10	43%	17	59%
Hispanic/Latino	3	6%	1	4%	2	7%
White	20	38%	11	48%	9	31%
Multi-Race	2	4%	1	4%	1	3%
Race/Ethnicity^b and Sex						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African American Males	19	37%	7	30%	12	41%
Black/African American Females	8	15%	3	13%	5	17%
Hispanic/Latino Males	2	4%	1	4%	1	3%
Hispanic/Latina Females	1	2%	-	-	1	3%
White Males	19	37%	11	48%	8	28%
White Females	1	2%	-	-	1	3%
Multi-Race Males	2	4%	1	4%	1	3%
Multi-Race Females	-	-	-	-	-	-
Total	52		23		29	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 9

Transmission Category ^b	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	21	50%	11	55%	10	45%
Injection drug use (IDU)	2	5%	1	5%	1	5%
Male-to-male sexual contact and IDU	2	5%	1	5%	1	5%
Heterosexual contact	3	7%	2	10%	1	5%
Other/unknown	14	33%	5	25%	9	41%
Subtotal	42	100%	20	100%	22	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	9	90%	2	67%	7	100%
Other/unknown	1	10%	1	33%	-	-
Subtotal	10	100%	3	100%	7	100%
Child (<13 years at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	52		23		29	

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 9

Exposure Category ^c	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	12	23%	6	26%	6	21%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	24	46%	7	30%	17	59%
Male-to-male sexual contact & IDU	2	4%	1	4%	1	3%
IDU & Heterosexual contact	2	4%	1	4%	1	3%
Male-to-male sexual contact & Heterosexual contact	9	17%	5	22%	4	14%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	3	6%	3	13%	-	-
Total	52		23		29	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 9

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	32	89%	27	75%	33	75%	35	73%	42	81%	1,885	83%
Females	4	11%	9	25%	11	25%	13	27%	10	19%	374	17%
Age at AIDS diagnosis (years)												
0-9	-	-	-	-	-	-	-	-	-	-	14	1%
10-14	-	-	-	-	-	-	-	-	-	-	7	<1%
15-19	1	3%	-	-	1	2%	-	-	1	2%	31	1%
20-24	5	14%	4	11%	3	7%	2	4%	2	4%	131	6%
25-29	7	19%	5	14%	7	16%	6	13%	7	13%	298	13%
30-34	5	14%	5	14%	5	11%	12	25%	5	10%	384	17%
35-39	7	19%	4	11%	6	14%	8	17%	10	19%	415	18%
40-44	3	8%	5	14%	5	11%	7	15%	9	17%	373	17%
45-49	2	6%	1	3%	8	18%	4	8%	8	15%	276	12%
50-54	3	8%	1	3%	1	2%	3	6%	3	6%	149	7%
55-64	3	8%	6	17%	4	9%	4	8%	4	8%	126	6%
65+	-	-	5	14%	4	9%	2	4%	3	6%	55	2%
Race/Ethnicity ^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	8	<1%
Black/African American	16	44%	13	36%	22	50%	20	42%	27	52%	877	39%
Hispanic/Latino	1	3%	-	-	2	5%	4	8%	3	6%	45	2%
White	17	47%	22	61%	18	41%	23	48%	20	38%	1,277	57%
Multi-Race	2	6%	1	3%	2	5%	1	2%	2	4%	52	2%
Race/Ethnicity ^a and Sex												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	6	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Black/African American Males	14	39%	10	28%	20	45%	13	27%	19	37%	665	29%
Black/African American Females	2	6%	3	8%	2	5%	7	15%	8	15%	212	9%
Hispanic/Latino Males	1	3%	-	-	2	5%	3	6%	2	4%	40	2%
Hispanic/Latina Females	-	-	-	-	-	-	1	2%	1	2%	5	<1%
White Males	15	42%	16	44%	9	20%	18	38%	19	37%	1,130	50%
White Females	2	6%	6	17%	9	20%	5	10%	1	2%	147	7%
Multi-Race Males	2	6%	1	3%	2	5%	1	2%	2	4%	44	2%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	8	<1%
Total	36		36		44		48		52		2,259	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 9

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	15	48%	18	67%	13	39%	18	51%	21	50%	1,297	69%
Injection drug use (IDU)	1	3%	-	-	3	9%	1	3%	2	5%	78	4%
Male-to-male sexual contact and IDU	1	3%	4	15%	1	3%	4	11%	2	5%	124	7%
Heterosexual contact	2	6%	1	4%	4	12%	-	-	3	7%	111	6%
Other/unknown	12	39%	4	15%	12	36%	12	34%	14	33%	258	14%
Subtotal	31	100%	27	100%	33	100%	35	100%	42	100%	1,868	100%
Female adult or adolescent												
Injection drug use	-	-	1	11%	4	36%	4	31%	-	-	63	17%
Heterosexual contact	4	100%	6	67%	6	55%	7	54%	9	90%	275	76%
Other/unknown	-	-	2	22%	1	9%	2	15%	1	10%	26	7%
Subtotal	4	100%	9	100%	11	100%	13	100%	10	100%	364	100%
Child (<13 years at diagnosis)												
Perinatal	1	100%	-	-	-	-	-	-	-	-	19	70%
Other/unknown	-	-	-	-	-	-	-	-	-	-	8	30%
Subtotal	1	100%	-	-	-	-	-	-	-	-	27	100%
Total	36		36		44		48		52		2,259	

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 9

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	12	33%	15	42%	12	27%	13	27%	12	23%	942	42%
Injection drug use (IDU) only	-	-	-	-	1	2%	2	4%	-	-	26	1%
Heterosexual contact only	8	22%	8	22%	15	34%	14	29%	24	46%	525	23%
Male-to-male sexual contact & IDU	-	-	4	11%	1	2%	-	-	2	4%	64	3%
IDU & Heterosexual contact	1	3%	1	3%	6	14%	3	6%	2	4%	114	5%
Male-to-male sexual contact & Heterosexual contact	3	8%	3	8%	1	2%	5	10%	9	17%	355	16%
Male-to-male sexual contact & IDU & Heterosexual contact	1	3%	-	-	-	-	4	8%	-	-	60	3%
Perinatal exposure	1	3%	-	-	-	-	-	-	-	-	20	1%
Other/unknown	10	28%	5	14%	8	18%	7	15%	3	6%	153	7%
Total	36		36		44		48		52		2,259	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 9

Counties: Clark, Darke, Greene, Miami, Montgomery, Preble

Reported persons living with diagnosed HIV infection in 2024 by current disease status and selected characteristics, HIV Prevention Planning Region 9

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2024			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex							
Males	362.1	1,861	77%	974	74%	887	81%
Females	102.4	548	23%	339	26%	209	19%
Age at end of year (years)							
0-9	*	1	<1%	1	<1%	-	-
10-14	*	1	<1%	1	<1%	-	-
15-19	30.8	21	1%	19	1%	2	<1%
20-24	96.1	65	3%	58	4%	7	1%
25-29	238.2	161	7%	129	10%	32	3%
30-34	407.6	287	12%	213	16%	74	7%
35-39	383.9	256	11%	162	12%	94	9%
40-44	398.9	256	11%	144	11%	112	10%
45-49	414.1	238	10%	127	10%	111	10%
50-54	399.3	244	10%	116	9%	128	12%
55-64	414.0	545	23%	208	16%	337	31%
65+	161.4	334	14%	135	10%	199	18%
Race/Ethnicity ^b							
American Indian/Alaska Native	*	2	<1%	2	<1%	-	-
Asian/Pacific Islander	67.3	17	1%	13	1%	4	<1%
Black/African American	766.6	1,120	46%	633	48%	487	44%
Hispanic/Latino	297.3	122	5%	71	5%	51	5%
White	127.9	1,028	43%	523	40%	505	46%
Multi-Race	377.7	116	5%	67	5%	49	4%
Unknown	*	4	<1%	4	<1%	-	-
Race/Ethnicity ^b and Sex							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	2	<1%	2	<1%	-	-
Asian/Pacific Islander Males	124.8	15	1%	12	1%	3	<1%
Asian/Pacific Islander Females	*	2	<1%	1	<1%	1	<1%
Black/African American Males	1,140.0	790	33%	427	33%	363	33%
Black/African American Females	429.6	330	14%	206	16%	124	11%
Hispanic/Latino Males	510.0	108	4%	61	5%	47	4%
Hispanic/Latina Females	70.5	14	1%	10	1%	4	<1%
White Males	214.7	848	35%	417	32%	431	39%
White Females	44.1	180	7%	106	8%	74	7%
Multi-Race Males	639.1	97	4%	54	4%	43	4%
Multi-Race Females	122.3	19	1%	13	1%	6	1%
Unknown	*	4	<1%	4	<1%	-	-
Total	229.6	2,409		1,313		1,096	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by current disease status and transmission category, HIV Prevention Planning Region 9

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,244	67%	651	67%	593	67%
Injection drug use (IDU)	49	3%	29	3%	20	2%
Male-to-male sexual contact and IDU	93	5%	42	4%	51	6%
Heterosexual contact	102	6%	37	4%	65	7%
Other/unknown	361	20%	206	21%	155	18%
Subtotal	1,849	100%	965	100%	884	100%
Female adult or adolescent						
Injection drug use	67	13%	36	11%	31	15%
Heterosexual contact	405	76%	243	75%	162	78%
Other/unknown	63	12%	47	14%	16	8%
Subtotal	535	100%	326	100%	209	100%
Child (<13 years at diagnosis)						
Perinatal	16	64%	16	73%	-	-
Other/unknown	9	36%	6	27%	3	100%
Subtotal	25	100%	22	100%	3	100%
Total	2,409		1,313		1,096	

Reported persons living with diagnosed HIV infection in 2024 by current disease status and exposure category, HIV Prevention Planning Region 9

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	952	40%	521	40%	431	39%
Injection drug use (IDU) only	11	<1%	7	1%	4	<1%
Heterosexual contact only	669	28%	358	27%	311	28%
Male-to-male sexual contact & IDU	52	2%	26	2%	26	2%
IDU & Heterosexual contact	105	4%	58	4%	47	4%
Male-to-male sexual contact & Heterosexual contact	292	12%	130	10%	162	15%
Male-to-male sexual contact & IDU & Heterosexual contact	41	2%	16	1%	25	2%
Perinatal exposure	20	1%	18	1%	2	<1%
Other/unknown	267	11%	179	14%	88	8%
Total	2,409		1,313		1,096	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and age, HIV Prevention Planning Region 9

Living with diagnosed HIV infection in 2024																				
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	-	-
10-14	*	-	-	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	132.8	15	1%	*	2	2%	*	3	<1%	*	1	1%	-	-
20-24	*	-	-	*	1	6%	429.2	46	4%	133.8	5	4%	20.9	10	1%	*	3	3%	-	-
25-29	*	-	-	*	3	18%	904.5	94	8%	471.0	16	13%	81.2	40	4%	321.4	8	7%	-	-
30-34	*	-	-	*	4	24%	1,301.7	151	13%	659.7	22	18%	183.5	94	9%	831.6	16	14%	-	-
35-39	*	1	50%	*	-	-	1,403.1	134	12%	758.1	22	18%	170.5	86	8%	917.4	13	11%	-	-
40-44	*	-	-	*	3	18%	1,414.5	118	11%	400.1	12	10%	217.9	108	11%	1,331.0	15	13%	-	-
45-49	*	-	-	*	2	12%	1,395.6	104	9%	596.9	13	11%	228.0	103	10%	1,588.9	16	14%	-	-
50-54	*	-	-	*	1	6%	1,437.4	114	10%	367.6	7	6%	221.7	108	11%	1,631.1	13	11%	1	25%
55-64	*	1	50%	*	3	18%	1,402.3	223	20%	517.1	15	12%	258.9	281	27%	1,537.3	20	17%	2	50%
65+	*	-	-	*	-	-	544.3	120	11%	261.7	7	6%	110.4	195	19%	647.4	11	9%	1	25%
Total	*	2		67.3	17		766.6	1,120		297.3	122		127.9	1,028		377.7	116		4	

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 9

Living with diagnosed HIV infection in 2024														
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	8	53%	490	63%	73	68%	606	72%	67	70%	-	-
Injection drug use (IDU)	-	-	-	-	13	2%	2	2%	29	3%	5	5%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	22	3%	4	4%	57	7%	10	10%	-	-
Heterosexual contact	-	-	-	-	70	9%	6	6%	24	3%	2	2%	-	-
Other/unknown	-	-	7	47%	189	24%	22	21%	128	15%	12	13%	3	100%
Subtotal	-	-	15	100%	784	100%	107	100%	844	100%	96	100%	3	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	15	5%	1	7%	51	28%	-	-	-	-
Heterosexual contact	1	50%	1	50%	262	82%	12	86%	112	63%	17	89%	-	-
Other/unknown	1	50%	1	50%	41	13%	1	7%	16	9%	2	11%	1	100%
Subtotal	2	100%	2	100%	318	100%	14	100%	179	100%	19	100%	1	100%
Child (<13 years at diagnosis)														
Perinatal	-	-	-	-	12	67%	1	100%	3	60%	-	-	-	-
Other/unknown	-	-	-	-	6	33%	-	-	2	40%	1	100%	-	-
Subtotal	-	-	-	-	18	100%	1	100%	5	100%	1	100%	-	-
Total	2		17		1,120		122		1,028		116		4	

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 9

Living with diagnosed HIV infection in 2024														
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
Exposure Category ^d	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	7	41%	357	32%	58	48%	473	46%	57	49%	-	-
Injection drug use (IDU) only	-	-	-	-	3	0%	1	1%	7	1%	-	-	-	-
Heterosexual contact only	1	50%	3	18%	428	38%	23	19%	189	18%	25	22%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	9	1%	3	2%	33	3%	7	6%	-	-
IDU & Heterosexual contact	-	-	-	-	25	2%	2	2%	73	7%	5	4%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	1	6%	133	12%	15	12%	133	13%	10	9%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	13	1%	1	1%	24	2%	3	3%	-	-
Perinatal exposure	-	-	-	-	15	1%	2	2%	3	0%	-	-	-	-
Other/unknown	1	50%	6	35%	137	12%	17	14%	93	9%	9	8%	4	100%
Total	2		17		1,120		122		1,028		116		4	

Notes:
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 9, 2020-2024

Living with Diagnosed HIV Infection															
Characteristic	2020			2021			2022			2023			2024		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	326.3	1,643	79%	328.5	1,675	79%	335.3	1,709	79%	345.1	1,758	78%	362.1	1,861	77%
Females	81.3	432	21%	84.8	451	21%	86.6	459	21%	93.9	499	22%	102.4	548	23%
Age at end of year (years)															
0-9	*	-	-	*	1	<1%	*	1	<1%	*	2	<1%	*	1	<1%
10-14	9.4	6	<1%	7.5	5	<1%	7.7	5	<1%	*	1	<1%	*	1	<1%
15-19	15.1	10	<1%	19.1	13	1%	13.5	9	<1%	22.1	15	1%	30.8	21	1%
20-24	99.5	67	3%	108.8	74	3%	104.5	73	3%	101.6	68	3%	96.1	65	3%
25-29	264.6	183	9%	241.6	162	8%	238.8	159	7%	244.6	162	7%	238.2	161	7%
30-34	314.0	208	10%	339.4	230	11%	335.5	230	11%	374.1	259	11%	407.6	287	12%
35-39	312.1	195	9%	318.2	204	10%	333.0	214	10%	374.2	243	11%	383.9	256	11%
40-44	330.2	191	9%	328.7	199	9%	338.9	209	10%	360.4	227	10%	398.9	256	11%
45-49	359.6	211	10%	342.6	195	9%	367.5	205	9%	373.7	210	9%	414.1	238	10%
50-54	414.9	261	13%	405.9	261	12%	384.4	243	11%	392.5	246	11%	399.3	244	10%
55-64	383.7	537	26%	391.7	549	26%	404.0	552	25%	398.6	534	24%	414.0	545	23%
65+	105.2	206	10%	119.7	233	11%	134.3	268	12%	143.3	290	13%	161.4	334	14%
Race/Ethnicity ^b															
American Indian/Alaska Native	*	1	<1%	*	2	<1%	*	1	<1%	*	1	<1%	*	2	<1%
Asian/Pacific Islander	64.5	14	1%	73.5	16	1%	85.1	19	1%	74.6	17	1%	67.3	17	1%
Black/African American	668.6	940	45%	687.6	978	46%	687.7	979	45%	713.3	1,026	45%	766.6	1,120	46%
Hispanic/Latino	242.1	77	4%	222.1	74	3%	246.3	85	4%	280.6	108	5%	297.3	122	5%
White	115.5	937	45%	115.8	942	44%	120.1	972	45%	123.3	991	44%	127.9	1,028	43%
Multi-Race	372.8	102	5%	388.2	110	5%	371.6	108	5%	369.5	110	5%	377.7	116	5%
Unknown	*	4	<1%	*	4	<1%	*	4	<1%	*	4	<1%	*	4	<1%
Race/Ethnicity ^b and Sex															
American Indian/Alaska Native Males	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	2	<1%
Asian/Pacific Islander Males	129.1	13	1%	146.9	15	1%	161.6	17	1%	140.1	15	1%	124.8	15	1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	2	<1%	*	2	<1%	*	2	<1%
Black/African American Males	1,023.0	676	33%	1,056.0	709	33%	1,059.8	713	33%	1,077.7	733	32%	1,140.0	790	33%
Black/African American Females	354.3	264	13%	358.3	269	13%	354.3	266	12%	386.4	293	13%	429.6	330	14%
Hispanic/Latino Males	449.5	73	4%	392.6	67	3%	440.5	78	4%	490.0	97	4%	510.0	108	4%
Hispanic/Latina Females	*	4	<1%	43.1	7	<1%	41.7	7	<1%	58.8	11	<1%	70.5	14	1%
White Males	201.0	797	38%	197.6	791	37%	203.1	810	37%	207.3	819	36%	214.7	848	35%
White Females	33.8	140	7%	36.5	151	7%	39.5	162	7%	42.1	172	8%	44.1	180	7%
Multi-Race Males	600.0	81	4%	638.7	89	4%	614.2	88	4%	617.5	91	4%	639.1	97	4%
Multi-Race Females	151.5	21	1%	145.8	21	1%	135.7	20	1%	126.4	19	1%	122.3	19	1%
Unknown	*	4	<1%	*	4	<1%	*	4	<1%	*	4	<1%	*	4	<1%
Total	200.5	2,075		204.1	2,126		208.6	2,168		216.8	2,257		229.6	2,409	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 9, 2020-2024

Living with Diagnosed HIV Infection										
Transmission Category ^a	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	1,181	72%	1,184	71%	1,179	69%	1,191	68%	1,244	67%
Injection drug use (IDU)	37	2%	41	2%	47	3%	45	3%	49	3%
Male-to-male sexual contact and IDU	82	5%	86	5%	90	5%	93	5%	93	5%
Heterosexual contact	81	5%	81	5%	84	5%	90	5%	102	6%
Other/unknown	251	15%	270	16%	297	18%	327	19%	361	20%
Subtotal	1,632	100%	1,662	100%	1,697	100%	1,746	100%	1,849	100%
Female adult or adolescent										
Injection drug use	53	13%	58	13%	60	13%	65	13%	67	13%
Heterosexual contact	340	80%	347	79%	346	77%	370	76%	405	76%
Other/unknown	30	7%	36	8%	44	10%	55	11%	63	12%
Subtotal	423	100%	441	100%	450	100%	490	100%	535	100%
Child (<13 years at diagnosis)										
Perinatal	13	65%	16	70%	14	67%	14	67%	16	64%
Other/unknown	7	35%	7	30%	7	33%	7	33%	9	36%
Subtotal	20	100%	23	100%	21	100%	21	100%	25	100%
Total	2,075		2,126		2,168		2,257		2,409	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 9, 2020-2024

Living with Diagnosed HIV Infection										
Exposure Category ^b	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	897	43%	894	42%	899	41%	915	41%	952	40%
Injection drug use (IDU) only	8	<1%	7	<1%	8	<1%	10	<1%	11	<1%
Heterosexual contact only	526	25%	540	25%	552	25%	591	26%	669	28%
Male-to-male sexual contact & IDU	42	2%	46	2%	51	2%	49	2%	52	2%
IDU & Heterosexual contact	82	4%	92	4%	99	5%	100	4%	105	4%
Male-to-male sexual contact & Heterosexual contact	284	14%	290	14%	280	13%	276	12%	292	12%
Male-to-male sexual contact & IDU & Heterosexual contact	40	2%	40	2%	39	2%	44	2%	41	2%
Perinatal exposure	15	1%	18	1%	16	1%	17	1%	20	1%
Other/unknown	181	9%	199	9%	224	10%	255	11%	267	11%
Total	2,075		2,126		2,168		2,257		2,409	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.