

Persons Living with Diagnosed HIV Infection Reported in Allen County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Allen County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	211.7	109	69%	54	65%	55	74%
Females	95.0	48	31%	29	35%	19	26%
Age at end of year							
<13	*	1	1%	1	1%	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	1	1%	1	1%	-	-
25-29	135.2	9	6%	8	10%	1	1%
30-34	332.8	20	13%	11	13%	9	12%
35-39	219.6	13	8%	9	11%	4	5%
40-44	387.1	23	15%	13	16%	10	14%
45-49	208.4	12	8%	8	10%	4	5%
50-54	355.8	21	13%	6	7%	15	20%
55-64	323.6	44	28%	21	25%	23	31%
65+	68.9	13	8%	5	6%	8	11%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	1	1%	-	-	1	1%
Black/African-American	521.9	66	42%	34	41%	32	43%
Hispanic/Latinx	249.8	9	6%	7	8%	2	3%
White	89.4	73	46%	38	46%	35	47%
Multi-Race	267.8	8	5%	4	5%	4	5%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	-	-	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African-American Males	604.7	40	25%	21	25%	19	26%
Black/African-American Females	431.2	26	17%	13	16%	13	18%
Hispanic/Latino Males	349.5	7	4%	5	6%	2	3%
Hispanic/Latina Females	*	2	1%	2	2%	-	-
White Males	137.2	56	36%	25	30%	31	42%
White Females	41.6	17	11%	13	16%	4	5%
Multi-Race Males	326.4	5	3%	3	4%	2	3%
Multi-Race Females	*	3	2%	1	1%	2	3%
Unknown	*	-	-	-	-	-	-
Total	154.0	157		83		74	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Allen County

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	65	61%	30	57%	35	65%
Injection drug use (IDU)	7	7%	3	6%	4	7%
Male-to-male sexual contact and IDU	10	9%	5	9%	5	9%
Heterosexual contact	12	11%	7	13%	5	9%
Other/unknown	13	12%	8	15%	5	9%
Subtotal	107	100%	53	100%	54	100%
Female adult or adolescent						
Injection drug use	6	13%	4	14%	2	11%
Heterosexual contact	39	83%	24	83%	15	83%
Other/unknown	2	4%	1	3%	1	6%
Subtotal	47	100%	29	100%	18	100%
Child (<13 yrs at diagnosis)						
Perinatal	3	100%	1	100%	2	100%
Other/unknown	-	-	-	-	-	-
Subtotal	3	100%	1	100%	2	100%
Total	157		83		74	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Allen County

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	57	36%	26	31%	31	42%
Injection drug use (IDU) only	6	4%	3	4%	3	4%
Heterosexual contact only	51	32%	31	37%	20	27%
Male-to-male sexual contact & IDU	9	6%	5	6%	4	5%
IDU & Heterosexual contact	7	4%	4	5%	3	4%
Male-to-male sexual contact & Heterosexual contact	8	5%	4	5%	4	5%
Male-to-male sexual contact & IDU & Heterosexual contact	1	1%	-	-	1	1%
Perinatal exposure	3	2%	1	1%	2	3%
Other/unknown	15	10%	9	11%	6	8%
Total	157		83		74	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Allen County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	1	2%	*	-	-	*	-	-	*	<5	-	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	<5	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	<5	-	-	-	-
20-24	*	-	-	*	-	-	*	-	-	*	-	-	*	1	1%	*	<5	-	-	-	-
25-29	*	-	-	*	-	-	500.5	5	8%	*	-	-	*	4	5%	*	<5	-	-	-	-
30-34	*	-	-	*	-	-	1,267.4	10	15%	*	2	22%	146.2	7	10%	*	<5	-	-	-	-
35-39	*	-	-	*	-	-	*	4	6%	*	1	11%	149.5	7	10%	*	<5	-	-	-	-
40-44	*	-	-	*	-	-	1,250.0	10	15%	*	4	44%	148.4	7	10%	*	<5	-	-	-	-
45-49	*	-	-	*	-	-	859.6	6	9%	*	-	-	*	4	5%	*	<5	-	-	-	-
50-54	*	-	-	*	-	-	1,771.1	13	20%	*	-	-	163.5	8	11%	*	<5	-	-	-	-
55-64	*	-	-	*	1	100%	1,030.2	15	23%	*	2	22%	205.8	24	33%	*	<5	-	-	-	-
65+	*	-	-	*	-	-	*	2	3%	*	-	-	65.6	11	15%	*	<5	-	-	-	-
Total	*	-	-	*	1	-	521.9	66	-	249.8	9	-	89.4	73	-	267.8	8	-	-	-	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Allen County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	24	62%	2	33%	35	63%	4	80%	-	-
Injection drug use (IDU)	-	-	-	-	4	10%	-	-	3	5%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	4	10%	-	-	6	11%	-	-	-	-
Heterosexual contact	-	-	1	100%	3	8%	3	50%	5	9%	-	-	-	-
Other/unknown	-	-	-	-	4	10%	1	17%	7	13%	1	20%	-	-
Subtotal	-	-	1	100%	39	100%	6	100%	56	100%	5	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	3	12%	-	-	3	19%	-	-	-	-
Heterosexual contact	-	-	-	-	21	81%	2	100%	13	81%	3	100%	-	-
Other/unknown	-	-	-	-	2	8%	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	26	100%	2	100%	16	100%	3	100%	-	-
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	1	100%	1	100%	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	1	100%	1	100%	-	-	-	-
Total	-	-	1	-	66	-	9	-	73	-	8	-	-	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Allen County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	21	32%	2	22%	31	42%	3	38%	-	-
Injection drug use (IDU) only	-	-	-	-	4	6%	-	-	2	3%	-	-	-	-
Heterosexual contact only	-	-	1	100%	24	36%	5	56%	18	25%	3	38%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	4	6%	-	-	5	7%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	3	5%	-	-	4	5%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	3	5%	-	-	4	5%	1	13%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	1%	-	-	-	-
Perinatal exposure	-	-	-	-	1	2%	1	11%	1	1%	-	-	-	-
Other/unknown	-	-	-	-	6	9%	1	11%	7	10%	1	13%	-	-
Total	-	-	1	-	66	-	9	-	73	-	8	-	-	

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Allen County, 2016-2020

Characteristic	Living with Diagnosed HIV Infection														
	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	204.2	107	69%	205.4	107	69%	197.0	102	67%	211.0	109	69%	211.7	109	69%
Females	95.4	49	31%	93.9	48	31%	100.2	51	33%	94.7	48	31%	95.0	48	31%
Age at end of year															
<13	*	-	-	*	-	-	*	1	1%	*	-	-	*	1	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	1%	*	-	-	*	-	-	*	-	-	*	-	-
20-24	82.0	6	4%	71.1	5	3%	*	3	2%	*	2	1%	*	1	1%
25-29	249.1	17	11%	215.4	15	10%	172.8	12	8%	116.4	8	5%	135.2	9	6%
30-34	255.7	15	10%	314.5	18	12%	280.6	16	10%	254.3	15	10%	332.8	20	13%
35-39	336.9	21	13%	287.5	18	12%	340.2	21	14%	331.5	20	13%	219.6	13	8%
40-44	209.8	12	8%	279.3	16	10%	258.0	15	10%	371.8	22	14%	387.1	23	15%
45-49	314.0	19	12%	313.1	19	12%	271.1	16	10%	243.2	14	9%	208.4	12	8%
50-54	317.6	22	14%	271.8	18	12%	284.0	18	12%	280.6	17	11%	355.8	21	13%
55-64	245.9	35	22%	269.6	38	25%	285.2	40	26%	339.0	47	30%	323.6	44	28%
65+	46.0	8	5%	45.0	8	5%	60.7	11	7%	64.8	12	8%	68.9	13	8%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Black/African-American	462.8	59	38%	471.1	60	39%	465.0	59	39%	502.8	63	40%	521.9	66	42%
Hispanic/Latinx	358.3	11	7%	376.1	12	8%	338.7	11	7%	292.4	10	6%	249.8	9	6%
White	94.0	79	51%	92.3	77	50%	90.5	75	49%	92.1	76	48%	89.4	73	46%
Multi-Race	218.0	6	4%	176.3	5	3%	243.0	7	5%	241.3	7	4%	267.8	8	5%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African-American Males	528.5	35	22%	541.4	36	23%	468.8	31	20%	534.2	35	22%	604.7	40	25%
Black/African-American Females	391.8	24	15%	394.3	24	15%	460.9	28	18%	468.5	28	18%	431.2	26	17%
Hispanic/Latino Males	404.9	7	4%	449.7	8	5%	387.6	7	5%	367.6	7	4%	349.5	7	4%
Hispanic/Latina Females	*	4	3%	*	4	3%	*	4	3%	*	3	2%	*	2	1%
White Males	142.4	60	38%	141.3	59	38%	142.3	59	39%	147.9	61	39%	137.2	56	36%
White Females	45.3	19	12%	43.2	18	12%	38.7	16	10%	36.4	15	10%	41.6	17	11%
Multi-Race Males	*	4	3%	*	3	2%	*	4	3%	339.4	5	3%	326.4	5	3%
Multi-Race Females	*	2	1%	*	2	1%	*	3	2%	*	2	1%	*	3	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	150.4	156		150.2	155		149.0	153		153.4	157		154.0	157	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Allen County, 2016-2020

Transmission Category ^a	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	70	66%	69	65%	61	61%	63	58%	65	61%
Injection drug use (IDU)	8	8%	8	8%	7	7%	6	6%	7	7%
Male-to-male sexual contact and IDU	4	4%	2	2%	6	6%	9	8%	10	9%
Heterosexual contact	10	9%	11	10%	11	11%	13	12%	12	11%
Other/unknown	14	13%	16	15%	15	15%	17	16%	13	12%
Subtotal	106	100%	106	100%	100	100%	108	100%	107	100%
Female adult or adolescent										
Injection drug use	7	15%	5	11%	7	14%	6	13%	6	13%
Heterosexual contact	40	83%	40	85%	40	80%	38	81%	39	83%
Other/unknown	1	2%	2	4%	3	6%	3	6%	2	4%
Subtotal	48	100%	47	100%	50	100%	47	100%	47	100%
Child (<13 yrs at diagnosis)										
Perinatal	2	100%	2	100%	3	100%	2	100%	3	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	2	100%	2	100%	3	100%	2	100%	3	100%
Total	156		155		153		157		157	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Allen County, 2016-2020

Exposure Category ^b	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	63	40%	63	41%	55	36%	54	34%	57	36%
Injection drug use (IDU) only	5	3%	4	3%	5	3%	6	4%	6	4%
Heterosexual contact only	50	32%	51	33%	51	33%	51	32%	51	32%
Male-to-male sexual contact & IDU	3	2%	1	1%	5	3%	8	5%	9	6%
IDU & Heterosexual contact	10	6%	9	6%	9	6%	6	4%	7	4%
Male-to-male sexual contact & Heterosexual contact	7	4%	6	4%	6	4%	9	6%	8	5%
Male-to-male sexual contact & IDU & Heterosexual contact	1	1%	1	1%	1	1%	1	1%	1	1%
Perinatal exposure	2	1%	2	1%	3	2%	2	1%	3	2%
Other/unknown	15	10%	18	12%	18	12%	20	13%	15	10%
Total	156		155		153		157		157	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.