



Department of
Health

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 2

Counties: Ashland, Crawford, Erie, Huron, Knox, Marion, Richland,
Seneca, Wyandot

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2025

Reported new diagnoses of HIV infection in 2024 by disease status and selected characteristics, HIV Prevention Planning Region 2

Characteristic	2024 diagnosis of HIV infection			Disease Status						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	3.6	10	83%	8	100%	2	67%	-	-	
Females	*	2	17%	-	-	1	33%	1	100%	
Age at diagnosis (years)										
0-9	*	-	-	-	-	-	-	-	-	
10-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	*	2	17%	2	25%	-	-	-	-	
25-29	*	-	-	-	-	-	-	-	-	
30-34	*	4	33%	2	25%	2	67%	-	-	
35-39	*	4	33%	3	38%	1	33%	-	-	
40-44	*	2	17%	1	13%	-	-	1	100%	
45-49	*	-	-	-	-	-	-	-	-	
50-54	*	-	-	-	-	-	-	-	-	
55-64	*	-	-	-	-	-	-	-	-	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b										
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
Black/African American	*	4	33%	2	25%	1	33%	1	100%	
Hispanic/Latino	*	-	-	-	-	-	-	-	-	
White	1.6	8	67%	6	75%	2	67%	-	-	
Multi-Race	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b and Sex										
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
Black/African American Males	*	2	17%	2	25%	-	-	-	-	
Black/African American Females	*	2	17%	-	-	1	33%	1	100%	
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
White Males	3.3	8	67%	6	75%	2	67%	-	-	
White Females	*	-	-	-	-	-	-	-	-	
Multi-Race Males	*	-	-	-	-	-	-	-	-	
Multi-Race Females	*	-	-	-	-	-	-	-	-	
Total	2.2	12		8		3		1		

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by disease status and transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	5	50%	4	50%	1	50%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	10%	1	13%	-	-	-	-
Other/unknown	4	40%	3	38%	1	50%	-	-
Subtotal	10	100%	8	100%	2	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	2	100%	-	-	1	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	2	100%	-	-	1	100%	1	100%
Child (<13 years at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	12		8		3		1	

Reported new diagnoses of HIV infection in 2024 by disease status and exposure category, HIV Prevention Planning Region 2

Exposure Category ^b	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	25%	2	25%	1	33%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	5	42%	3	38%	1	33%	1	100%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	2	17%	2	25%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	2	17%	1	13%	1	33%	-	-
Total	12		8		3		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 2

2024 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
0-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	1	25%	*	-	-	*	1	13%	*	-	-
25-29	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
30-34	*	-	-	*	-	-	*	1	25%	*	-	-	*	3	38%	*	-	-
35-39	*	-	-	*	-	-	*	1	25%	*	-	-	*	3	38%	*	-	-
40-44	*	-	-	*	-	-	*	1	25%	*	-	-	*	1	13%	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	*	4	-	*	-	-	1.6	8	-	*	-	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 2

Transmission Category ^c	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	1	50%	-	-	4	50%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	1	50%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	4	50%	-	-
Subtotal	-	-	-	-	2	100%	-	-	8	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	2	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	2	100%	-	-	-	-	-	-
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	4	-	-	-	8	-	-	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 2

Exposure Category ^d	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-	-	-	3	38%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	3	75%	-	-	2	25%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	1	25%	-	-	1	13%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	25%	-	-
Total	-	-	-	-	4	-	-	-	8	-	-	-

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
Asterisk (*) indicates rate not calculated. for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 2

Diagnoses of HIV Infection by Year																	
Characteristic	2020			2021			2022			2023			2024			Cumulative Diagnoses	
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex																	
Males	5.8	16	94%	3.2	9	75%	3.6	10	83%	5.0	14	78%	3.6	10	83%	618	83%
Females	*	1	6%	*	3	25%	*	2	17%	*	4	22%	*	2	17%	131	17%
Age at diagnosis (years)																	
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	9	1%
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	1%
15-19	*	2	12%	*	-	-	*	-	-	*	-	-	*	-	-	23	3%
20-24	*	2	12%	*	3	25%	*	2	17%	*	3	17%	*	2	17%	91	12%
25-29	14.6	5	29%	*	1	8%	*	2	17%	*	4	22%	*	-	-	126	17%
30-34	*	1	6%	*	2	17%	14.8	5	42%	*	2	11%	*	4	33%	125	17%
35-39	*	-	-	*	2	17%	*	2	17%	*	1	6%	*	4	33%	145	19%
40-44	*	1	6%	*	1	8%	*	-	-	*	2	11%	*	2	17%	83	11%
45-49	*	1	6%	*	1	8%	*	-	-	*	1	6%	*	-	-	58	8%
50-54	*	2	12%	*	-	-	*	1	8%	*	2	11%	*	-	-	31	4%
55-64	*	3	18%	*	1	8%	*	-	-	*	2	11%	*	-	-	42	6%
65+	*	-	-	*	1	8%	*	-	-	*	1	6%	*	-	-	12	2%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Black/African American	*	2	12%	*	4	33%	*	4	33%	*	3	17%	*	4	33%	126	17%
Hispanic/Latino	*	2	12%	*	2	17%	*	-	-	*	3	17%	*	-	-	45	6%
White	2.2	11	65%	1.2	6	50%	1.6	8	67%	2.4	12	67%	1.6	8	67%	556	74%
Multi-Race	*	2	12%	*	-	-	*	-	-	*	-	-	*	-	-	15	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	5	1%
Race/Ethnicity ^b and Sex																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Black/African American Males	*	2	12%	*	2	17%	*	3	25%	*	1	6%	*	2	17%	86	11%
Black/African American Females	*	-	-	*	2	17%	*	1	8%	*	2	11%	*	2	17%	40	5%
Hispanic/Latino Males	*	2	12%	*	2	17%	*	-	-	*	3	17%	*	-	-	41	5%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	1%
White Males	4.1	10	59%	2.0	5	42%	2.8	7	58%	4.1	10	56%	3.3	8	67%	474	63%
White Females	*	1	6%	*	1	8%	*	1	8%	*	2	11%	*	-	-	82	11%
Multi-Race Males	*	2	12%	*	-	-	*	-	-	*	-	-	*	-	-	12	2%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	5	1%
Total	3.1	17		2.2	12		2.2	12		3.2	18		2.2	12		749	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	11	69%	5	56%	8	80%	8	57%	5	50%	405	66%
Injection drug use (IDU)	-	-	1	11%	-	-	-	-	-	-	28	5%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	2	14%	-	-	45	7%
Heterosexual contact	-	-	-	-	-	-	-	-	1	10%	24	4%
Other/unknown	5	31%	3	33%	2	20%	4	29%	4	40%	110	18%
Subtotal	16	100%	9	100%	10	100%	14	100%	10	100%	612	100%
Female adult or adolescent												
Injection drug use	-	-	1	33%	1	50%	1	25%	-	-	18	14%
Heterosexual contact	-	-	2	67%	1	50%	2	50%	2	100%	94	75%
Other/unknown	1	100%	-	-	-	-	1	25%	-	-	14	11%
Subtotal	1	100%	3	100%	2	100%	4	100%	2	100%	126	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	8	73%
Other/unknown	-	-	-	-	-	-	-	-	-	-	3	27%
Subtotal	-	-	-	-	-	-	-	-	-	-	11	100%
Total	17		12		12		18		12		749	

Trends in reported diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 2

Exposure Category ^b	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	8	47%	4	33%	7	58%	7	39%	3	25%	303	40%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	9	1%
Heterosexual contact only	1	6%	4	33%	2	17%	3	17%	5	42%	151	20%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	2	11%	-	-	25	3%
IDU & Heterosexual contact	-	-	2	17%	1	8%	1	6%	-	-	37	5%
Male-to-male sexual contact & Heterosexual contact	3	18%	1	8%	1	8%	1	6%	2	17%	102	14%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	20	3%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	9	1%
Other/unknown	5	29%	1	8%	1	8%	4	22%	2	17%	93	12%
Total	17		12		12		18		12		749	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 2

Characteristic	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex						
Males	7	70%	4	100%	3	50%
Females	3	30%	-	-	3	50%
Age at AIDS diagnosis (years)						
0-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	4	40%	1	25%	3	50%
35-39	1	10%	-	-	1	17%
40-44	1	10%	-	-	1	17%
45-49	1	10%	1	25%	-	-
50-54	-	-	-	-	-	-
55-64	1	10%	1	25%	-	-
65+	2	20%	1	25%	1	17%
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African American	4	40%	1	25%	3	50%
Hispanic/Latino	2	20%	2	50%	-	-
White	4	40%	1	25%	3	50%
Multi-Race	-	-	-	-	-	-
Race/Ethnicity^b and Sex						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African American Males	1	10%	1	25%	-	-
Black/African American Females	3	30%	-	-	3	50%
Hispanic/Latino Males	2	20%	2	50%	-	-
Hispanic/Latina Females	-	-	-	-	-	-
White Males	4	40%	1	25%	3	50%
White Females	-	-	-	-	-	-
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
Total	10		4		6	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 2

Transmission Category ^b	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	3	43%	1	25%	2	67%
Injection drug use (IDU)	1	14%	1	25%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Other/unknown	3	43%	2	50%	1	33%
Subtotal	7	100%	4	100%	3	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	3	100%	-	-	3	100%
Other/unknown	-	-	-	-	-	-
Subtotal	3	100%	-	-	3	100%
Child (<13 years at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	10		4		6	

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 2

Exposure Category ^c	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	30%	1	25%	2	33%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	3	30%	-	-	3	50%
Male-to-male sexual contact & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	1	10%	1	25%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	3	30%	2	50%	1	17%
Total	10		4		6	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 2

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	9	90%	3	75%	6	86%	5	63%	7	70%	430	83%
Females	1	10%	1	25%	1	14%	3	38%	3	30%	87	17%
Age at AIDS diagnosis (years)												
0-9	-	-	-	-	-	-	-	-	-	-	6	1%
10-14	-	-	-	-	1	14%	-	-	-	-	3	1%
15-19	-	-	-	-	-	-	-	-	-	-	2	<1%
20-24	1	10%	1	25%	-	-	-	-	-	-	24	5%
25-29	4	40%	1	25%	1	14%	2	25%	-	-	79	15%
30-34	2	20%	-	-	2	29%	2	25%	4	40%	87	17%
35-39	-	-	-	-	-	-	-	-	1	10%	102	20%
40-44	-	-	-	-	1	14%	2	25%	1	10%	77	15%
45-49	-	-	-	-	-	-	-	-	1	10%	65	13%
50-54	2	20%	-	-	1	14%	-	-	-	-	25	5%
55-64	1	10%	1	25%	-	-	1	13%	1	10%	34	7%
65+	-	-	1	25%	1	14%	1	13%	2	20%	13	3%
Race/Ethnicity ^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
Black/African American	2	20%	1	25%	2	29%	3	38%	4	40%	85	16%
Hispanic/Latino	-	-	-	-	-	-	-	-	2	20%	21	4%
White	7	70%	3	75%	5	71%	5	63%	4	40%	400	77%
Multi-Race	1	10%	-	-	-	-	-	-	-	-	10	2%
Race/Ethnicity ^a and Sex												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
Black/African American Males	1	10%	1	25%	2	29%	2	25%	1	10%	60	12%
Black/African American Females	1	10%	-	-	-	-	1	13%	3	30%	25	5%
Hispanic/Latino Males	-	-	-	-	-	-	-	-	2	20%	17	3%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	4	1%
White Males	7	70%	2	50%	4	57%	3	38%	4	40%	345	67%
White Females	-	-	1	25%	1	14%	2	25%	-	-	55	11%
Multi-Race Males	1	10%	-	-	-	-	-	-	-	-	7	1%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	3	1%
Total	10		4		7		8		10		517	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	6	67%	1	33%	5	83%	5	100%	3	43%	284	67%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	14%	22	5%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	35	8%
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	20	5%
Other/unknown	3	33%	2	67%	1	17%	-	-	3	43%	65	15%
Subtotal	9	100%	3	100%	6	100%	5	100%	7	100%	426	100%
Female adult or adolescent												
Injection drug use	-	-	1	100%	-	-	-	-	-	-	13	16%
Heterosexual contact	1	100%	-	-	-	-	3	100%	3	100%	66	80%
Other/unknown	-	-	-	-	-	-	-	-	-	-	4	5%
Subtotal	1	100%	1	100%	-	-	3	100%	3	100%	83	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	63%
Other/unknown	-	-	-	-	1	100%	-	-	-	-	3	38%
Subtotal	-	-	-	-	1	100%	-	-	-	-	8	100%
Total	10		4		7		8		10		517	

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 2

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	6	60%	1	25%	5	71%	4	50%	3	30%	204	39%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	6	1%
Heterosexual contact only	2	20%	1	25%	1	14%	3	38%	3	30%	107	21%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	16	3%
IDU & Heterosexual contact	-	-	1	25%	-	-	-	-	1	10%	29	6%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	1	13%	-	-	80	15%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	19	4%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	2	20%	1	25%	1	14%	-	-	3	30%	51	10%
Total	10		4		7		8		10		517	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 2

Counties: Ashland, Crawford, Erie, Huron, Knox, Marion, Richland,
Seneca, Wyandot

Reported persons living with diagnosed HIV infection in 2024 by current disease status and selected characteristics, HIV Prevention Planning Region 2

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2024			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex							
Males	131.3	368	78%	181	79%	187	78%
Females	37.0	102	22%	49	21%	53	22%
Age at end of year (years)							
0-9	*	-	-	-	-	-	-
10-14	*	-	-	-	-	-	-
15-19	*	3	1%	2	1%	1	<1%
20-24	31.8	11	2%	11	5%	-	-
25-29	57.0	19	4%	12	5%	7	3%
30-34	111.4	39	8%	29	13%	10	4%
35-39	109.5	36	8%	25	11%	11	5%
40-44	151.5	50	11%	24	10%	26	11%
45-49	152.4	49	10%	24	10%	25	10%
50-54	147.1	49	10%	21	9%	28	12%
55-64	188.2	135	29%	58	25%	77	32%
65+	67.4	79	17%	24	10%	55	23%
Race/Ethnicity ^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	2	<1%	1	<1%	1	<1%
Black/African American	300.3	79	17%	39	17%	40	17%
Hispanic/Latino	234.6	49	10%	28	12%	21	9%
White	64.8	318	68%	153	67%	165	69%
Multi-Race	173.2	21	4%	8	3%	13	5%
Unknown	*	1	<1%	1	<1%	-	-
Race/Ethnicity ^b and Sex							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	2	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African American Males	336.2	55	12%	29	13%	26	11%
Black/African American Females	241.2	24	5%	10	4%	14	6%
Hispanic/Latino Males	378.3	42	9%	27	12%	15	6%
Hispanic/Latina Females	71.5	7	1%	1	<1%	6	3%
White Males	102.9	251	53%	117	51%	134	56%
White Females	27.2	67	14%	36	16%	31	13%
Multi-Race Males	291.1	18	4%	7	3%	11	5%
Multi-Race Females	*	3	1%	1	<1%	2	1%
Unknown	*	1	<1%	1	<1%	-	-
Total	84.6	470		230		240	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^bHispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by current disease status and transmission category, HIV Prevention Planning Region 2

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	260	71%	127	71%	133	71%
Injection drug use (IDU)	6	2%	2	1%	4	2%
Male-to-male sexual contact and IDU	21	6%	8	4%	13	7%
Heterosexual contact	24	7%	8	4%	16	9%
Other/unknown	56	15%	35	19%	21	11%
Subtotal	367	100%	180	100%	187	100%
Female adult or adolescent						
Injection drug use	14	15%	7	15%	7	14%
Heterosexual contact	71	74%	30	65%	41	82%
Other/unknown	11	11%	9	20%	2	4%
Subtotal	96	100%	46	100%	50	100%
Child (<13 years at diagnosis)						
Perinatal	6	86%	4	100%	2	67%
Other/unknown	1	14%	-	-	1	33%
Subtotal	7	100%	4	100%	3	100%
Total	470		230		240	

Reported persons living with diagnosed HIV infection in 2024 by current disease status and exposure category, HIV Prevention Planning Region 2

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	196	42%	98	43%	98	41%
Injection drug use (IDU) only	2	<1%	1	<1%	1	<1%
Heterosexual contact only	116	25%	48	21%	68	28%
Male-to-male sexual contact & IDU	9	2%	5	2%	4	2%
IDU & Heterosexual contact	18	4%	8	3%	10	4%
Male-to-male sexual contact & Heterosexual contact	64	14%	29	13%	35	15%
Male-to-male sexual contact & IDU & Heterosexual contact	12	3%	3	1%	9	4%
Perinatal exposure	6	1%	4	2%	2	1%
Other/unknown	47	10%	34	15%	13	5%
Total	470		230		240	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and age, HIV Prevention Planning Region 2

Living with diagnosed HIV infection in 2024																				
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	205.7	5	6%	*	2	4%	*	4	1%	*	-	-	-	-
25-29	*	-	-	*	-	-	*	4	5%	318.5	5	10%	24.7	7	2%	*	3	14%	-	-
30-34	*	-	-	*	2	100%	315.8	8	10%	*	4	8%	84.2	25	8%	*	-	-	-	-
35-39	*	-	-	*	-	-	344.5	7	9%	413.8	6	12%	73.7	21	7%	*	2	10%	-	-
40-44	*	-	-	*	-	-	595.2	11	14%	344.6	5	10%	111.0	32	10%	*	2	10%	-	-
45-49	*	-	-	*	-	-	452.7	8	10%	413.9	5	10%	119.7	34	11%	*	2	10%	-	-
50-54	*	-	-	*	-	-	450.2	7	9%	640.3	6	12%	119.1	36	11%	*	-	-	-	-
55-64	*	-	-	*	-	-	646.6	17	22%	669.8	10	20%	153.4	102	32%	1,222.0	6	29%	-	-
65+	*	-	-	*	-	-	370.6	12	15%	398.4	6	12%	48.8	54	17%	710.9	6	29%	1	100%
Total	*	-	-	*	2	-	300.3	79	-	234.6	49	-	64.8	318	-	173.2	21	-	1	-

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 2

Living with diagnosed HIV infection in 2024														
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	1	50%	35	64%	22	52%	188	75%	14	78%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	1	2%	5	2%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	2%	4	10%	14	6%	2	11%	-	-
Heterosexual contact	-	-	-	-	9	16%	6	14%	9	4%	-	-	-	-
Other/unknown	-	-	1	50%	10	18%	9	21%	34	14%	2	11%	-	-
Subtotal	-	-	2	100%	55	100%	42	100%	250	100%	18	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	-	-	-	-	12	19%	2	67%	-	-
Heterosexual contact	-	-	-	-	20	87%	5	83%	44	70%	1	33%	1	100%
Other/unknown	-	-	-	-	3	13%	1	17%	7	11%	-	-	-	-
Subtotal	-	-	-	-	23	100%	6	100%	63	100%	3	100%	1	100%
Child (<13 years at diagnosis)														
Perinatal	-	-	-	-	1	100%	1	100%	4	80%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	20%	-	-	-	-
Subtotal	-	-	-	-	1	100%	1	100%	5	100%	-	-	-	-
Total	-	-	2	-	79	-	49	-	318	-	21	-	1	-

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 2

Living with diagnosed HIV infection in 2024														
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
Exposure Category ^d	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	1	50%	17	22%	17	35%	149	47%	12	57%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	2	1%	-	-	-	-
Heterosexual contact only	-	-	-	-	34	43%	14	29%	65	20%	2	10%	1	100%
Male-to-male sexual contact & IDU	-	-	-	-	1	1%	1	2%	6	2%	1	5%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	1	2%	15	5%	2	10%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	18	23%	5	10%	39	12%	2	10%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	3	6%	8	3%	1	5%	-	-
Perinatal exposure	-	-	-	-	1	1%	1	2%	4	1%	-	-	-	-
Other/unknown	-	-	1	50%	8	10%	7	14%	30	9%	1	5%	-	-
Total	-	-	2	-	79	-	49	-	318	-	21	-	1	-

Notes:
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 2, 2020-2024

Living with Diagnosed HIV Infection															
Characteristic	2020			2021			2022			2023			2024		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	125.3	345	79%	120.5	338	78%	124.6	350	79%	129.9	364	78%	131.3	368	78%
Females	33.4	92	21%	33.6	93	22%	33.1	91	21%	36.4	100	22%	37.0	102	22%
Age at end of year (years)															
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
10-14	*	4	1%	*	4	1%	*	1	<1%	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	2	<1%	*	3	1%	*	3	1%
20-24	35.1	12	3%	31.7	11	3%	33.3	12	3%	32.1	11	2%	31.8	11	2%
25-29	96.1	33	8%	85.9	29	7%	62.6	21	5%	78.2	26	6%	57.0	19	4%
30-34	133.7	43	10%	111.6	37	9%	118.6	40	9%	92.7	32	7%	111.4	39	8%
35-39	98.6	31	7%	98.9	32	7%	118.2	38	9%	136.0	44	9%	109.5	36	8%
40-44	138.7	44	10%	131.0	43	10%	118.6	39	9%	121.2	40	9%	151.5	50	11%
45-49	121.1	39	9%	141.1	45	10%	136.4	43	10%	151.2	48	10%	152.4	49	10%
50-54	191.9	65	15%	149.1	52	12%	160.1	55	12%	173.7	59	13%	147.1	49	10%
55-64	151.0	116	27%	155.8	120	28%	168.1	126	29%	171.5	126	27%	188.2	135	29%
65+	45.3	50	11%	52.4	58	13%	56.7	64	15%	65.2	75	16%	67.4	79	17%
Race/Ethnicity ^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	2	<1%
Black/African American	331.9	83	19%	308.9	80	19%	310.2	81	18%	320.7	84	18%	300.3	79	17%
Hispanic/Latino	179.8	33	8%	188.7	36	8%	178.5	35	8%	201.6	40	9%	234.6	49	10%
White	61.6	303	69%	60.1	298	69%	62.0	306	69%	64.6	318	69%	64.8	318	68%
Multi-Race	160.6	17	4%	145.6	16	4%	156.4	18	4%	178.1	21	5%	173.2	21	4%
Unknown	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Race/Ethnicity ^b and Sex															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	2	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African American Males	407.9	62	14%	377.0	60	14%	383.4	62	14%	385.1	63	14%	336.2	55	12%
Black/African American Females	214.1	21	5%	200.4	20	5%	191.1	19	4%	213.6	21	5%	241.2	24	5%
Hispanic/Latino Males	299.5	29	7%	306.7	31	7%	286.9	30	7%	342.4	36	8%	378.3	42	9%
Hispanic/Latina Females	*	4	1%	55.7	5	1%	54.6	5	1%	*	4	1%	71.5	7	1%
White Males	98.9	240	55%	94.9	234	54%	98.8	243	55%	100.9	247	53%	102.9	251	53%
White Females	25.3	63	14%	25.6	64	15%	25.5	63	14%	28.7	71	15%	27.2	67	14%
Multi-Race Males	261.5	14	3%	233.3	13	3%	256.5	15	3%	301.9	18	4%	291.1	18	4%
Multi-Race Females	*	3	1%	*	3	1%	*	3	1%	*	3	1%	*	3	1%
Unknown	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Total	79.3	437		77.3	431		79.4	441		83.6	464		84.6	470	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 2, 2020-2024

Living with Diagnosed HIV Infection										
Transmission Category ^a	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	245	71%	243	72%	252	72%	263	72%	260	71%
Injection drug use (IDU)	9	3%	11	3%	9	3%	7	2%	6	2%
Male-to-male sexual contact and IDU	21	6%	17	5%	20	6%	21	6%	21	6%
Heterosexual contact	24	7%	21	6%	22	6%	24	7%	24	7%
Other/unknown	45	13%	45	13%	46	13%	48	13%	56	15%
Subtotal	344	100%	337	100%	349	100%	363	100%	367	100%
Female adult or adolescent										
Injection drug use	15	18%	14	16%	14	17%	19	20%	14	15%
Heterosexual contact	61	72%	62	73%	61	73%	64	69%	71	74%
Other/unknown	9	11%	9	11%	9	11%	10	11%	11	11%
Subtotal	85	100%	85	100%	84	100%	93	100%	96	100%
Child (<13 years at diagnosis)										
Perinatal	6	75%	7	78%	7	88%	7	88%	6	86%
Other/unknown	2	25%	2	22%	1	13%	1	13%	1	14%
Subtotal	8	100%	9	100%	8	100%	8	100%	7	100%
Total	437		431		441		464		470	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 2, 2020-2024

Living with Diagnosed HIV Infection										
Exposure Category ^b	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	178	41%	178	41%	186	42%	195	42%	196	42%
Injection drug use (IDU) only	3	1%	2	<1%	2	<1%	1	<1%	2	<1%
Heterosexual contact only	100	23%	98	23%	98	22%	105	23%	116	25%
Male-to-male sexual contact & IDU	12	3%	8	2%	7	2%	10	2%	9	2%
IDU & Heterosexual contact	21	5%	23	5%	21	5%	25	5%	18	4%
Male-to-male sexual contact & Heterosexual contact	67	15%	65	15%	66	15%	68	15%	64	14%
Male-to-male sexual contact & IDU & Heterosexual contact	9	2%	9	2%	13	3%	11	2%	12	3%
Perinatal exposure	6	1%	7	2%	7	2%	7	2%	6	1%
Other/unknown	41	9%	41	10%	41	9%	42	9%	47	10%
Total	437		431		441		464		470	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.