

Persons Living with Diagnosed HIV Infection Reported in Clark County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Clark County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	191.2	124	76%	57	74%	67	77%
Females	58.2	40	24%	20	26%	20	23%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	4	2%	4	5%	-	-
25-29	186.3	15	9%	11	14%	4	5%
30-34	166.4	13	8%	8	10%	5	6%
35-39	228.1	17	10%	12	16%	5	6%
40-44	260.7	19	12%	9	12%	10	11%
45-49	193.7	15	9%	6	8%	9	10%
50-54	282.8	24	15%	7	9%	17	20%
55-64	228.7	42	26%	12	16%	30	34%
65+	55.6	15	9%	8	10%	7	8%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	1	1%	-	-	1	1%
Black/African-American	428.2	50	30%	24	31%	26	30%
Hispanic/Latinx	100.8	5	3%	2	3%	3	3%
White	89.5	100	61%	46	60%	54	62%
Multi-Race	179.5	7	4%	4	5%	3	3%
Unknown	*	1	1%	1	1%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	-	-	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African-American Males	553.4	32	20%	15	19%	17	20%
Black/African-American Females	305.3	18	11%	9	12%	9	10%
Hispanic/Latino Males	184.0	5	3%	2	3%	3	3%
Hispanic/Latina Females	*	-	-	-	-	-	-
White Males	148.5	80	49%	36	47%	44	51%
White Females	34.6	20	12%	10	13%	10	11%
Multi-Race Males	312.8	6	4%	4	5%	2	2%
Multi-Race Females	*	1	1%	-	-	1	1%
Unknown	*	1	1%	1	1%	-	-
Total	122.8	164		77		87	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Clark County

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	92	74%	44	77%	48	72%
Injection drug use (IDU)	2	2%	1	2%	1	1%
Male-to-male sexual contact and IDU	5	4%	1	2%	4	6%
Heterosexual contact	9	7%	2	4%	7	10%
Other/unknown	16	13%	9	16%	7	10%
Subtotal	124	100%	57	100%	67	100%
Female adult or adolescent						
Injection drug use	6	15%	4	20%	2	10%
Heterosexual contact	31	78%	15	75%	16	80%
Other/unknown	3	8%	1	5%	2	10%
Subtotal	40	100%	20	100%	20	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	164		77		87	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Clark County

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	91	55%	44	57%	47	54%
Injection drug use (IDU) only	2	1%	1	1%	1	1%
Heterosexual contact only	40	24%	17	22%	23	26%
Male-to-male sexual contact & IDU	4	2%	1	1%	3	3%
IDU & Heterosexual contact	6	4%	4	5%	2	2%
Male-to-male sexual contact & Heterosexual contact	1	1%	-	-	1	1%
Male-to-male sexual contact & IDU & Heterosexual contact	1	1%	-	-	1	1%
Perinatal exposure	-	-	-	-	-	-
Other/unknown	19	12%	10	13%	9	10%
Total	164		77		87	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Clark County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	<5	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	<5	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	<5	-	-	-
20-24	*	-	-	*	-	-	*	1	2%	*	-	-	*	3	3%	*	<5	-	-	-
25-29	*	-	-	*	-	-	683.4	6	12%	*	1	20%	123.2	8	8%	*	<5	-	-	-
30-34	*	-	-	*	-	-	995.7	7	14%	*	1	20%	77.8	5	5%	*	<5	-	-	-
35-39	*	-	-	*	-	-	1,067.1	7	14%	*	-	-	129.6	8	8%	*	<5	-	-	-
40-44	*	-	-	*	1	100%	894.2	6	12%	*	1	20%	162.7	10	10%	*	<5	-	-	-
45-49	*	-	-	*	-	-	*	3	6%	*	1	20%	149.8	10	10%	*	<5	-	-	-
50-54	*	-	-	*	-	-	1,240.3	8	16%	*	-	-	201.6	15	15%	*	<5	-	-	-
55-64	*	-	-	*	-	-	569.8	8	16%	*	1	20%	189.9	31	31%	*	<5	-	-	-
65+	*	-	-	*	-	-	*	4	8%	*	-	-	41.0	10	10%	*	<5	-	1	100%
Total	*	-	-	*	1	-	428.2	50	-	100.8	5	-	89.5	100	-	179.5	7	-	1	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Clark County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	23	72%	4	80%	59	74%	6	100%	-	-
Injection drug use (IDU)	-	-	-	-	1	3%	1	20%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	3%	-	-	4	5%	-	-	-	-
Heterosexual contact	-	-	-	-	4	13%	-	-	5	6%	-	-	-	-
Other/unknown	-	-	1	100%	3	9%	-	-	12	15%	-	-	-	-
Subtotal	-	-	1	100%	32	100%	5	100%	80	100%	6	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	2	11%	-	-	4	20%	-	-	-	-
Heterosexual contact	-	-	-	-	15	83%	-	-	15	75%	1	100%	-	-
Other/unknown	-	-	-	-	1	6%	-	-	1	5%	-	-	1	100%
Subtotal	-	-	-	-	18	100%	-	-	20	100%	1	100%	1	100%
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	1	-	50	-	5	-	100	-	7	-	1	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Clark County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	23	46%	4	80%	58	58%	6	86%	-	-
Injection drug use (IDU) only	-	-	-	-	1	2%	1	20%	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	19	38%	-	-	20	20%	1	14%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	4	4%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	2	4%	-	-	4	4%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	1	1%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	1	2%	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	100%	4	8%	-	-	13	13%	-	-	1	100%
Total	-	-	1	-	50	-	5	-	100	-	7	-	1	-

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Clark County, 2016-2020

Characteristic	Living with Diagnosed HIV Infection														
	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	156.0	102	74%	153.4	100	74%	161.2	105	73%	173.8	113	73%	191.2	124	76%
Females	51.9	36	26%	51.9	36	26%	54.7	38	27%	59.4	41	27%	58.2	40	24%
Age at end of year															
<13	*	1	1%	*	1	1%	*	1	1%	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	2	1%	*	2	1%	*	3	2%	*	2	1%	*	4	2%
25-29	208.4	17	12%	147.3	12	9%	123.0	10	7%	148.1	12	8%	186.3	15	9%
30-34	91.8	7	5%	133.1	10	7%	170.7	13	9%	220.1	17	11%	166.4	13	8%
35-39	148.6	11	8%	148.7	11	8%	135.4	10	7%	216.0	16	10%	228.1	17	10%
40-44	204.5	15	11%	153.4	11	8%	167.5	12	8%	206.0	15	10%	260.7	19	12%
45-49	220.7	19	14%	260.5	22	16%	263.7	22	15%	249.8	20	13%	193.7	15	9%
50-54	302.0	28	20%	289.3	26	19%	217.3	19	13%	232.0	20	13%	282.8	24	15%
55-64	163.4	31	22%	159.4	30	22%	224.7	42	29%	199.6	37	24%	228.7	42	26%
65+	27.7	7	5%	43.0	11	8%	42.1	11	8%	56.7	15	10%	55.6	15	9%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Black/African-American	340.5	40	29%	338.8	40	29%	357.7	42	29%	384.9	45	29%	428.2	50	30%
Hispanic/Latinx	*	3	2%	*	3	2%	*	2	1%	*	4	3%	100.8	5	3%
White	79.2	90	65%	77.7	88	65%	81.3	92	64%	84.5	95	62%	89.5	100	61%
Multi-Race	*	3	2%	*	3	2%	136.5	5	3%	216.5	8	5%	179.5	7	4%
Unknown	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African-American Males	416.9	24	17%	394.8	23	17%	433.3	25	17%	452.6	26	17%	553.4	32	20%
Black/African-American Females	267.0	16	12%	284.3	17	13%	284.7	17	12%	319.6	19	12%	305.3	18	11%
Hispanic/Latino Males	*	3	2%	*	3	2%	*	2	1%	*	4	3%	184.0	5	3%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
White Males	131.1	72	52%	130.0	71	52%	134.2	73	51%	138.4	75	49%	148.5	80	49%
White Females	30.7	18	13%	29.0	17	13%	32.4	19	13%	34.3	20	13%	34.6	20	12%
Multi-Race Males	*	2	1%	*	2	1%	*	4	3%	382.3	7	5%	312.8	6	4%
Multi-Race Females	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Unknown	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Total	102.4	138		101.1	136		106.3	143		114.9	154		122.8	164	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Clark County, 2016-2020

Living with Diagnosed HIV Infection										
Transmission Category ^a	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	76	75%	71	72%	78	75%	87	77%	92	74%
Injection drug use (IDU)	3	3%	3	3%	2	2%	3	3%	2	2%
Male-to-male sexual contact and IDU	-	-	2	2%	3	3%	3	3%	5	4%
Heterosexual contact	10	10%	11	11%	8	8%	9	8%	9	7%
Other/unknown	12	12%	12	12%	13	13%	11	10%	16	13%
Subtotal	101	100%	99	100%	104	100%	113	100%	124	100%
Female adult or adolescent										
Injection drug use	5	14%	5	14%	5	13%	8	20%	6	15%
Heterosexual contact	28	78%	28	78%	30	79%	30	73%	31	78%
Other/unknown	3	8%	3	8%	3	8%	3	7%	3	8%
Subtotal	36	100%	36	100%	38	100%	41	100%	40	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	1	100%	-	-	-	-
Total	138		136		143		154		164	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Clark County, 2016-2020

Living with Diagnosed HIV Infection										
Exposure Category ^b	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	73	53%	69	51%	77	54%	86	56%	91	55%
Injection drug use (IDU) only	3	2%	3	2%	2	1%	4	3%	2	1%
Heterosexual contact only	38	28%	39	29%	38	27%	39	25%	40	24%
Male-to-male sexual contact & IDU	-	-	2	1%	3	2%	3	2%	4	2%
IDU & Heterosexual contact	5	4%	5	4%	5	3%	7	5%	6	4%
Male-to-male sexual contact & Heterosexual contact	3	2%	2	1%	1	1%	1	1%	1	1%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	16	12%	16	12%	17	12%	14	9%	19	12%
Total	138		136		143		154		164	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.