



Department of
Health

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 7

Counties: Adams, Fayette, Gallia, Hocking, Jackson, Lawrence,
Pike, Ross, Scioto, Vinton

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2025

Reported new diagnoses of HIV infection in 2024 by disease status and selected characteristics, HIV Prevention Planning Region 7

Characteristic	2024 diagnosis of HIV infection			Disease Status						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	*	3	33%	3	43%	-	-	-	-	
Females	3.1	6	67%	4	57%	2	100%	-	-	
Age at diagnosis (years)										
0-9	*	-	-	-	-	-	-	-	-	
10-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	*	1	11%	1	14%	-	-	-	-	
25-29	*	-	-	-	-	-	-	-	-	
30-34	*	3	33%	1	14%	2	100%	-	-	
35-39	*	3	33%	3	43%	-	-	-	-	
40-44	*	1	11%	1	14%	-	-	-	-	
45-49	*	1	11%	1	14%	-	-	-	-	
50-54	*	-	-	-	-	-	-	-	-	
55-64	*	-	-	-	-	-	-	-	-	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b										
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
Black/African American	*	-	-	-	-	-	-	-	-	
Hispanic/Latino	*	-	-	-	-	-	-	-	-	
White	2.5	9	100%	7	100%	2	100%	-	-	
Multi-Race	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b and Sex										
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
Black/African American Males	*	-	-	-	-	-	-	-	-	
Black/African American Females	*	-	-	-	-	-	-	-	-	
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
White Males	*	3	33%	3	43%	-	-	-	-	
White Females	3.3	6	67%	4	57%	2	100%	-	-	
Multi-Race Males	*	-	-	-	-	-	-	-	-	
Multi-Race Females	*	-	-	-	-	-	-	-	-	
Total	2.3	9		7		2		-		

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by disease status and transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	3	100%	3	100%	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	3	100%	3	100%	-	-	-	-
Female adult or adolescent								
Injection drug use	2	33%	2	50%	-	-	-	-
Heterosexual contact	3	50%	1	25%	2	100%	-	-
Other/unknown	1	17%	1	25%	-	-	-	-
Subtotal	6	100%	4	100%	2	100%	-	-
Child (<13 years at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	9		7		2		-	

Reported new diagnoses of HIV infection in 2024 by disease status and exposure category, HIV Prevention Planning Region 7

Exposure Category ^b	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2	22%	2	29%	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	3	33%	1	14%	2	100%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	2	22%	2	29%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	1	11%	1	14%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	1	11%	1	14%	-	-	-	-
Total	9		7		2		-	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 7

2024 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
0-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	-	-	*	-	-	*	1	11%	*	-	-
25-29	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
30-34	*	-	-	*	-	-	*	-	-	*	-	-	*	3	33%	*	-	-
35-39	*	-	-	*	-	-	*	-	-	*	-	-	*	3	33%	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	1	11%	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	1	11%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	*	-	-	*	-	-	2.5	9	-	*	-	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 7

Transmission Category ^c	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	-	-	-	-	3	100%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	3	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	2	33%	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	3	50%	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	17%	-	-
Subtotal	-	-	-	-	-	-	-	-	6	100%	-	-
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-		-		-		-		9		-	

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 7

Exposure Category ^d	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-	-	-	2	22%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	-	-	-	-	3	33%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	2	22%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	1	11%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	11%	-	-
Total	-	-	-	-	-	-	-	-	9	-	-	-

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
Asterisk (*) indicates rate not calculated. for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 7

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2020			2021			2022			2023			2024				
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex																	
Males	5.1	10	83%	5.1	10	56%	5.6	11	85%	4.6	9	82%	*	3	33%	446	81%
Females	*	2	17%	4.1	8	44%	*	2	15%	*	2	18%	3.1	6	67%	106	19%
Age at diagnosis (years)																	
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	6	1%
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	1%
15-19	*	1	8%	*	-	-	*	-	-	*	-	-	*	-	-	7	1%
20-24	*	1	8%	*	1	6%	*	1	8%	*	1	9%	*	1	11%	53	10%
25-29	*	2	17%	*	4	22%	*	3	23%	*	-	-	*	-	-	87	16%
30-34	*	4	33%	*	3	17%	*	4	31%	*	4	36%	*	3	33%	123	22%
35-39	*	2	17%	*	3	17%	*	3	23%	*	1	9%	*	3	33%	95	17%
40-44	*	1	8%	*	-	-	*	1	8%	*	-	-	*	1	11%	72	13%
45-49	*	1	8%	*	3	17%	*	1	8%	*	1	9%	*	1	11%	39	7%
50-54	*	-	-	*	2	11%	*	-	-	*	3	27%	*	-	-	37	7%
55-64	*	-	-	*	2	11%	*	-	-	*	-	-	*	-	-	22	4%
65+	*	-	-	*	-	-	*	-	-	*	1	9%	*	-	-	8	1%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	9%	*	-	-	1	<1%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Black/African American	*	-	-	*	-	-	*	2	15%	*	-	-	*	-	-	35	6%
Hispanic/Latino	*	1	8%	*	1	6%	*	-	-	*	1	9%	*	-	-	11	2%
White	2.7	10	83%	4.1	15	83%	3.0	11	85%	2.5	9	82%	2.5	9	100%	485	88%
Multi-Race	*	1	8%	*	2	11%	*	-	-	*	-	-	*	-	-	18	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Race/Ethnicity ^b and Sex																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	1	9%	*	-	-	1	<1%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African American Males	*	-	-	*	-	-	*	1	8%	*	-	-	*	-	-	23	4%
Black/African American Females	*	-	-	*	-	-	*	1	8%	*	-	-	*	-	-	12	2%
Hispanic/Latino Males	*	1	8%	*	-	-	*	-	-	*	1	9%	*	-	-	8	1%
Hispanic/Latina Females	*	-	-	*	1	6%	*	-	-	*	-	-	*	-	-	3	1%
White Males	4.4	8	67%	4.9	9	50%	5.5	10	77%	4.5	8	73%	*	3	33%	399	72%
White Females	*	2	17%	3.2	6	33%	*	1	8%	*	1	9%	3.3	6	67%	86	16%
Multi-Race Males	*	1	8%	*	1	6%	*	-	-	*	-	-	*	-	-	14	3%
Multi-Race Females	*	-	-	*	1	6%	*	-	-	*	-	-	*	-	-	4	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Total	3.0	12		4.6	18		3.3	13		2.8	11		2.3	9		552	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	2	20%	7	70%	3	27%			3	100%	278	63%
Injection drug use (IDU)	5	50%	1	10%	4	36%	2	22%	-	-	35	8%
Male-to-male sexual contact and IDU	1	10%	1	10%	1	9%	-	-	-	-	27	6%
Heterosexual contact	-	-	1	10%	-	-	-	-	-	-	17	4%
Other/unknown	2	20%	-	-	3	27%	4	44%	-	-	84	19%
Subtotal	10	100%	10	100%	11	100%	9	100%	3	100%	441	100%
Female adult or adolescent												
Injection drug use	2	100%	3	38%	1	50%	-	-	2	33%	25	24%
Heterosexual contact	-	-	4	50%	1	50%	2	100%	3	50%	59	57%
Other/unknown	-	-	1	13%	-	-	-	-	1	17%	20	19%
Subtotal	2	100%	8	100%	2	100%	2	100%	6	100%	104	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	2	29%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	71%
Subtotal	-	-	-	-	-	-	-	-	-	-	7	100%
Total	12		18		13		11		9		552	

Trends in reported diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 7

Diagnoses of HIV Infection by Year												
Exposure Category ^b	2020		2021		2022		2023		2024		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	8%	4	22%	1	8%	1	9%	2	22%	209	38%
Injection drug use (IDU) only	2	17%	1	6%	-	-	-	-	-	-	10	2%
Heterosexual contact only	-	-	5	28%	1	8%	5	45%	3	33%	108	20%
Male-to-male sexual contact & IDU	-	-	-	-	1	8%	-	-	-	-	14	3%
IDU & Heterosexual contact	5	42%	3	17%	5	38%	2	18%	2	22%	48	9%
Male-to-male sexual contact & Heterosexual contact	1	8%	3	17%	2	15%	2	18%	1	11%	69	13%
Male-to-male sexual contact & IDU & Heterosexual contact	1	8%	1	6%	-	-	-	-	-	-	13	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	2	<1%
Other/unknown	2	17%	1	6%	3	23%	1	9%	1	11%	79	14%
Total	12		18		13		11		9		552	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 7

Characteristic	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex						
Males	1	33%	1	100%	-	-
Females	2	67%	-	-	2	100%
Age at AIDS diagnosis (years)						
0-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	1	33%	-	-	1	50%
35-39	-	-	-	-	-	-
40-44	1	33%	1	100%	-	-
45-49	-	-	-	-	-	-
50-54	-	-	-	-	-	-
55-64	-	-	-	-	-	-
65+	1	33%	-	-	1	50%
Race/Ethnicity^b						
American Indian/Alaska Native	1	33%	-	-	1	50%
Asian/Pacific Islander	-	-	-	-	-	-
Black/African American	-	-	-	-	-	-
Hispanic/Latino	-	-	-	-	-	-
White	2	67%	1	100%	1	50%
Multi-Race	-	-	-	-	-	-
Race/Ethnicity^b and Sex						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	1	33%	-	-	1	50%
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African American Males	-	-	-	-	-	-
Black/African American Females	-	-	-	-	-	-
Hispanic/Latino Males	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-
White Males	1	33%	1	100%	-	-
White Females	1	33%	-	-	1	50%
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
Total	3		1		2	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 7

Transmission Category ^b	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1	100%	1	100%	-	-
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	2	100%	-	-	2	100%
Other/unknown	-	-	-	-	-	-
Subtotal	2	100%	-	-	2	100%
Child (<13 years at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	3		1		2	

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 7

Exposure Category ^c	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	33%	1	100%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	2	67%	-	-	2	100%
Male-to-male sexual contact & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Total	3		1		2	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 7

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	2	50%	6	55%	8	89%	8	89%	1	33%	317	84%
Females	2	50%	5	45%	1	11%	1	11%	2	67%	61	16%
Age at AIDS diagnosis (years)												
0-9	-	-	-	-	-	-	-	-	-	-	2	1%
10-14	-	-	-	-	-	-	-	-	-	-	1	<1%
15-19	-	-	-	-	-	-	-	-	-	-	3	1%
20-24	-	-	1	9%	1	11%	1	11%	-	-	13	3%
25-29	1	25%	-	-	3	33%	-	-	-	-	39	10%
30-34	-	-	3	27%	1	11%	2	22%	1	33%	79	21%
35-39	1	25%	2	18%	-	-	1	11%	-	-	65	17%
40-44	1	25%	-	-	2	22%	-	-	1	33%	76	20%
45-49	1	25%	2	18%	-	-	1	11%	-	-	42	11%
50-54	-	-	2	18%	-	-	3	33%	-	-	28	7%
55-64	-	-	1	9%	2	22%	1	11%	-	-	23	6%
65+	-	-	-	-	-	-	-	-	1	33%	7	2%
Race/Ethnicity ^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	1	33%	1	<1%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	2	1%
Black/African American	-	-	-	-	2	22%	-	-	-	-	25	7%
Hispanic/Latino	-	-	1	9%	-	-	1	11%	-	-	6	2%
White	4	100%	10	91%	7	78%	8	89%	2	67%	335	89%
Multi-Race	-	-	-	-	-	-	-	-	-	-	9	2%
Race/Ethnicity ^a and Sex												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	1	33%	1	<1%
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	2	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
Black/African American Males	-	-	-	-	2	22%	-	-	-	-	19	5%
Black/African American Females	-	-	-	-	-	-	-	-	-	-	6	2%
Hispanic/Latino Males	-	-	-	-	-	-	1	11%	-	-	4	1%
Hispanic/Latina Females	-	-	1	9%	-	-	-	-	-	-	2	1%
White Males	2	50%	6	55%	6	67%	7	78%	1	33%	285	75%
White Females	2	50%	4	36%	1	11%	1	11%	1	33%	50	13%
Multi-Race Males	-	-	-	-	-	-	-	-	-	-	7	2%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	2	1%
Total	4		11		9		9		3		378	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	4	67%	5	63%	3	38%	1	100%	214	68%
Injection drug use (IDU)	-	-	1	17%	-	-	1	13%	-	-	16	5%
Male-to-male sexual contact and IDU	1	50%	1	17%	1	13%	1	13%	-	-	16	5%
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	11	4%
Other/unknown	1	50%	-	-	2	25%	3	38%	-	-	56	18%
Subtotal	2	100%	6	100%	8	100%	8	100%	1	100%	313	100%
Female adult or adolescent												
Injection drug use	1	50%	2	40%	1	100%	1	100%	-	-	18	30%
Heterosexual contact	1	50%	2	40%	-	-	-	-	2	100%	34	57%
Other/unknown	-	-	1	20%	-	-	-	-	-	-	8	13%
Subtotal	2	100%	5	100%	1	100%	1	100%	2	100%	60	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	1	20%
Other/unknown	-	-	-	-	-	-	-	-	-	-	4	80%
Subtotal	-	-	-	-	-	-	-	-	-	-	5	100%
Total	4		11		9		9		3		378	

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 7

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	2	18%	3	33%	-	-	1	33%	157	42%
Injection drug use (IDU) only	-	-	1	9%	-	-	-	-	-	-	5	1%
Heterosexual contact only	1	25%	2	18%	-	-	2	22%	2	67%	66	17%
Male-to-male sexual contact & IDU	1	25%	-	-	1	11%	1	11%	-	-	7	2%
IDU & Heterosexual contact	1	25%	2	18%	1	11%	2	22%	-	-	28	7%
Male-to-male sexual contact & Heterosexual contact	-	-	2	18%	2	22%	3	33%	-	-	57	15%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	9%	-	-	-	-	-	-	9	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	1	<1%
Other/unknown	1	25%	1	9%	2	22%	1	11%	-	-	48	13%
Total	4		11		9		9		3		378	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 7

Counties: Adams, Fayette, Gallia, Hocking, Jackson, Lawrence,
Pike, Ross, Scioto, Vinton

Reported persons living with diagnosed HIV infection in 2024 by current disease status and selected characteristics, HIV Prevention Planning Region 7

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2024			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex							
Males	160.4	313	78%	159	76%	154	80%
Females	45.4	88	22%	50	24%	38	20%
Age at end of year (years)							
0-9	*	-	-	-	-	-	-
10-14	*	1	<1%	1	<1%	-	-
15-19	*	-	-	-	-	-	-
20-24	*	4	1%	4	2%	-	-
25-29	59.2	14	3%	11	5%	3	2%
30-34	176.4	43	11%	28	13%	15	8%
35-39	216.2	50	12%	34	16%	16	8%
40-44	217.5	52	13%	27	13%	25	13%
45-49	174.0	42	10%	24	11%	18	9%
50-54	189.4	49	12%	24	11%	25	13%
55-64	193.4	100	25%	34	16%	66	34%
65+	60.4	46	11%	22	11%	24	13%
Race/Ethnicity ^b							
American Indian/Alaska Native	*	1	<1%	-	-	1	1%
Asian/Pacific Islander	*	-	-	-	-	-	-
Black/African American	331.2	33	8%	18	9%	15	8%
Hispanic/Latino	179.1	10	2%	2	1%	8	4%
White	93.4	337	84%	179	86%	158	82%
Multi-Race	241.9	20	5%	10	5%	10	5%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity ^b and Sex							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	1%
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African American Males	319.6	22	5%	11	5%	11	6%
Black/African American Females	357.1	11	3%	7	3%	4	2%
Hispanic/Latino Males	194.4	6	1%	1	<1%	5	3%
Hispanic/Latina Females	*	4	1%	1	<1%	3	2%
White Males	149.2	267	67%	139	67%	128	67%
White Females	38.5	70	17%	40	19%	30	16%
Multi-Race Males	424.2	18	4%	8	4%	10	5%
Multi-Race Females	*	2	<1%	2	1%	-	-
Unknown	*	-	-	-	-	-	-
Total	103.1	401		209		192	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by current disease status and transmission category, HIV Prevention Planning Region 7

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	204	65%	99	62%	105	69%
Injection drug use (IDU)	32	10%	20	13%	12	8%
Male-to-male sexual contact and IDU	28	9%	14	9%	14	9%
Heterosexual contact	11	4%	8	5%	3	2%
Other/unknown	37	12%	18	11%	19	12%
Subtotal	312	100%	159	100%	153	100%
Female adult or adolescent						
Injection drug use	27	33%	18	38%	9	26%
Heterosexual contact	47	57%	24	50%	23	66%
Other/unknown	9	11%	6	13%	3	9%
Subtotal	83	100%	48	100%	35	100%
Child (<13 years at diagnosis)						
Perinatal	5	83%	2	100%	3	75%
Other/unknown	1	17%	-	-	1	25%
Subtotal	6	100%	2	100%	4	100%
Total	401		209		192	

Reported persons living with diagnosed HIV infection in 2024 by current disease status and exposure category, HIV Prevention Planning Region 7

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	159	40%	80	38%	79	41%
Injection drug use (IDU) only	7	2%	4	2%	3	2%
Heterosexual contact only	78	19%	40	19%	38	20%
Male-to-male sexual contact & IDU	16	4%	9	4%	7	4%
IDU & Heterosexual contact	51	13%	34	16%	17	9%
Male-to-male sexual contact & Heterosexual contact	45	11%	19	9%	26	14%
Male-to-male sexual contact & IDU & Heterosexual contact	12	3%	5	2%	7	4%
Perinatal exposure	5	1%	2	1%	3	2%
Other/unknown	28	7%	16	8%	12	6%
Total	401		209		192	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and age, HIV Prevention Planning Region 7

Living with diagnosed HIV infection in 2024																				
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
10-14	*	-	-	*	-	-	*	1	3%	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	*	1	3%	*	-	-	*	3	1%	*	-	-	-	-
25-29	*	-	-	*	-	-	*	4	12%	*	-	-	46.5	10	3%	*	-	-	-	-
30-34	*	-	-	*	-	-	*	4	12%	*	2	20%	166.4	37	11%	*	-	-	-	-
35-39	*	-	-	*	-	-	*	2	6%	*	1	10%	210.7	45	13%	*	2	10%	-	-
40-44	*	-	-	*	-	-	634.5	5	15%	*	-	-	185.2	41	12%	2,173.9	6	30%	-	-
45-49	*	-	-	*	-	-	*	1	3%	*	3	30%	154.7	35	10%	*	3	15%	-	-
50-54	*	-	-	*	-	-	*	3	9%	*	1	10%	168.0	41	12%	*	4	20%	-	-
55-64	*	-	-	*	-	-	662.9	7	21%	*	2	20%	174.8	86	26%	969.0	5	25%	-	-
65+	*	1	100%	*	-	-	372.6	5	15%	*	1	10%	53.8	39	12%	*	-	-	-	-
Total	*	1		*	-	-	331.2	33		179.1	10		93.4	337		241.9	20		-	-

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 7

Living with diagnosed HIV infection in 2024														
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	14	64%	4	67%	173	65%	13	72%	-	-
Injection drug use (IDU)	-	-	-	-	4	18%	-	-	27	10%	1	6%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	26	10%	2	11%	-	-
Heterosexual contact	-	-	-	-	2	9%	1	17%	8	3%	-	-	-	-
Other/unknown	-	-	-	-	2	9%	1	17%	32	12%	2	11%	-	-
Subtotal	-	-	-	-	22	100%	6	100%	266	100%	18	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	2	25%	-	-	24	35%	1	50%	-	-
Heterosexual contact	1	100%	-	-	5	63%	3	75%	37	54%	1	50%	-	-
Other/unknown	-	-	-	-	1	13%	1	25%	7	10%	-	-	-	-
Subtotal	1	100%	-	-	8	100%	4	100%	68	100%	2	100%	-	-
Child (<13 years at diagnosis)														
Perinatal	-	-	-	-	3	100%	-	-	2	67%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	33%	-	-	-	-
Subtotal	-	-	-	-	3	100%	-	-	3	100%	-	-	-	-
Total	1		-		33		10		337		20		-	

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 7

Living with diagnosed HIV infection in 2024														
Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	9	27%	3	30%	138	41%	9	45%	-	-
Injection drug use (IDU) only	-	-	-	-	1	3%	-	-	6	2%	-	-	-	-
Heterosexual contact only	1	100%	-	-	8	24%	5	50%	63	19%	1	5%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	16	5%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	5	15%	-	-	44	13%	2	10%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	5	15%	1	10%	35	10%	4	20%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	10	3%	2	10%	-	-
Perinatal exposure	-	-	-	-	3	9%	-	-	2	1%	-	-	-	-
Other/unknown	-	-	-	-	2	6%	1	10%	23	7%	2	10%	-	-
Total	1		-		33		10		337		20		-	

Notes:
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 7, 2020-2024

Living with Diagnosed HIV Infection															
Characteristic	2020			2021			2022			2023			2024		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	144.1	285	82%	144.3	285	81%	153.2	301	80%	158.7	310	81%	160.4	313	78%
Females	32.1	64	18%	34.7	68	19%	39.1	76	20%	38.1	74	19%	45.4	88	22%
Age at end of year (years)															
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	1	<1%
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	22.4	5	1%	22.6	5	1%	38.8	9	2%	*	4	1%	*	4	1%
25-29	69.4	17	5%	76.2	18	5%	64.0	15	4%	77.1	18	5%	59.2	14	3%
30-34	117.9	28	8%	137.1	33	9%	164.2	40	11%	192.0	47	12%	176.4	43	11%
35-39	158.4	37	11%	164.3	38	11%	185.5	42	11%	171.7	39	10%	216.2	50	12%
40-44	157.0	38	11%	151.0	37	10%	176.6	43	11%	189.8	46	12%	217.5	52	13%
45-49	171.3	44	13%	169.2	42	12%	168.9	41	11%	165.8	40	10%	174.0	42	10%
50-54	222.2	58	17%	185.2	49	14%	163.1	43	11%	163.3	43	11%	189.4	49	12%
55-64	169.6	95	27%	175.2	97	27%	201.9	108	29%	201.5	106	28%	193.4	100	25%
65+	36.2	27	8%	46.5	34	10%	48.7	36	10%	54.8	41	11%	60.4	46	11%
Race/Ethnicity ^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	<1%	*	1	<1%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	1	<1%	*	-	-
Black/African American	331.5	32	9%	290.4	29	8%	318.8	31	8%	333.6	33	9%	331.2	33	8%
Hispanic/Latino	92.9	5	1%	148.0	8	2%	159.4	9	2%	207.5	11	3%	179.1	10	2%
White	78.2	290	83%	80.4	295	84%	87.1	317	84%	87.7	318	83%	93.4	337	84%
Multi-Race	291.5	22	6%	270.9	21	6%	252.4	20	5%	247.4	20	5%	241.9	20	5%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity ^b and Sex															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	1	<1%	*	1	<1%
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	1	<1%	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African American Males	332.6	22	6%	304.8	21	6%	312.7	21	6%	323.9	22	6%	319.6	22	5%
Black/African American Females	329.2	10	3%	258.2	8	2%	332.2	10	3%	355.0	11	3%	357.1	11	3%
Hispanic/Latino Males	*	3	1%	172.8	5	1%	195.9	6	2%	279.3	8	2%	194.4	6	1%
Hispanic/Latina Females	*	2	1%	*	3	1%	*	3	1%	*	3	1%	*	4	1%
White Males	131.3	240	69%	131.9	240	68%	142.1	257	68%	144.7	260	68%	149.2	267	67%
White Females	26.6	50	14%	29.7	55	16%	32.8	60	16%	31.7	58	15%	38.5	70	17%
Multi-Race Males	526.9	20	6%	484.1	19	5%	420.0	17	5%	458.4	19	5%	424.2	18	4%
Multi-Race Females	*	2	1%	*	2	1%	*	3	1%	*	1	<1%	*	2	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	87.9	349		89.7	353		96.5	377		98.6	384		103.1	401	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 7, 2020-2024

Living with Diagnosed HIV Infection										
Transmission Category ^a	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	195	69%	193	68%	196	65%	199	64%	204	65%
Injection drug use (IDU)	23	8%	22	8%	27	9%	34	11%	32	10%
Male-to-male sexual contact and IDU	23	8%	24	8%	28	9%	25	8%	28	9%
Heterosexual contact	13	5%	12	4%	14	5%	14	5%	11	4%
Other/unknown	30	11%	33	12%	35	12%	37	12%	37	12%
Subtotal	284	100%	284	100%	300	100%	309	100%	312	100%
Female adult or adolescent										
Injection drug use	20	32%	23	35%	25	35%	22	31%	27	33%
Heterosexual contact	34	55%	34	52%	39	54%	42	60%	47	57%
Other/unknown	8	13%	8	12%	8	11%	6	9%	9	11%
Subtotal	62	100%	65	100%	72	100%	70	100%	83	100%
Child (<13 years at diagnosis)										
Perinatal	2	67%	3	75%	4	80%	4	80%	5	83%
Other/unknown	1	33%	1	25%	1	20%	1	20%	1	17%
Subtotal	3	100%	4	100%	5	100%	5	100%	6	100%
Total	349		353		377		384		401	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 7, 2020-2024

Living with Diagnosed HIV Infection										
Exposure Category ^b	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	147	42%	144	41%	145	38%	150	39%	159	40%
Injection drug use (IDU) only	7	2%	6	2%	5	1%	6	2%	7	2%
Heterosexual contact only	61	17%	62	18%	68	18%	73	19%	78	19%
Male-to-male sexual contact & IDU	11	3%	11	3%	17	5%	16	4%	16	4%
IDU & Heterosexual contact	35	10%	38	11%	46	12%	49	13%	51	13%
Male-to-male sexual contact & Heterosexual contact	48	14%	49	14%	51	14%	49	13%	45	11%
Male-to-male sexual contact & IDU & Heterosexual contact	12	3%	13	4%	11	3%	9	2%	12	3%
Perinatal exposure	2	1%	3	1%	4	1%	4	1%	5	1%
Other/unknown	26	7%	27	8%	30	8%	28	7%	28	7%
Total	349		353		377		384		401	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.