

# New Diagnoses of HIV Infection Reported in Lucas County

As of June 30, 2020

## Reported new diagnoses of HIV infection in 2019 by disease status and selected characteristics, Lucas County

Characteristic	2019 diagnosis of HIV infection			Disease Status						
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
<b>Sex at birth</b>										
Males	17.4	36	90%	26	90%	8	89%	2	100%	
Females	*	4	10%	3	10%	1	11%	-	-	
<b>Age at diagnosis (yr)</b>										
<13	*	-	-	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	33.9	9	23%	7	24%	1	11%	1	50%	
25-29	*	2	5%	2	7%	-	-	-	-	
30-34	27.4	8	20%	8	28%	-	-	-	-	
35-39	*	4	10%	3	10%	1	11%	-	-	
40-44	21.3	5	13%	3	10%	2	22%	-	-	
45-49	*	4	10%	1	3%	2	22%	1	50%	
50-54	*	1	3%	1	3%	-	-	-	-	
55-64	8.5	5	13%	3	10%	2	22%	-	-	
65+	*	2	5%	1	3%	1	11%	-	-	
<b>Race/Ethnicity<sup>b</sup></b>										
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
Black/African-American	22.6	19	48%	12	41%	6	67%	1	50%	
Hispanic/Latinx	19.0	6	15%	6	21%	-	-	-	-	
White	4.8	14	35%	10	34%	3	33%	1	50%	
Multi-Race	*	1	3%	1	3%	-	-	-	-	
<b>Race/Ethnicity<sup>b</sup> and Sex at birth</b>										
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
Black/African-American Males	40.5	16	40%	10	34%	5	56%	1	50%	
Black/African-American Females	*	3	8%	2	7%	1	11%	-	-	
Hispanic/Latino Males	38.4	6	15%	6	21%	-	-	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
White Males	9.1	13	33%	9	31%	3	33%	1	50%	
White Females	*	1	3%	1	3%	-	-	-	-	
Multi-Race Males	*	1	3%	1	3%	-	-	-	-	
Multi-Race Females	*	-	-	-	-	-	-	-	-	
<b>Total</b>	<b>9.3</b>	<b>40</b>		<b>29</b>		<b>9</b>		<b>2</b>		

**Notes:**

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2019 U.S. Census estimates.

<sup>b</sup> Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.

## Reported new diagnoses of HIV infection in 2019 by disease status and transmission category, Lucas County

Transmission Category <sup>a</sup>	2019 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	26	72%	20	77%	4	50%	2	100%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	11%	3	12%	1	13%	-	-
Heterosexual contact	1	3%	1	4%	-	-	-	-
Other/unknown	5	14%	2	8%	3	38%	-	-
Subtotal	36	100%	26	100%	8	100%	2	100%
<b>Female adult or adolescent</b>								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	4	100%	3	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	4	100%	3	100%	1	100%	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
<b>Total</b>	<b>40</b>		<b>29</b>		<b>9</b>		<b>2</b>	

## Reported new diagnoses of HIV infection in 2019 by disease status and exposure category, Lucas County

Exposure Category <sup>b</sup>	2019 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	23	58%	17	59%	4	44%	2	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	5	13%	4	14%	1	11%	-	-
Male-to-male sexual contact & IDU	3	8%	3	10%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	3	8%	3	10%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	1	3%	-	-	1	11%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	5	13%	2	7%	3	33%	-	-
<b>Total</b>	<b>40</b>		<b>29</b>		<b>9</b>		<b>2</b>	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.

Reported new diagnoses of HIV infection in 2019 by race/ethnicity and age at diagnosis, Lucas County

2019 diagnosis of HIV infection

Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx <sup>a</sup>			White			Multi-Race		
	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	84.1	5	26%	*	2	33%	*	2	14%	*	-	-
25-29	*	-	-	*	-	-	*	-	-	*	1	17%	*	1	7%	*	-	-
30-34	*	-	-	*	-	-	*	3	16%	*	2	33%	*	3	21%	*	-	-
35-39	*	-	-	*	-	-	*	2	11%	*	-	-	*	2	14%	*	-	-
40-44	*	-	-	*	-	-	*	4	21%	*	-	-	*	-	-	*	1	100%
45-49	*	-	-	*	-	-	*	2	11%	*	-	-	*	2	14%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	1	7%	*	-	-
55-64	*	-	-	*	-	-	*	1	5%	*	1	17%	*	3	21%	*	-	-
65+	*	-	-	*	-	-	*	2	11%	*	-	-	*	-	-	*	-	-
<b>Total</b>	*	-	-	*	-	-	<b>22.6</b>	<b>19</b>		<b>19.0</b>	<b>6</b>		<b>4.8</b>	<b>14</b>		*	<b>1</b>	

Reported new diagnoses of HIV infection in 2019 by race/ethnicity and transmission category, Lucas County

2019 diagnosis of HIV infection

Transmission Category <sup>c</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	-	-	-	-	10	63%	5	83%	10	77%	1	100%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	6%	1	17%	2	15%	-	-
Heterosexual contact	-	-	-	-	1	6%	-	-	-	-	-	-
Other/unknown	-	-	-	-	4	25%	-	-	1	8%	-	-
Subtotal	-	-	-	-	16	100%	6	100%	13	100%	1	100%
<b>Female adult or adolescent</b>												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	3	100%	-	-	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	3	100%	-	-	1	100%	-	-
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	-	-	-	-	<b>19</b>		<b>6</b>		<b>14</b>		<b>1</b>	

Reported new diagnoses of HIV infection in 2019 by race/ethnicity and exposure category, Lucas County

2019 diagnosis of HIV infection

Exposure Category <sup>d</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	9	47%	4	67%	9	64%	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	4	21%	-	-	1	7%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	1	17%	2	14%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	1	5%	1	17%	1	7%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	1	5%	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	4	21%	-	-	1	7%	-	-
<b>Total</b>	-	-	-	-	<b>19</b>		<b>6</b>		<b>14</b>		<b>1</b>	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (\*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

<sup>b</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2019 U.S. Census estimates.

<sup>c</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>d</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.

**Trends in reported new diagnoses of HIV infection by year of diagnosis (2015-2019) and cumulative diagnoses by selected characteristics, Lucas County**

Characteristic	Diagnosis of HIV Infection by Year															Cumulative Diagnoses	
	2015			2016			2017			2018			2019			No.	%
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%		
<b>Sex at birth</b>																	
Males	10.9	23	82%	19.1	40	93%	16.8	35	90%	18.7	39	89%	17.4	36	90%	1,334	78%
Females	2.2	5	18%	*	3	7%	*	4	10%	2.3	5	11%	*	4	10%	368	22%
<b>Age at diagnosis (yr)</b>																	
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	23	1%
13-14	*	-	-	*	-	-	*	1	3%	*	-	-	*	-	-	3	<1%
15-19	*	3	11%	*	2	5%	*	2	5%	*	2	5%	*	-	-	69	4%
20-24	16.5	5	18%	30.9	9	21%	17.8	5	13%	36.7	10	23%	33.9	9	23%	250	15%
25-29	15.4	5	18%	30.0	10	23%	33.2	11	28%	21.2	7	16%	*	2	5%	293	17%
30-34	*	2	7%	18.2	5	12%	18.1	5	13%	28.2	8	18%	27.4	8	20%	297	17%
35-39	*	3	11%	*	4	9%	*	2	5%	*	3	7%	*	4	10%	282	17%
40-44	*	4	14%	*	3	7%	25.0	6	15%	*	3	7%	21.3	5	13%	192	11%
45-49	*	1	4%	*	3	7%	*	1	3%	18.9	5	11%	*	4	10%	127	7%
50-54	*	3	11%	*	4	9%	*	1	3%	*	1	2%	*	1	3%	80	5%
55-64	*	2	7%	*	3	7%	*	4	10%	*	4	9%	8.5	5	13%	74	4%
65+	*	-	-	*	-	-	*	1	3%	*	1	2%	*	2	5%	12	1%
<b>Race/Ethnicity<sup>b</sup></b>																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	6	<1%
Black/African-American	19.3	16	57%	26.3	22	51%	25.1	21	54%	30.8	26	59%	22.6	19	48%	734	43%
Hispanic/Latinx	*	2	7%	16.7	5	12%	*	3	8%	*	2	5%	19.0	6	15%	98	6%
White	3.3	10	36%	4.3	13	30%	4.7	14	36%	4.8	14	32%	4.8	14	35%	825	48%
Multi-Race	*	-	-	*	3	7%	*	1	3%	*	2	5%	*	1	3%	33	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	6	<1%
<b>Race/Ethnicity<sup>b</sup> and Sex at birth</b>																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	5	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Black/African-American Males	38.2	15	54%	53.0	21	49%	53.4	21	54%	60.3	24	55%	40.5	16	40%	521	31%
Black/African-American Females	*	1	4%	*	1	2%	*	-	-	*	2	5%	*	3	8%	213	13%
Hispanic/Latino Males	*	-	-	33.8	5	12%	*	3	8%	*	2	5%	38.4	6	15%	81	5%
Hispanic/Latina Females	*	2	7%	*	-	-	*	-	-	*	-	-	*	-	-	17	1%
White Males	5.4	8	29%	8.2	12	28%	6.9	10	26%	7.7	11	25%	9.1	13	33%	695	41%
White Females	*	2	7%	*	1	2%	*	4	10%	*	3	7%	*	1	3%	130	8%
Multi-Race Males	*	-	-	*	2	5%	*	1	3%	*	2	5%	*	1	3%	28	2%
Multi-Race Females	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-	5	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	6	<1%
<b>Total</b>	<b>6.5</b>	<b>28</b>		<b>9.9</b>	<b>43</b>		<b>9.1</b>	<b>39</b>		<b>10.2</b>	<b>44</b>		<b>9.3</b>	<b>40</b>		<b>1,702</b>	

Notes:  
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2015-2019) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2019.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

<sup>b</sup> Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.

**Trends in reported new diagnoses of HIV infection by year of diagnosis (2015-2019) and cumulative diagnoses by transmission category, Lucas County**

Transmission Category <sup>a</sup>	Diagnosis of HIV Infection by Year											
	2015		2016		2017		2018		2019		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	18	78%	30	75%	28	80%	31	79%	26	72%	950	72%
Injection drug use (IDU)	-	-	1	3%	-	-	-	-	-	-	76	6%
Male-to-male sexual contact and IDU	1	4%	2	5%	1	3%	4	10%	4	11%	95	7%
Heterosexual contact	1	4%	3	8%	2	6%	2	5%	1	3%	92	7%
Other/unknown	3	13%	4	10%	4	11%	2	5%	5	14%	111	8%
Subtotal	23	100%	40	100%	35	100%	39	100%	36	100%	1,324	100%
<b>Female adult or adolescent</b>												
Injection drug use	-	-	-	-	1	25%	-	-	-	-	42	12%
Heterosexual contact	4	80%	3	100%	3	75%	5	100%	4	100%	301	85%
Other/unknown	1	20%	-	-	-	-	-	-	-	-	12	3%
Subtotal	5	100%	3	100%	4	100%	5	100%	4	100%	355	100%
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	22	96%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	4%
Subtotal	-	-	-	-	-	-	-	-	-	-	23	100%
<b>Total</b>	<b>28</b>		<b>43</b>		<b>39</b>		<b>44</b>		<b>40</b>		<b>1,702</b>	

**Trends in reported diagnoses of HIV infection by year of diagnosis (2015-2019) and cumulative diagnoses by exposure category, Lucas County**

Exposure Category <sup>b</sup>	Diagnosis of HIV Infection by Year											
	2015		2016		2017		2018		2019		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	16	57%	29	67%	26	67%	28	64%	23	58%	892	52%
Injection drug use (IDU) only	-	-	1	2%	-	-	-	-	-	-	58	3%
Heterosexual contact only	5	18%	6	14%	5	13%	7	16%	5	13%	393	23%
Male-to-male sexual contact & IDU	1	4%	-	-	1	3%	3	7%	3	8%	74	4%
IDU & Heterosexual contact	-	-	-	-	1	3%	-	-	-	-	60	4%
Male-to-male sexual contact & Heterosexual contact	2	7%	1	2%	2	5%	3	7%	3	8%	58	3%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	2	5%	-	-	1	2%	1	3%	21	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	22	1%
Other/unknown	4	14%	4	9%	4	10%	2	5%	5	13%	124	7%
<b>Total</b>	<b>28</b>		<b>43</b>		<b>39</b>		<b>44</b>		<b>40</b>		<b>1,702</b>	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2015-2019) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2019.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.

# Persons Living with Diagnosed HIV Infection Reported in Lucas County

As of June 30, 2020

## Reported persons living with diagnosed HIV infection in 2019 by current disease status and selected characteristics, Lucas County

Characteristic	Living with diagnosed HIV infection			Current Disease Status			
	in 2019			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Sex at birth</b>							
Males	395.3	819	76%	408	76%	411	75%
Females	117.6	260	24%	126	24%	134	25%
<b>Age at end of year</b>							
<13	*	1	<1%	1	<1%	-	-
13-14	*	1	<1%	1	<1%	-	-
15-19	22.2	6	1%	3	1%	3	1%
20-24	158.0	42	4%	32	6%	10	2%
25-29	223.5	72	7%	52	10%	20	4%
30-34	304.8	89	8%	63	12%	26	5%
35-39	304.9	79	7%	48	9%	31	6%
40-44	480.4	113	10%	64	12%	49	9%
45-49	646.0	168	16%	68	13%	100	18%
50-54	596.3	154	14%	69	13%	85	16%
55-64	455.8	267	25%	92	17%	175	32%
65+	121.6	87	8%	41	8%	46	8%
<b>Race/Ethnicity<sup>b</sup></b>							
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander	82.0	6	1%	4	1%	2	<1%
Black/African-American	633.5	533	49%	254	48%	279	51%
Hispanic/Latinx	231.0	73	7%	32	6%	41	8%
White	146.8	429	40%	223	42%	206	38%
Multi-Race	265.0	32	3%	16	3%	16	3%
Unknown	*	5	<1%	5	1%	-	-
<b>Race/Ethnicity<sup>b</sup> and Sex at birth</b>							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Males	144.4	5	<1%	3	1%	2	<1%
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
Black/African-American Males	966.7	382	35%	185	35%	197	36%
Black/African-American Females	338.5	151	14%	69	13%	82	15%
Hispanic/Latino Males	364.8	57	5%	27	5%	30	6%
Hispanic/Latina Females	100.1	16	1%	5	1%	11	2%
White Males	241.8	344	32%	175	33%	169	31%
White Females	56.6	85	8%	48	9%	37	7%
Multi-Race Males	477.1	28	3%	15	3%	13	2%
Multi-Race Females	*	4	<1%	1	<1%	3	1%
Unknown	*	5	<1%	5	1%	-	-
<b>Total</b>	<b>251.9</b>	<b>1,079</b>		<b>534</b>		<b>545</b>	

**Notes:**

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2019. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2019, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2019 U.S. Census estimates.

<sup>b</sup>Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.



## Reported persons living with diagnosed HIV infection in 2019 by current disease status and transmission category, Lucas County

Transmission Category <sup>a</sup>	Living with diagnosed HIV infection		Current Disease Status			
	in 2019		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	597	73%	304	75%	293	72%
Injection drug use (IDU)	27	3%	14	3%	13	3%
Male-to-male sexual contact and IDU	58	7%	23	6%	35	9%
Heterosexual contact	68	8%	29	7%	39	10%
Other/unknown	64	8%	35	9%	29	7%
Subtotal	814	100%	405	100%	409	100%
<b>Female adult or adolescent</b>						
Injection drug use	19	8%	8	6%	11	9%
Heterosexual contact	222	89%	108	87%	114	90%
Other/unknown	9	4%	8	6%	1	1%
Subtotal	250	100%	124	100%	126	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	15	100%	5	100%	10	100%
Other/unknown	-	-	-	-	-	-
Subtotal	15	100%	5	100%	10	100%
<b>Total</b>	<b>1,079</b>		<b>534</b>		<b>545</b>	

## Reported persons living with diagnosed HIV infection in 2019 by current disease status and exposure category, Lucas County

Exposure Category <sup>b</sup>	Living with diagnosed HIV infection		Current Disease Status			
	in 2019		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	537	50%	269	50%	268	49%
Injection drug use (IDU) only	20	2%	12	2%	8	1%
Heterosexual contact only	290	27%	137	26%	153	28%
Male-to-male sexual contact & IDU	41	4%	18	3%	23	4%
IDU & Heterosexual contact	26	2%	10	2%	16	3%
Male-to-male sexual contact & Heterosexual contact	60	6%	35	7%	25	5%
Male-to-male sexual contact & IDU & Heterosexual contact	17	2%	5	1%	12	2%
Perinatal exposure	15	1%	5	1%	10	2%
Other/unknown	73	7%	43	8%	30	6%
<b>Total</b>	<b>1,079</b>		<b>534</b>		<b>545</b>	

### Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2019. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2019, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.

Reported persons living with diagnosed HIV infection in 2019 by race/ethnicity and age, Lucas County

Living with diagnosed HIV infection in 2019

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx <sup>a</sup>			White			Multi-Race			Unknown	
	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	No.	%
<13	*	-	-	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-	-	-
13-14	*	-	-	*	-	-	*	1	<1%	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	3	1%	*	-	-	*	2	<1%	*	1	3%	-	-
20-24	*	-	-	*	-	-	487.7	29	5%	185.2	5	7%	37.3	6	1%	*	2	6%	-	-
25-29	*	-	-	*	-	-	615.7	46	9%	217.9	6	8%	89.0	18	4%	*	2	6%	-	-
30-34	*	-	-	*	-	-	806.2	48	9%	451.2	11	15%	133.4	26	6%	*	4	13%	-	-
35-39	*	-	-	*	1	17%	842.4	42	8%	*	4	5%	169.7	30	7%	*	2	6%	-	-
40-44	*	-	-	*	-	-	1,340.0	59	11%	444.4	8	11%	262.2	43	10%	*	3	9%	-	-
45-49	*	-	-	*	4	67%	1,694.6	84	16%	609.3	9	12%	359.4	67	16%	*	4	13%	-	-
50-54	*	-	-	*	1	17%	1,630.1	72	14%	673.7	9	12%	348.6	67	16%	1,858.7	5	16%	-	-
55-64	*	-	-	*	-	-	1,218.8	119	22%	689.4	16	22%	275.0	124	29%	1,043.8	5	16%	3	60%
65+	*	1	100%	*	-	-	288.9	29	5%	256.0	5	7%	79.3	46	11%	*	4	13%	2	40%
<b>Total</b>	*	<b>1</b>		<b>82.0</b>	<b>6</b>		<b>633.5</b>	<b>533</b>		<b>231.0</b>	<b>73</b>		<b>146.8</b>	<b>429</b>		<b>265.0</b>	<b>32</b>		<b>5</b>	

Reported persons living with diagnosed HIV infection in 2019 by race/ethnicity and transmission category, Lucas County

Living with diagnosed HIV infection in 2019

Transmission Category <sup>c</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx <sup>a</sup>		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>														
Male-to-male sexual contact	-	-	3	60%	254	67%	45	79%	276	81%	19	68%	-	-
Injection drug use (IDU)	-	-	-	-	12	3%	4	7%	9	3%	2	7%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	24	6%	7	12%	23	7%	4	14%	-	-
Heterosexual contact	-	-	-	-	51	13%	-	-	14	4%	3	11%	-	-
Other/unknown	-	-	2	40%	38	10%	1	2%	20	6%	-	-	3	100%
Subtotal	-	-	5	100%	379	100%	57	100%	342	100%	28	100%	3	100%
<b>Female adult or adolescent</b>														
Injection drug use	-	-	-	-	12	8%	1	7%	6	7%	-	-	-	-
Heterosexual contact	1	100%	1	100%	127	88%	14	93%	76	92%	3	100%	-	-
Other/unknown	-	-	-	-	6	4%	-	-	1	1%	-	-	2	100%
Subtotal	1	100%	1	100%	145	100%	15	100%	83	100%	3	100%	2	100%
<b>Child (&lt;13 yrs at diagnosis)</b>														
Perinatal	-	-	-	-	9	100%	1	100%	4	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	9	100%	1	100%	4	100%	1	100%	-	-
<b>Total</b>	<b>1</b>		<b>6</b>		<b>533</b>		<b>73</b>		<b>429</b>		<b>32</b>		<b>5</b>	

Reported persons living with diagnosed HIV infection in 2019 by race/ethnicity and exposure category, Lucas County

Living with diagnosed HIV infection in 2019

Exposure Category <sup>d</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx <sup>a</sup>		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	3	50%	223	42%	42	58%	250	58%	19	59%	-	-
Injection drug use (IDU) only	-	-	-	-	10	2%	1	1%	8	2%	1	3%	-	-
Heterosexual contact only	1	100%	1	17%	178	33%	14	19%	90	21%	6	19%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	17	3%	6	8%	15	3%	3	9%	-	-
IDU & Heterosexual contact	-	-	-	-	14	3%	4	5%	7	2%	1	3%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	31	6%	3	4%	26	6%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	7	1%	1	1%	8	2%	1	3%	-	-
Perinatal exposure	-	-	-	-	9	2%	1	1%	4	1%	1	3%	-	-
Other/unknown	-	-	2	33%	44	8%	1	1%	21	5%	-	-	5	100%
<b>Total</b>	<b>1</b>		<b>6</b>		<b>533</b>		<b>73</b>		<b>429</b>		<b>32</b>		<b>5</b>	

Notes:  
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2019. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2019, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.  
 Asterisk (\*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.  
<sup>b</sup> The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.  
<sup>c</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.  
<sup>d</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
 Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.

## Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Lucas County, 2015-2019

Characteristic	Living with Diagnosed HIV Infection														
	2015			2016			2017			2018			2019		
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%
<b>Sex at birth</b>															
Males	339.6	714	74%	359.7	755	74%	373.8	780	75%	381.0	794	75%	395.3	819	76%
Females	112.8	252	26%	117.3	261	26%	117.4	261	25%	119.6	265	25%	117.6	260	24%
<b>Age at end of year</b>															
<13	*	2	<1%	*	2	<1%	*	1	<1%	*	1	<1%	*	1	<1%
13-14	*	2	<1%	*	1	<1%	*	1	<1%	*	2	<1%	*	1	<1%
15-19	42.5	12	1%	36.1	10	1%	32.8	9	1%	29.6	8	1%	22.2	6	1%
20-24	105.9	32	3%	120.3	35	3%	132.0	37	4%	146.7	40	4%	158.0	42	4%
25-29	206.1	67	7%	240.2	80	8%	274.3	91	9%	238.8	79	7%	223.5	72	7%
30-34	206.5	56	6%	233.4	64	6%	268.2	74	7%	292.8	83	8%	304.8	89	8%
35-39	373.1	93	10%	361.3	91	9%	301.4	77	7%	320.7	83	8%	304.9	79	7%
40-44	549.5	144	15%	527.7	132	13%	521.0	125	12%	511.5	121	11%	480.4	113	10%
45-49	611.9	162	17%	627.4	168	17%	610.0	163	16%	634.5	168	16%	646.0	168	16%
50-54	556.5	167	17%	573.2	165	16%	596.6	165	16%	560.0	149	14%	596.3	154	14%
55-64	320.0	190	20%	369.4	220	22%	395.6	236	23%	427.2	253	24%	455.8	267	25%
65+	60.0	39	4%	72.3	48	5%	91.0	62	6%	102.8	72	7%	121.6	87	8%
<b>Race/Ethnicity<sup>b</sup></b>															
American Indian/Alaska Native	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander	66.9	5	1%	81.2	6	1%	79.4	6	1%	79.5	6	1%	82.0	6	1%
Black/African-American	540.3	449	46%	576.1	482	47%	601.6	503	48%	617.0	520	49%	633.5	533	49%
Hispanic/Latinx	206.5	61	6%	223.6	67	7%	225.5	69	7%	230.2	72	7%	231.0	73	7%
White	138.9	419	43%	142.0	425	42%	143.7	426	41%	144.2	424	40%	146.8	429	40%
Multi-Race	238.2	26	3%	265.6	30	3%	265.5	31	3%	260.1	31	3%	265.0	32	3%
Unknown	*	5	1%	*	5	<1%	*	5	<1%	*	5	<1%	*	5	<1%
<b>Race/Ethnicity<sup>b</sup> and Sex at birth</b>															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander Males	*	4	<1%	141.4	5	<1%	139.4	5	<1%	139.5	5	<1%	144.4	5	<1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Black/African-American Males	777.7	305	32%	840.4	333	33%	899.6	354	34%	927.8	369	35%	966.7	382	35%
Black/African-American Females	328.2	144	15%	338.3	149	15%	336.6	149	14%	339.2	151	14%	338.5	151	14%
Hispanic/Latino Males	308.6	45	5%	337.5	50	5%	330.5	50	5%	335.1	52	5%	364.8	57	5%
Hispanic/Latina Females	107.0	16	2%	112.2	17	2%	122.9	19	2%	127.0	20	2%	100.1	16	1%
White Males	227.3	334	35%	233.0	340	33%	236.1	341	33%	235.9	338	32%	241.8	344	32%
White Females	54.9	85	9%	55.4	85	8%	55.9	85	8%	57.1	86	8%	56.6	85	8%
Multi-Race Males	427.9	23	2%	436.0	24	2%	476.5	27	3%	466.9	27	3%	477.1	28	3%
Multi-Race Females	*	3	<1%	103.6	6	1%	*	4	<1%	*	4	<1%	*	4	<1%
Unknown	*	5	1%	*	5	<1%	*	5	<1%	*	5	<1%	*	5	<1%
<b>Total</b>	<b>222.7</b>	<b>966</b>		<b>234.9</b>	<b>1,016</b>		<b>241.6</b>	<b>1,041</b>		<b>246.3</b>	<b>1,059</b>		<b>251.9</b>	<b>1,079</b>	

**Notes:**

Living with diagnosed HIV infection by year (2015-2019) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

<sup>b</sup> Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.

## Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Lucas County, 2015-2019

Transmission Category <sup>a</sup>	Living with Diagnosed HIV Infection									
	2015		2016		2017		2018		2019	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	506	71%	535	71%	555	71%	571	72%	597	73%
Injection drug use (IDU)	36	5%	35	5%	32	4%	27	3%	27	3%
Male-to-male sexual contact and IDU	51	7%	58	8%	59	8%	61	8%	58	7%
Heterosexual contact	67	9%	70	9%	71	9%	69	9%	68	8%
Other/unknown	51	7%	55	7%	60	8%	62	8%	64	8%
Subtotal	711	100%	753	100%	777	100%	790	100%	814	100%
<b>Female adult or adolescent</b>										
Injection drug use	20	8%	22	9%	21	8%	18	7%	19	8%
Heterosexual contact	213	88%	219	88%	220	88%	226	89%	222	89%
Other/unknown	9	4%	9	4%	9	4%	10	4%	9	4%
Subtotal	242	100%	250	100%	250	100%	254	100%	250	100%
<b>Child (&lt;13 yrs at diagnosis)</b>										
Perinatal	13	100%	13	100%	14	100%	15	100%	15	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	13	100%	13	100%	14	100%	15	100%	15	100%
<b>Total</b>	<b>966</b>		<b>1,016</b>		<b>1,041</b>		<b>1,059</b>		<b>1,079</b>	

## Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Lucas County, 2015-2019

Exposure Category <sup>b</sup>	Living with Diagnosed HIV Infection									
	2015		2016		2017		2018		2019	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	465	48%	489	48%	505	49%	520	49%	537	50%
Injection drug use (IDU) only	20	2%	20	2%	21	2%	19	2%	20	2%
Heterosexual contact only	280	29%	289	28%	291	28%	295	28%	290	27%
Male-to-male sexual contact & IDU	36	4%	40	4%	41	4%	43	4%	41	4%
IDU & Heterosexual contact	36	4%	37	4%	32	3%	26	2%	26	2%
Male-to-male sexual contact & Heterosexual contact	41	4%	46	5%	50	5%	51	5%	60	6%
Male-to-male sexual contact & IDU & Heterosexual contact	15	2%	18	2%	18	2%	18	2%	17	2%
Perinatal exposure	13	1%	13	1%	14	1%	15	1%	15	1%
Other/unknown	60	6%	64	6%	69	7%	72	7%	73	7%
<b>Total</b>	<b>966</b>		<b>1,016</b>		<b>1,041</b>		<b>1,059</b>		<b>1,079</b>	

### Notes:

Living with diagnosed HIV infection by year (2015-2019) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2020.