

New Diagnoses of HIV Infection Reported in Clermont County

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2021

Reported new diagnoses of HIV infection in 2020 by disease status and selected characteristics, Clermont County

Characteristic	2020 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	*	4	67%	2	100%	2	50%	-	-
Females	*	2	33%	-	-	2	50%	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	1	17%	1	50%	-	-	-	-
20-24	*	1	17%	1	50%	-	-	-	-
25-29	*	-	-	-	-	-	-	-	-
30-34	*	1	17%	-	-	1	25%	-	-
35-39	*	-	-	-	-	-	-	-	-
40-44	*	-	-	-	-	-	-	-	-
45-49	*	1	17%	-	-	1	25%	-	-
50-54	*	1	17%	-	-	1	25%	-	-
55-64	*	-	-	-	-	-	-	-	-
65+	*	1	17%	-	-	1	25%	-	-
Race/Ethnicity^b									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
Black/African-American	*	1	17%	1	50%	-	-	-	-
Hispanic/Latinx	*	1	17%	-	-	1	25%	-	-
White	*	4	67%	1	50%	3	75%	-	-
Multi-Race	*	-	-	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African-American Males	*	1	17%	1	50%	-	-	-	-
Black/African-American Females	*	-	-	-	-	-	-	-	-
Hispanic/Latino Males	*	1	17%	-	-	1	25%	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
White Males	*	2	33%	1	50%	1	25%	-	-
White Females	*	2	33%	-	-	2	50%	-	-
Multi-Race Males	*	-	-	-	-	-	-	-	-
Multi-Race Females	*	-	-	-	-	-	-	-	-
Total		2.9	6	2	4	4	-	-	-

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by disease status and transmission category, Clermont County

Transmission Category ^a	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	3	75%	2	100%	1	50%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	1	25%	-	-	1	50%	-	-
Subtotal	4	100%	2	100%	2	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	1	50%	-	-	1	50%	-	-
Other/unknown	1	50%	-	-	1	50%	-	-
Subtotal	2	100%	-	-	2	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	6		2		4		-	

Reported new diagnoses of HIV infection in 2020 by disease status and exposure category, Clermont County

Exposure Category ^b	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	50%	2	100%	1	25%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	1	17%	-	-	1	25%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	2	33%	-	-	2	50%	-	-
Total	6		2		4		-	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and age at diagnosis, Clermont County

2020 diagnosis of HIV infection

Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	1	25%	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	1	100%	*	-	-	*	-	-	*	-	-
25-29	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
30-34	*	-	-	*	-	-	*	-	-	*	1	100%	*	-	-	*	-	-
35-39	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	1	25%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	1	25%	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	1	25%	*	-	-
Total	*	-	-	*	-	-	*	1	-	*	1	-	*	4	-	*	-	-

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and transmission category, Clermont County

2020 diagnosis of HIV infection

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	1	100%	-	-	2	100%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	1	100%	-	-	-	-
Subtotal	-	-	-	-	1	100%	1	100%	2	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	1	50%	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	50%	-	-
Subtotal	-	-	-	-	-	-	-	-	2	100%	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	1	-	1	-	4	-	-	-

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and exposure category, Clermont County

2020 diagnosis of HIV infection

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	1	100%	-	-	2	50%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	-	-	-	-	1	25%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	1	100%	1	25%	-	-
Total	-	-	-	-	1	-	1	-	4	-	-	-

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by selected characteristics, Clermont County

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2016			2017			2018			2019			2020			No.	%
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%		
Sex at birth																	
Males	*	4	44%	*	1	100%	11.8	12	100%	9.8	10	83%	*	4	67%	158	81%
Females	4.9	5	56%	*	-	-	*	-	-	*	2	17%	*	2	33%	36	19%
Age at diagnosis (yr)																	
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	1	8%	*	1	17%	5	3%
20-24	*	1	11%	*	-	-	43.8	5	42%	*	3	25%	*	1	17%	35	18%
25-29	*	1	11%	*	-	-	*	2	17%	*	4	33%	*	-	-	35	18%
30-34	*	1	11%	*	1	100%	*	2	17%	*	-	-	*	1	17%	38	20%
35-39	*	3	33%	*	-	-	*	-	-	*	2	17%	*	-	-	26	13%
40-44	*	1	11%	*	-	-	*	-	-	*	-	-	*	-	-	19	10%
45-49	*	-	-	*	-	-	*	1	8%	*	1	8%	*	1	17%	18	9%
50-54	*	1	11%	*	-	-	*	2	17%	*	1	8%	*	1	17%	11	6%
55-64	*	1	11%	*	-	-	*	-	-	*	-	-	*	-	-	5	3%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	1	17%	2	1%
Race/Ethnicity^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	1%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	1%
Black/African-American	*	1	11%	*	-	-	*	-	-	*	1	8%	*	1	17%	16	8%
Hispanic/Latinx	*	-	-	*	-	-	*	2	17%	*	1	8%	*	1	17%	14	7%
White	3.7	7	78%	*	1	100%	5.2	10	83%	5.2	10	83%	*	4	67%	156	80%
Multi-Race	*	1	11%	*	-	-	*	-	-	*	-	-	*	-	-	5	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Race/Ethnicity^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African-American Males	*	-	-	*	-	-	*	-	-	*	1	8%	*	1	17%	13	7%
Black/African-American Females	*	1	11%	*	-	-	*	-	-	*	-	-	*	-	-	3	2%
Hispanic/Latino Males	*	-	-	*	-	-	*	2	17%	*	1	8%	*	1	17%	14	7%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
White Males	*	4	44%	*	1	100%	10.6	10	83%	8.5	8	67%	*	2	33%	126	65%
White Females	*	3	33%	*	-	-	*	-	-	*	2	17%	*	2	33%	30	15%
Multi-Race Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	1%
Multi-Race Females	*	1	11%	*	-	-	*	-	-	*	-	-	*	-	-	3	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Total	4.4	9		*	1		5.8	12		5.8	12		2.9	6		194	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by transmission category, Clermont County

Transmission Category ^a	Diagnoses of HIV Infection by Year											
	2016		2017		2018		2019		2020		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	3	75%	-	-	8	67%	4	40%	3	75%	103	65%
Injection drug use (IDU)	1	25%	-	-	1	8%	2	20%	-	-	11	7%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	1	10%	-	-	9	6%
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	7	4%
Other/unknown	-	-	1	100%	3	25%	3	30%	1	25%	28	18%
Subtotal	4	100%	1	100%	12	100%	10	100%	4	100%	158	100%
Female adult or adolescent												
Injection drug use	1	20%	-	-	-	-	2	100%	-	-	6	17%
Heterosexual contact	3	60%	-	-	-	-	-	-	1	50%	26	72%
Other/unknown	1	20%	-	-	-	-	-	-	1	50%	4	11%
Subtotal	5	100%	-	-	-	-	2	100%	2	100%	36	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	9		1		12		12		6		194	

Trends in reported diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by exposure category, Clermont County

Exposure Category ^b	Diagnoses of HIV Infection by Year											
	2016		2017		2018		2019		2020		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	33%	-	-	8	67%	4	33%	3	50%	101	52%
Injection drug use (IDU) only	2	22%	-	-	-	-	3	25%	-	-	9	5%
Heterosexual contact only	3	33%	-	-	-	-	-	-	1	17%	33	17%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	1	8%	-	-	9	5%
IDU & Heterosexual contact	-	-	-	-	1	8%	1	8%	-	-	8	4%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	2	1%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	11%	1	100%	3	25%	3	25%	2	33%	32	16%
Total	9		1		12		12		6		194	

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.
 Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Persons Living with Diagnosed HIV Infection Reported in Clermont County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Clermont County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	134.8	138	79%	72	81%	66	77%
Females	35.2	37	21%	17	19%	20	23%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	43.1	5	3%	5	6%	-	-
25-29	83.9	11	6%	8	9%	3	3%
30-34	197.9	26	15%	15	17%	11	13%
35-39	152.6	20	11%	13	15%	7	8%
40-44	149.4	19	11%	7	8%	12	14%
45-49	157.1	20	11%	11	12%	9	10%
50-54	158.8	22	13%	10	11%	12	14%
55-64	134.1	40	23%	17	19%	23	27%
65+	32.8	12	7%	3	3%	9	10%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	3	2%	3	3%	-	-
Black/African-American	504.9	18	10%	9	10%	9	10%
Hispanic/Latinx	201.4	9	5%	5	6%	4	5%
White	73.2	141	81%	71	80%	70	81%
Multi-Race	*	4	2%	1	1%	3	3%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	1%	1	1%	-	-
Asian/Pacific Islander Females	*	2	1%	2	2%	-	-
Black/African-American Males	574.7	11	6%	5	6%	6	7%
Black/African-American Females	424.0	7	4%	4	4%	3	3%
Hispanic/Latino Males	340.9	8	5%	5	6%	3	3%
Hispanic/Latina Females	*	1	1%	-	-	1	1%
White Males	121.5	115	66%	60	67%	55	64%
White Females	26.5	26	15%	11	12%	15	17%
Multi-Race Males	*	3	2%	1	1%	2	2%
Multi-Race Females	*	1	1%	-	-	1	1%
Unknown	*	-	-	-	-	-	-
Total	84.4	175		89		86	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Clermont County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	*	-	-	*	1	11%	*	4	3%	*	-	-	-	-
25-29	*	-	-	*	-	-	*	2	11%	*	-	-	75.7	9	6%	*	-	-	-	-
30-34	*	-	-	*	-	-	*	3	17%	*	1	11%	184.4	22	16%	*	-	-	-	-
35-39	*	-	-	*	-	-	2,074.7	5	28%	*	3	33%	91.2	11	8%	*	<5	-	-	-
40-44	*	-	-	*	1	33%	*	2	11%	*	-	-	126.6	15	11%	*	<5	-	-	-
45-49	*	-	-	*	-	-	*	2	11%	*	2	22%	134.3	16	11%	*	-	-	-	-
50-54	*	-	-	*	-	-	*	3	17%	*	2	22%	129.5	17	12%	*	-	-	-	-
55-64	*	-	-	*	1	33%	*	1	6%	*	-	-	129.6	37	26%	*	<5	-	-	-
65+	*	-	-	*	1	33%	*	-	-	*	-	-	28.3	10	7%	*	<5	-	-	-
Total	*	-	-	*	3	-	504.9	18	-	201.4	9	-	73.2	141	-	4	-	-	-	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Clermont County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	7	64%	3	38%	86	75%	3	100%	-	-
Injection drug use (IDU)	-	-	-	-	1	9%	-	-	5	4%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	10	9%	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	1	13%	2	2%	-	-	-	-
Other/unknown	-	-	1	100%	3	27%	4	50%	12	10%	-	-	-	-
Subtotal	-	-	1	100%	11	100%	8	100%	115	100%	3	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	-	-	-	-	6	23%	-	-	-	-
Heterosexual contact	-	-	2	100%	7	100%	-	-	16	62%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	1	100%	4	15%	-	-	-	-
Subtotal	-	-	2	100%	7	100%	1	100%	26	100%	1	100%	-	-
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	3	-	18	-	9	-	141	-	4	-	-	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Clermont County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	6	33%	3	33%	82	58%	2	50%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	6	4%	-	-	-	-
Heterosexual contact only	-	-	2	67%	7	39%	1	11%	18	13%	1	25%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	7	5%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	1	6%	-	-	5	4%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	1	6%	-	-	4	3%	1	25%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	3	2%	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	1	33%	3	17%	5	56%	16	11%	-	-	-	-
Total	-	-	3	-	18	-	9	-	141	-	4	-	-	-

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Clermont County, 2016-2020

Characteristic	Living with Diagnosed HIV Infection														
	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	104.8	105	77%	107.4	108	77%	124.3	126	78%	126.6	129	80%	134.8	138	79%
Females	30.1	31	23%	31.8	33	23%	33.6	35	22%	31.6	33	20%	35.2	37	21%
Age at end of year															
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	1	1%	*	-	-
20-24	*	3	2%	*	3	2%	43.8	5	3%	43.6	5	3%	43.1	5	3%
25-29	102.6	13	10%	77.8	10	7%	76.4	10	6%	68.3	9	6%	83.9	11	6%
30-34	118.9	15	11%	118.7	15	11%	174.4	22	14%	154.5	20	12%	197.9	26	15%
35-39	139.6	18	13%	136.1	18	13%	136.8	18	11%	167.8	22	14%	152.6	20	11%
40-44	137.5	17	13%	164.8	20	14%	153.5	19	12%	144.4	18	11%	149.4	19	11%
45-49	85.3	12	9%	101.8	14	10%	177.4	24	15%	175.7	23	14%	157.1	20	11%
50-54	203.1	30	22%	152.1	22	16%	127.3	18	11%	143.4	20	12%	158.8	22	13%
55-64	81.7	24	18%	104.4	31	22%	116.9	35	22%	113.2	34	21%	134.1	40	23%
65+	*	4	3%	24.8	8	6%	29.8	10	6%	28.6	10	6%	32.8	12	7%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	2	1%	*	3	2%	*	3	2%	*	3	2%	*	3	2%
Black/African-American	428.3	13	10%	508.3	16	11%	496.1	16	10%	352.6	12	7%	504.9	18	10%
Hispanic/Latinx	214.8	8	6%	149.8	6	4%	238.6	10	6%	234.0	10	6%	201.4	9	5%
White	57.7	110	81%	59.7	114	81%	67.2	129	80%	69.1	133	82%	73.2	141	81%
Multi-Race	*	3	2%	*	2	1%	*	3	2%	*	4	2%	*	4	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Asian/Pacific Islander Females	*	1	1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
Black/African-American Males	560.4	9	7%	670.3	11	8%	530.3	9	6%	332.0	6	4%	574.7	11	6%
Black/African-American Females	*	4	3%	331.8	5	4%	458.1	7	4%	375.9	6	4%	424.0	7	4%
Hispanic/Latino Males	411.5	8	6%	296.4	6	4%	419.4	9	6%	407.4	9	6%	340.9	8	5%
Hispanic/Latina Females	*	-	-	*	-	-	*	1	1%	*	1	1%	*	1	1%
White Males	91.7	86	63%	94.7	89	63%	111.2	105	65%	116.2	110	68%	121.5	115	66%
White Females	24.8	24	18%	25.7	25	18%	24.6	24	15%	23.5	23	14%	26.5	26	15%
Multi-Race Males	*	1	1%	*	1	1%	*	2	1%	*	3	2%	*	3	2%
Multi-Race Females	*	2	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	67.0	136		69.0	141		78.4	161		78.5	162		84.4	175	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Clermont County, 2016-2020

Transmission Category ^a	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	74	70%	70	65%	87	69%	85	66%	99	72%
Injection drug use (IDU)	2	2%	4	4%	5	4%	5	4%	6	4%
Male-to-male sexual contact and IDU	11	10%	15	14%	13	10%	15	12%	10	7%
Heterosexual contact	4	4%	3	3%	3	2%	3	2%	3	2%
Other/unknown	14	13%	16	15%	18	14%	21	16%	20	14%
Subtotal	105	100%	108	100%	126	100%	129	100%	138	100%
Female adult or adolescent										
Injection drug use	6	19%	5	15%	4	11%	6	18%	6	16%
Heterosexual contact	22	71%	25	76%	26	74%	23	70%	26	70%
Other/unknown	3	10%	3	9%	5	14%	4	12%	5	14%
Subtotal	31	100%	33	100%	35	100%	33	100%	37	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-
Total	136		141		161		162		175	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Clermont County, 2016-2020

Exposure Category ^b	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	70	51%	65	46%	82	51%	79	49%	93	53%
Injection drug use (IDU) only	5	4%	5	4%	4	2%	6	4%	6	3%
Heterosexual contact only	26	19%	28	20%	29	18%	26	16%	29	17%
Male-to-male sexual contact & IDU	9	7%	10	7%	10	6%	11	7%	7	4%
IDU & Heterosexual contact	3	2%	4	3%	5	3%	5	3%	6	3%
Male-to-male sexual contact & Heterosexual contact	4	3%	5	4%	5	3%	6	4%	6	3%
Male-to-male sexual contact & IDU & Heterosexual contact	2	1%	5	4%	3	2%	4	2%	3	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-
Other/unknown	17	13%	19	13%	23	14%	25	15%	25	14%
Total	136		141		161		162		175	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.