

New Diagnoses of HIV Infection Reported in Stark County

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2021

Reported new diagnoses of HIV infection in 2020 by disease status and selected characteristics, Stark County

Characteristic	2020 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	14.5	26	90%	22	88%	4	100%	-	-
Females	*	3	10%	3	12%	-	-	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	4	14%	4	16%	-	-	-	-
20-24	36.3	8	28%	7	28%	1	25%	-	-
25-29	21.7	5	17%	4	16%	1	25%	-	-
30-34	*	3	10%	2	8%	1	25%	-	-
35-39	*	3	10%	3	12%	-	-	-	-
40-44	*	1	3%	1	4%	-	-	-	-
45-49	*	2	7%	1	4%	1	25%	-	-
50-54	*	1	3%	1	4%	-	-	-	-
55-64	*	1	3%	1	4%	-	-	-	-
65+	*	1	3%	1	4%	-	-	-	-
Race/Ethnicity^b									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
Black/African-American	48.2	14	48%	14	56%	-	-	-	-
Hispanic/Latinx	*	-	-	-	-	-	-	-	-
White	4.7	15	52%	11	44%	4	100%	-	-
Multi-Race	*	-	-	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African-American Males	77.6	11	38%	11	44%	-	-	-	-
Black/African-American Females	*	3	10%	3	12%	-	-	-	-
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
White Males	9.7	15	52%	11	44%	4	100%	-	-
White Females	*	-	-	-	-	-	-	-	-
Multi-Race Males	*	-	-	-	-	-	-	-	-
Multi-Race Females	*	-	-	-	-	-	-	-	-
Total	7.8	29		25		4			

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by disease status and transmission category, Stark County

Transmission Category ^a	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	19	73%	16	73%	3	75%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	2	8%	2	9%	-	-	-	-
Other/unknown	5	19%	4	18%	1	25%	-	-
Subtotal	26	100%	22	100%	4	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	3	100%	3	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	3	100%	3	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	29		25		4		-	

Reported new diagnoses of HIV infection in 2020 by disease status and exposure category, Stark County

Exposure Category ^b	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	18	62%	15	60%	3	75%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	5	17%	5	20%	-	-	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	1	3%	1	4%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	5	17%	4	16%	1	25%	-	-
Total	29		25		4		-	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and age at diagnosis, Stark County

2020 diagnosis of HIV infection

Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	3	21%	*	-	-	*	1	7%	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	234.3	5	36%	*	-	-	*	3	20%	*	-	-
25-29	*	-	-	*	-	-	*	3	21%	*	-	-	*	2	13%	*	-	-
30-34	*	-	-	*	-	-	*	2	14%	*	-	-	*	1	7%	*	-	-
35-39	*	-	-	*	-	-	*	-	-	*	-	-	*	3	20%	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	1	7%	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	2	13%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	1	7%	*	-	-
55-64	*	-	-	*	-	-	*	1	7%	*	-	-	*	-	-	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	1	7%	*	-	-
Total	*	-	-	*	-	-	48.2	14		*	-	-	4.7	15		*	-	-

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and transmission category, Stark County

2020 diagnosis of HIV infection

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	5	45%	-	-	14	93%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	2	18%	-	-	-	-	-	-
Other/unknown	-	-	-	-	4	36%	-	-	1	7%	-	-
Subtotal	-	-	-	-	11	100%	-	-	15	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	3	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	3	100%	-	-	-	-	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	14		-	-	15		-	-

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and exposure category, Stark County

2020 diagnosis of HIV infection

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	5	36%	-	-	13	87%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	5	36%	-	-	-	-	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	1	7%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	4	29%	-	-	1	7%	-	-
Total	-	-	-	-	14		-	-	15		-	-

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by selected characteristics, Stark County

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2016			2017			2018			2019			2020			No.	%
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%		
Sex at birth																	
Males	15.4	28	88%	13.8	25	93%	15.5	28	90%	6.1	11	73%	14.5	26	90%	671	83%
Females	*	4	13%	*	2	7%	*	3	10%	*	4	27%	*	3	10%	136	17%
Age at diagnosis (yr)																	
<13	*	1	3%	*	-	-	*	-	-	*	-	-	*	-	-	11	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	2	6%	*	3	11%	21.3	5	16%	*	1	7%	*	4	14%	37	5%
20-24	34.5	8	25%	26.2	6	22%	39.8	9	29%	*	1	7%	36.3	8	28%	132	16%
25-29	*	4	13%	39.1	9	33%	21.4	5	16%	*	3	20%	21.7	5	17%	131	16%
30-34	42.1	9	28%	23.8	5	19%	*	2	6%	*	4	27%	*	3	10%	143	18%
35-39	*	-	-	*	1	4%	*	2	6%	*	-	-	*	3	10%	113	14%
40-44	24.0	5	16%	*	3	11%	*	2	6%	*	1	7%	*	1	3%	89	11%
45-49	*	3	9%	*	-	-	*	4	13%	*	1	7%	*	2	7%	65	8%
50-54	*	-	-	*	-	-	*	2	6%	*	2	13%	*	1	3%	40	5%
55-64	*	-	-	*	-	-	*	-	-	*	1	7%	*	1	3%	37	5%
65+	*	-	-	*	-	-	*	-	-	*	1	7%	*	1	3%	9	1%
Race/Ethnicity^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	1	3%	*	-	-	*	-	-	4	<1%
Black/African-American	24.3	7	22%	27.6	8	30%	48.3	14	45%	27.6	8	53%	48.2	14	48%	225	28%
Hispanic/Latinx	*	1	3%	*	-	-	*	-	-	*	-	-	*	-	-	14	2%
White	7.1	23	72%	5.3	17	63%	4.4	14	45%	1.9	6	40%	4.7	15	52%	543	67%
Multi-Race	*	1	3%	*	2	7%	*	2	6%	*	1	7%	*	-	-	18	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	<1%
Race/Ethnicity^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	1	3%	*	-	-	*	-	-	2	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Black/African-American Males	49.7	7	22%	49.3	7	26%	91.9	13	42%	42.4	6	40%	77.6	11	38%	168	21%
Black/African-American Females	*	-	-	*	1	4%	*	1	3%	*	2	13%	*	3	10%	57	7%
Hispanic/Latino Males	*	1	3%	*	-	-	*	-	-	*	-	-	*	-	-	14	2%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
White Males	12.1	19	59%	10.9	17	63%	7.7	12	39%	*	4	27%	9.7	15	52%	468	58%
White Females	*	4	13%	*	-	-	*	2	6%	*	2	13%	*	-	-	75	9%
Multi-Race Males	*	1	3%	*	1	4%	*	2	6%	*	1	7%	*	-	-	16	2%
Multi-Race Females	*	-	-	*	1	4%	*	-	-	*	-	-	*	-	-	2	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	<1%
Total	8.6	32		7.2	27		8.3	31		4.0	15		7.8	29		807	

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by transmission category, Stark County

Transmission Category ^a	Diagnoses of HIV Infection by Year											
	2016		2017		2018		2019		2020		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	20	71%	14	56%	22	79%	7	64%	19	73%	474	71%
Injection drug use (IDU)	-	-	3	12%	-	-	2	18%	-	-	22	3%
Male-to-male sexual contact and IDU	2	7%	1	4%	-	-	-	-	-	-	27	4%
Heterosexual contact	3	11%	1	4%	-	-	-	-	2	8%	28	4%
Other/unknown	3	11%	6	24%	6	21%	2	18%	5	19%	117	18%
Subtotal	28	100%	25	100%	28	100%	11	100%	26	100%	668	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	1	33%	-	-	-	-	18	14%
Heterosexual contact	3	100%	2	100%	2	67%	4	100%	3	100%	99	77%
Other/unknown	-	-	-	-	-	-	-	-	-	-	12	9%
Subtotal	3	100%	2	100%	3	100%	4	100%	3	100%	129	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	-	-	-	-	-	-	-	-	10	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-	10	100%
Total	32		27		31		15		29		807	

Trends in reported diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by exposure category, Stark County

Exposure Category ^b	Diagnoses of HIV Infection by Year											
	2016		2017		2018		2019		2020		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	19	59%	14	52%	20	65%	7	47%	18	62%	453	56%
Injection drug use (IDU) only	-	-	1	4%	1	3%	1	7%	-	-	21	3%
Heterosexual contact only	6	19%	3	11%	2	6%	4	27%	5	17%	127	16%
Male-to-male sexual contact & IDU	1	3%	-	-	-	-	-	-	-	-	17	2%
IDU & Heterosexual contact	-	-	2	7%	-	-	1	7%	-	-	19	2%
Male-to-male sexual contact & Heterosexual contact	1	3%	-	-	2	6%	-	-	1	3%	21	3%
Male-to-male sexual contact & IDU & Heterosexual contact	1	3%	1	4%	-	-	-	-	-	-	10	1%
Perinatal exposure	1	3%	-	-	-	-	-	-	-	-	10	1%
Other/unknown	3	9%	6	22%	6	19%	2	13%	5	17%	129	16%
Total	32		27		31		15		29		807	

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.
 Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Persons Living with Diagnosed HIV Infection Reported in Stark County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Stark County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	234.8	422	82%	241	83%	181	82%
Females	47.4	90	18%	51	17%	39	18%
Age at end of year							
<13	*	2	<1%	2	1%	-	-
13-14	*	1	<1%	1	<1%	-	-
15-19	26.5	6	1%	6	2%	-	-
20-24	99.8	22	4%	19	7%	3	1%
25-29	199.8	46	9%	34	12%	12	5%
30-34	253.4	56	11%	42	14%	14	6%
35-39	212.2	46	9%	31	11%	15	7%
40-44	188.3	39	8%	21	7%	18	8%
45-49	186.4	40	8%	21	7%	19	9%
50-54	334.8	79	15%	42	14%	37	17%
55-64	219.0	115	22%	50	17%	65	30%
65+	79.7	60	12%	23	8%	37	17%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	3	1%	2	1%	1	<1%
Black/African-American	509.7	148	29%	81	28%	67	30%
Hispanic/Latinx	141.2	12	2%	7	2%	5	2%
White	105.1	334	65%	190	65%	144	65%
Multi-Race	129.1	13	3%	10	3%	3	1%
Unknown	*	2	<1%	2	1%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	2	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
Black/African-American Males	790.6	112	22%	65	22%	47	21%
Black/African-American Females	242.1	36	7%	16	5%	20	9%
Hispanic/Latino Males	251.7	11	2%	7	2%	4	2%
Hispanic/Latina Females	*	1	<1%	-	-	1	<1%
White Males	184.3	284	55%	158	54%	126	57%
White Females	30.6	50	10%	32	11%	18	8%
Multi-Race Males	217.6	11	2%	8	3%	3	1%
Multi-Race Females	*	2	<1%	2	1%	-	-
Unknown	*	2	<1%	2	1%	-	-
Total	138.5	512		292		220	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Stark County

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	305	73%	169	70%	136	76%
Injection drug use (IDU)	10	2%	6	3%	4	2%
Male-to-male sexual contact and IDU	25	6%	13	5%	12	7%
Heterosexual contact	19	5%	10	4%	9	5%
Other/unknown	61	15%	42	18%	19	11%
Subtotal	420	100%	240	100%	180	100%
Female adult or adolescent						
Injection drug use	6	7%	2	4%	4	11%
Heterosexual contact	72	86%	41	85%	31	86%
Other/unknown	6	7%	5	10%	1	3%
Subtotal	84	100%	48	100%	36	100%
Child (<13 yrs at diagnosis)						
Perinatal	8	100%	4	100%	4	100%
Other/unknown	-	-	-	-	-	-
Subtotal	8	100%	4	100%	4	100%
Total	512		292		220	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Stark County

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	285	56%	160	55%	125	57%
Injection drug use (IDU) only	8	2%	3	1%	5	2%
Heterosexual contact only	91	18%	51	17%	40	18%
Male-to-male sexual contact & IDU	18	4%	10	3%	8	4%
IDU & Heterosexual contact	8	2%	5	2%	3	1%
Male-to-male sexual contact & Heterosexual contact	20	4%	9	3%	11	5%
Male-to-male sexual contact & IDU & Heterosexual contact	7	1%	3	1%	4	2%
Perinatal exposure	8	2%	4	1%	4	2%
Other/unknown	67	13%	47	16%	20	9%
Total	512		292		220	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Stark County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	2	1%	*	-	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	1	<1%	*	-	-	-	-
15-19	*	-	-	*	-	-	*	4	3%	*	-	-	*	2	1%	*	-	-	-	-
20-24	*	-	-	*	-	-	515.5	11	7%	*	-	-	50.3	9	3%	*	2	15%	-	-
25-29	*	-	-	*	-	-	785.0	18	12%	*	-	-	137.3	26	8%	*	2	15%	-	-
30-34	*	-	-	*	-	-	1,164.0	23	16%	*	1	8%	155.0	29	9%	*	3	23%	-	-
35-39	*	-	-	*	1	33%	544.5	9	6%	*	1	8%	186.8	35	10%	*	-	-	-	-
40-44	*	-	-	*	-	-	627.7	10	7%	*	-	-	160.6	29	9%	*	-	-	-	-
45-49	*	-	-	*	1	33%	899.7	14	9%	*	-	-	132.7	25	7%	*	-	-	-	-
50-54	*	-	-	*	1	33%	1,319.9	21	14%	*	4	33%	237.2	50	15%	*	3	23%	-	-
55-64	*	-	-	*	-	-	747.6	25	17%	*	4	33%	172.5	82	25%	*	3	23%	1	50%
65+	*	-	-	*	-	-	323.8	13	9%	*	2	17%	63.2	44	13%	*	-	-	1	50%
Total	*	-	-	*	3	-	509.7	148	-	141.2	12	-	105.1	334	-	129.1	13	-	2	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Stark County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	1	50%	77	69%	6	55%	214	76%	7	64%	-	-
Injection drug use (IDU)	-	-	-	-	5	5%	-	-	5	2%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	3	3%	3	27%	19	7%	-	-	-	-
Heterosexual contact	-	-	-	-	4	4%	1	9%	11	4%	3	27%	-	-
Other/unknown	-	-	1	50%	22	20%	1	9%	34	12%	1	9%	2	100%
Subtotal	-	-	2	100%	111	100%	11	100%	283	100%	11	100%	2	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	1	3%	1	100%	4	9%	-	-	-	-
Heterosexual contact	-	-	1	100%	33	97%	-	-	36	78%	2	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	6	13%	-	-	-	-
Subtotal	-	-	1	100%	34	100%	1	100%	46	100%	2	100%	-	-
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	3	100%	-	-	5	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	3	100%	-	-	5	100%	-	-	-	-
Total	-	-	3	-	148	-	12	-	334	-	13	-	2	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Stark County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	1	33%	74	50%	6	50%	198	59%	6	46%	-	-
Injection drug use (IDU) only	-	-	-	-	3	2%	-	-	5	1%	-	-	-	-
Heterosexual contact only	-	-	1	33%	37	25%	1	8%	47	14%	5	38%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	1	1%	2	17%	15	4%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	3	2%	1	8%	4	1%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	3	2%	-	-	16	5%	1	8%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	2	1%	1	8%	4	1%	-	-	-	-
Perinatal exposure	-	-	-	-	3	2%	-	-	5	1%	-	-	-	-
Other/unknown	-	-	1	33%	22	15%	1	8%	40	12%	1	8%	2	100%
Total	-	-	3	-	148	-	12	-	334	-	13	-	2	-

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Stark County, 2016-2020

Characteristic	Living with Diagnosed HIV Infection														
	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	210.7	382	82%	217.7	394	82%	233.7	422	83%	237.6	428	82%	234.8	422	82%
Females	44.2	85	18%	43.9	84	18%	45.5	87	17%	48.3	92	18%	47.4	90	18%
Age at end of year															
<13	*	3	1%	*	3	1%	*	3	1%	*	2	<1%	*	2	<1%
13-14	*	-	-	*	-	-	*	-	-	*	1	<1%	*	1	<1%
15-19	20.7	5	1%	*	4	1%	34.1	8	2%	*	4	1%	26.5	6	1%
20-24	99.3	23	5%	117.8	27	6%	132.5	30	6%	107.5	24	5%	99.8	22	4%
25-29	219.9	50	11%	212.8	49	10%	222.5	52	10%	222.1	52	10%	199.8	46	9%
30-34	229.3	49	10%	237.7	50	10%	252.5	53	10%	247.1	53	10%	253.4	56	11%
35-39	122.3	26	6%	129.7	28	6%	169.8	37	7%	203.0	44	8%	212.2	46	9%
40-44	143.8	30	6%	162.5	33	7%	168.3	34	7%	171.2	35	7%	188.3	39	8%
45-49	367.8	89	19%	330.3	79	17%	308.8	72	14%	266.4	60	12%	186.4	40	8%
50-54	246.7	64	14%	258.5	65	14%	283.1	69	14%	331.9	79	15%	334.8	79	15%
55-64	172.7	94	20%	187.9	102	21%	204.2	110	22%	218.2	116	22%	219.0	115	22%
65+	49.1	34	7%	53.7	38	8%	56.7	41	8%	67.7	50	10%	79.7	60	12%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	2	<1%	*	2	<1%	*	3	1%	*	3	1%	*	3	1%
Black/African-American	406.4	117	25%	425.1	123	26%	479.9	139	27%	485.7	141	27%	509.7	148	29%
Hispanic/Latinx	122.6	9	2%	104.1	8	2%	137.3	11	2%	121.2	10	2%	141.2	12	2%
White	99.6	323	69%	101.1	326	68%	104.7	336	66%	108.3	346	67%	105.1	334	65%
Multi-Race	158.4	14	3%	188.3	17	4%	193.7	18	4%	190.4	18	3%	129.1	13	3%
Unknown	*	2	<1%	*	2	<1%	*	2	<1%	*	2	<1%	*	2	<1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	<1%	*	1	<1%	*	2	<1%	*	2	<1%	*	2	<1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Black/African-American Males	624.2	88	19%	662.4	94	20%	756.1	107	21%	756.2	107	21%	790.6	112	22%
Black/African-American Females	197.4	29	6%	196.7	29	6%	216.0	32	6%	228.5	34	7%	242.1	36	7%
Hispanic/Latino Males	212.1	8	2%	179.4	7	1%	244.8	10	2%	239.1	10	2%	251.7	11	2%
Hispanic/Latina Females	*	1	<1%	*	1	<1%	*	1	<1%	*	-	-	*	1	<1%
White Males	171.7	270	58%	177.1	277	58%	183.8	286	56%	188.4	292	56%	184.3	284	55%
White Females	31.7	53	11%	29.5	49	10%	30.3	50	10%	32.8	54	10%	30.6	50	10%
Multi-Race Males	297.3	13	3%	289.2	13	3%	321.1	15	3%	315.9	15	3%	217.6	11	2%
Multi-Race Females	*	1	<1%	*	4	1%	*	3	1%	*	3	1%	*	2	<1%
Unknown	*	2	<1%	*	2	<1%	*	2	<1%	*	2	<1%	*	2	<1%
Total	125.0	467		128.3	478		137.0	509		140.3	520		138.5	512	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Stark County, 2016-2020

Transmission Category ^a	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	274	72%	282	72%	301	72%	306	72%	305	73%
Injection drug use (IDU)	9	2%	12	3%	13	3%	16	4%	10	2%
Male-to-male sexual contact and IDU	24	6%	21	5%	23	5%	24	6%	25	6%
Heterosexual contact	20	5%	21	5%	23	5%	19	4%	19	5%
Other/unknown	53	14%	56	14%	60	14%	61	14%	61	15%
Subtotal	380	100%	392	100%	420	100%	426	100%	420	100%
Female adult or adolescent										
Injection drug use	8	10%	8	10%	6	7%	6	7%	6	7%
Heterosexual contact	65	82%	65	83%	70	86%	75	87%	72	86%
Other/unknown	6	8%	5	6%	5	6%	5	6%	6	7%
Subtotal	79	100%	78	100%	81	100%	86	100%	84	100%
Child (<13 yrs at diagnosis)										
Perinatal	8	100%	8	100%	8	100%	8	100%	8	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	8	100%	8	100%	8	100%	8	100%	8	100%
Total	467		478		509		520		512	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Stark County, 2016-2020

Exposure Category ^b	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	256	55%	268	56%	286	56%	288	55%	285	56%
Injection drug use (IDU) only	9	2%	9	2%	9	2%	12	2%	8	2%
Heterosexual contact only	85	18%	86	18%	93	18%	94	18%	91	18%
Male-to-male sexual contact & IDU	14	3%	13	3%	15	3%	16	3%	18	4%
IDU & Heterosexual contact	8	2%	11	2%	10	2%	10	2%	8	2%
Male-to-male sexual contact & Heterosexual contact	18	4%	14	3%	15	3%	18	3%	20	4%
Male-to-male sexual contact & IDU & Heterosexual contact	10	2%	8	2%	8	2%	8	2%	7	1%
Perinatal exposure	9	2%	8	2%	8	2%	8	2%	8	2%
Other/unknown	58	12%	61	13%	65	13%	66	13%	67	13%
Total	467		478		509		520		512	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.