

New Diagnoses of HIV Infection Reported in Stark County

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2024

Reported new diagnoses of HIV infection in 2023 by disease status and selected characteristics, Stark County

Characteristic	2023 diagnosis of HIV infection			Disease Status						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex at birth										
Males	6.0	11	79%	5	63%	6	100%	-	-	
Females	*	3	21%	3	38%	-	-	-	-	
Age at diagnosis (years)										
0-9	*	-	-	-	-	-	-	-	-	
10-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	*	1	7%	1	13%	-	-	-	-	
25-29	*	1	7%	1	13%	-	-	-	-	
30-34	25.1	6	43%	2	25%	4	67%	-	-	
35-39	*	2	14%	2	25%	-	-	-	-	
40-44	*	2	14%	1	13%	1	17%	-	-	
45-49	*	-	-	-	-	-	-	-	-	
50-54	*	-	-	-	-	-	-	-	-	
55-64	*	2	14%	1	13%	1	17%	-	-	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b										
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
Black/African American	20.3	6	43%	3	38%	3	50%	-	-	
Hispanic/Latino	*	-	-	-	-	-	-	-	-	
White	2.5	8	57%	5	63%	3	50%	-	-	
Multi-Race	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b and Sex at birth										
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
Black/African American Males	41.8	6	43%	3	38%	3	50%	-	-	
Black/African American Females	*	-	-	-	-	-	-	-	-	
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
White Males	3.2	5	36%	2	25%	3	50%	-	-	
White Females	*	3	21%	3	38%	-	-	-	-	
Multi-Race Males	*	-	-	-	-	-	-	-	-	
Multi-Race Females	*	-	-	-	-	-	-	-	-	
Total	3.8	14		8		6		-		

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported new diagnoses of HIV infection in 2023 by disease status and transmission category, Stark County

Transmission Category ^a	2023 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	8	73%	4	80%	4	67%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	2	18%	1	20%	1	17%	-	-
Heterosexual contact	1	9%	-	-	1	17%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	11	100%	5	100%	6	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	3	100%	3	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	3	100%	3	100%	-	-	-	-
Child (<13 years at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	14		8		6		-	

Reported new diagnoses of HIV infection in 2023 by disease status and exposure category, Stark County

Exposure Category ^b	2023 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	4	29%	2	25%	2	33%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	4	29%	3	38%	1	17%	-	-
Male-to-male sexual contact & IDU	1	7%	1	13%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	4	29%	2	25%	2	33%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	1	7%	-	-	1	17%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Total	14		8		6		-	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and age at diagnosis, Stark County

2023 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	1	17%	*	-	-	*	-	-	*	-	-
25-29	*	-	-	*	-	-	*	1	17%	*	-	-	*	-	-	*	-	-
30-34	*	-	-	*	-	-	*	3	50%	*	-	-	*	3	38%	*	-	-
35-39	*	-	-	*	-	-	*	1	17%	*	-	-	*	1	13%	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	2	25%	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	2	25%	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	20.3	6		*	-	-	2.5	8		*	-	-

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and transmission category, Stark County

Transmission Category ^c	2023 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	6	100%	-	-	2	40%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	2	40%	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	1	20%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	6	100%	-	-	5	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	3	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	3	100%	-	-
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	6	-	-	-	8	-	-	-

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and exposure category, Stark County

Exposure Category ^d	2023 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	3	50%	-	-	1	13%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	-	-	-	-	4	50%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	1	13%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	3	50%	-	-	1	13%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	13%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	6		-		8		-	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by selected characteristics, Stark County

Diagnoses of HIV Infection by Year																	
Characteristic	2019			2020			2021			2022			2023			Cumulative Diagnoses	
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex at birth																	
Males	6.1	11	73%	13.4	24	89%	9.3	17	74%	11.5	21	78%	6.0	11	79%	719	83%
Females	*	4	27%	*	3	11%	3.1	6	26%	3.2	6	22%	*	3	21%	150	17%
Age at diagnosis (years)																	
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	11	1%
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	1	7%	*	4	15%	*	-	-	*	-	-	*	-	-	37	4%
20-24	*	1	7%	36.3	8	30%	*	3	13%	21.7	5	19%	*	1	7%	141	16%
25-29	*	3	20%	*	4	15%	22.3	5	22%	*	3	11%	*	1	7%	139	16%
30-34	*	4	27%	*	3	11%	30.5	7	30%	*	4	15%	25.1	6	43%	158	18%
35-39	*	-	-	*	3	11%	*	-	-	*	3	11%	*	2	14%	118	14%
40-44	*	1	7%	*	1	4%	*	2	9%	*	3	11%	*	2	14%	96	11%
45-49	*	1	7%	*	1	4%	*	2	9%	24.6	5	19%	*	-	-	72	8%
50-54	*	2	13%	*	1	4%	*	1	4%	*	1	4%	*	-	-	42	5%
55-64	*	1	7%	*	1	4%	*	3	13%	*	2	7%	*	2	14%	45	5%
65+	*	1	7%	*	1	4%	*	-	-	*	1	4%	*	-	-	10	1%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	<1%
Black/African American	27.6	8	53%	41.3	12	44%	20.3	6	26%	33.8	10	37%	20.3	6	43%	241	28%
Hispanic/Latino	*	-	-	*	1	4%	*	1	4%	*	3	11%	*	-	-	20	2%
White	1.9	6	40%	4.4	14	52%	4.7	15	65%	4.4	14	52%	2.5	8	57%	579	67%
Multi-Race	*	1	7%	*	-	-	*	1	4%	*	-	-	*	-	-	24	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Race/Ethnicity ^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Black/African American Males	42.4	6	40%	63.5	9	33%	*	3	13%	48.5	7	26%	41.8	6	43%	181	21%
Black/African American Females	*	2	13%	*	3	11%	*	3	13%	*	3	11%	*	-	-	60	7%
Hispanic/Latino Males	*	-	-	*	1	4%	*	1	4%	*	3	11%	*	-	-	20	2%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
White Males	*	4	27%	9.1	14	52%	8.3	13	57%	7.1	11	41%	3.2	5	36%	496	57%
White Females	*	2	13%	*	-	-	*	2	9%	*	3	11%	*	3	21%	83	10%
Multi-Race Males	*	1	7%	*	-	-	*	-	-	*	-	-	*	-	-	19	2%
Multi-Race Females	*	-	-	*	-	-	*	1	4%	*	-	-	*	-	-	5	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Total	4.0	15		7.3	27		6.2	23		7.2	27		3.8	14		869	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2019-2023) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2023.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by transmission category, Stark County

Transmission Category ^a	Diagnoses of HIV Infection by Year											
	2019		2020		2021		2022		2023		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	7	64%	18	75%	9	53%	8	38%	8	73%	501	70%
Injection drug use (IDU)	2	18%	-	-	-	-	2	10%	-	-	23	3%
Male-to-male sexual contact and IDU	-	-	-	-	1	6%	2	10%	2	18%	32	4%
Heterosexual contact	-	-	2	8%	2	12%	5	24%	1	9%	36	5%
Other/unknown	2	18%	4	17%	5	29%	4	19%	-	-	124	17%
Subtotal	11	100%	24	100%	17	100%	21	100%	11	100%	716	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	18	13%
Heterosexual contact	4	100%	3	100%	6	100%	6	100%	3	100%	113	79%
Other/unknown	-	-	-	-	-	-	-	-	-	-	12	8%
Subtotal	4	100%	3	100%	6	100%	6	100%	3	100%	143	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	10	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	10	100%
Total	15		27		23		27		14		869	

Trends in reported diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by exposure category, Stark County

Exposure Category ^b	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2019		2020		2021		2022		2023			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	5	33%	11	41%	8	35%	6	22%	4	29%	309	36%
Injection drug use (IDU) only	1	7%	-	-	-	-	1	4%	-	-	9	1%
Heterosexual contact only	6	40%	6	22%	9	39%	12	44%	4	29%	197	23%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	2	7%	1	7%	12	1%
IDU & Heterosexual contact	1	7%	-	-	-	-	1	4%	-	-	32	4%
Male-to-male sexual contact & Heterosexual contact	2	13%	7	26%	1	4%	2	7%	4	29%	191	22%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	1	4%	-	-	1	7%	20	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	10	1%
Other/unknown	-	-	3	11%	4	17%	3	11%	-	-	89	10%
Total	15		27		23		27		14		869	

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2019-2023) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2023.
 Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Persons Living with Diagnosed HIV Infection Reported in Stark County

As of June 30, 2024

Reported persons living with diagnosed HIV infection in 2023 by current disease status and selected characteristics, Stark County

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2023			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex at birth							
Males	236.0	430	80%	251	78%	179	82%
Females	57.7	110	20%	71	22%	39	18%
Age at end of year (years)							
0-9	*	1	<1%	1	<1%	-	-
10-14	*	-	-	-	-	-	-
15-19	*	3	1%	3	1%	-	-
20-24	86.8	19	4%	17	5%	2	1%
25-29	177.6	39	7%	30	9%	9	4%
30-34	280.4	67	12%	47	15%	20	9%
35-39	236.0	52	10%	35	11%	17	8%
40-44	216.8	49	9%	35	11%	14	6%
45-49	200.3	41	8%	22	7%	19	9%
50-54	331.0	77	14%	45	14%	32	15%
55-64	237.7	119	22%	49	15%	70	32%
65+	93.4	73	14%	38	12%	35	16%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	127.8	5	1%	5	2%	-	-
Black/African American	537.6	159	29%	91	28%	68	31%
Hispanic/Latino	180.9	21	4%	11	3%	10	5%
White	105.5	333	62%	203	63%	130	60%
Multi-Race	187.0	21	4%	11	3%	10	5%
Unknown	*	1	<1%	1	<1%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	3	1%	3	1%	-	-
Asian/Pacific Islander Females	*	2	<1%	2	1%	-	-
Black/African American Males	802.1	115	21%	65	20%	50	23%
Black/African American Females	288.7	44	8%	26	8%	18	8%
Hispanic/Latino Males	300.7	18	3%	10	3%	8	4%
Hispanic/Latina Females	*	3	1%	1	<1%	2	1%
White Males	181.1	279	52%	164	51%	115	53%
White Females	33.4	54	10%	39	12%	15	7%
Multi-Race Males	250.1	14	3%	8	2%	6	3%
Multi-Race Females	124.4	7	1%	3	1%	4	2%
Unknown	*	1	<1%	1	<1%	-	-
Total	144.9	540		322		218	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported persons living with diagnosed HIV infection in 2023 by current disease status and transmission category, Stark County

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2023		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	304	71%	170	68%	134	76%
Injection drug use (IDU)	10	2%	6	2%	4	2%
Male-to-male sexual contact and IDU	28	7%	17	7%	11	6%
Heterosexual contact	21	5%	14	6%	7	4%
Other/unknown	63	15%	42	17%	21	12%
Subtotal	426	100%	249	100%	177	100%
Female adult or adolescent						
Injection drug use	7	7%	2	3%	5	13%
Heterosexual contact	95	90%	62	91%	33	87%
Other/unknown	4	4%	4	6%	-	-
Subtotal	106	100%	68	100%	38	100%
Child (<13 years at diagnosis)						
Perinatal	8	100%	5	100%	3	100%
Other/unknown	-	-	-	-	-	-
Subtotal	8	100%	5	100%	3	100%
Total	540		322		218	

Reported persons living with diagnosed HIV infection in 2023 by current disease status and exposure category, Stark County

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2023		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	192	36%	114	35%	78	36%
Injection drug use (IDU) only	4	1%	3	1%	1	<1%
Heterosexual contact only	145	27%	91	28%	54	25%
Male-to-male sexual contact & IDU	12	2%	10	3%	2	1%
IDU & Heterosexual contact	13	2%	5	2%	8	4%
Male-to-male sexual contact & Heterosexual contact	111	21%	56	17%	55	25%
Male-to-male sexual contact & IDU & Heterosexual contact	16	3%	7	2%	9	4%
Perinatal exposure	8	1%	5	2%	3	1%
Other/unknown	39	7%	31	10%	8	4%
Total	540		322		218	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and age, Stark County

Living with diagnosed HIV infection in 2023																		
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	-	-	*	1	20%	*	-	-	*	-	-	*	-	-	*	-	-
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	3	1%	*	-	-
20-24	*	-	-	*	-	-	664.5	14	9%	*	-	-	28.9	5	2%	*	-	-
25-29	*	-	-	*	-	-	881.5	18	11%	*	2	10%	89.8	16	5%	*	3	14%
30-34	*	-	-	*	1	20%	851.1	20	13%	*	-	-	208.8	41	12%	726.7	5	24%
35-39	*	-	-	*	-	-	972.5	17	11%	*	1	5%	170.8	32	10%	*	2	10%
40-44	*	-	-	*	2	40%	567.5	10	6%	*	1	5%	175.3	34	10%	*	2	10%
45-49	*	-	-	*	-	-	658.8	10	6%	*	2	10%	152.4	27	8%	*	2	10%
50-54	*	-	-	*	1	20%	1,021.1	16	10%	*	4	19%	272.7	56	17%	*	-	-
55-64	*	-	-	*	-	-	1,153.5	36	23%	781.3	7	33%	155.5	70	21%	1,749.3	6	29%
65+	*	-	-	*	-	-	430.2	18	11%	*	4	19%	68.3	49	15%	*	1	5%
Total	*	-	-	127.8	5	-	537.6	159	-	180.9	21	-	105.5	333	-	187.0	21	-

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and transmission category, Stark County

Living with diagnosed HIV infection in 2023														
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	1	50%	78	69%	13	72%	202	73%	10	71%	-	-
Injection drug use (IDU)	-	-	-	-	4	4%	-	-	5	2%	1	7%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	2	2%	3	17%	22	8%	1	7%	-	-
Heterosexual contact	-	-	-	-	8	7%	-	-	11	4%	2	14%	-	-
Other/unknown	-	-	1	50%	21	19%	2	11%	38	14%	-	-	1	100%
Subtotal	-	-	2	100%	113	100%	18	100%	278	100%	14	100%	1	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	2	5%	1	33%	3	6%	1	14%	-	-
Heterosexual contact	-	-	2	100%	41	95%	2	67%	44	86%	6	86%	-	-
Other/unknown	-	-	-	-	-	-	-	-	4	8%	-	-	-	-
Subtotal	-	-	2	100%	43	100%	3	100%	51	100%	7	100%	-	-
Child (<13 years at diagnosis)														
Perinatal	-	-	1	100%	3	100%	-	-	4	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	3	100%	-	-	4	100%	-	-	-	-
Total	-	-	5		159		21		333		21		1	

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and exposure category, Stark County

Living with diagnosed HIV infection in 2023														
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/ Latino ^a		White		Multi-Race		Unknown	
Exposure Category ^d	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	1	20%	45	28%	11	52%	130	39%	5	24%	-	-
Injection drug use (IDU) only	-	-	-	-	2	1%	-	-	2	1%	-	-	-	-
Heterosexual contact only	-	-	3	60%	58	36%	4	19%	72	22%	8	38%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	11	3%	1	5%	-	-
IDU & Heterosexual contact	-	-	-	-	4	3%	1	5%	6	2%	2	10%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	33	21%	2	10%	71	21%	5	24%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	2	1%	3	14%	11	3%	-	-	-	-
Perinatal exposure	-	-	1	20%	3	2%	-	-	4	1%	-	-	-	-
Other/unknown	-	-	-	-	12	8%	-	-	26	8%	-	-	1	100%
Total	-	-	5	-	159	-	21	-	333	-	21	-	1	-

Notes:
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Stark County, 2019-2023

Characteristic	Living with Diagnosed HIV Infection														
	2019			2020			2021			2022			2023		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	234.3	422	82%	237.0	426	82%	233.4	427	81%	229.9	420	79%	236.0	430	80%
Females	48.3	92	18%	47.9	91	18%	52.9	101	19%	58.4	111	21%	57.7	110	20%
Age at end of year (years)															
0-9	*	-	-	*	-	-	*	-	-	*	1	<1%	*	1	<1%
10-14	*	3	1%	*	3	1%	*	2	<1%	*	2	<1%	*	-	-
15-19	*	4	1%	22.1	5	1%	25.7	6	1%	*	3	1%	*	3	1%
20-24	107.5	24	5%	108.9	24	5%	85.1	19	4%	86.7	20	4%	86.8	19	4%
25-29	222.1	52	10%	204.1	47	9%	209.7	47	9%	193.9	43	8%	177.6	39	7%
30-34	247.1	53	10%	262.4	58	11%	283.3	65	12%	260.3	61	11%	280.4	67	12%
35-39	203.0	44	9%	230.7	50	10%	217.2	48	9%	242.0	53	10%	236.0	52	10%
40-44	166.3	34	7%	188.3	39	8%	159.7	35	7%	161.6	36	7%	216.8	49	9%
45-49	266.4	60	12%	200.4	43	8%	186.4	39	7%	226.4	46	9%	200.3	41	8%
50-54	323.5	77	15%	330.6	78	15%	314.1	76	14%	289.7	69	13%	331.0	77	14%
55-64	216.3	115	22%	213.3	112	22%	237.1	125	24%	245.0	125	24%	237.7	119	22%
65+	65.0	48	9%	77.0	58	11%	87.7	66	13%	93.7	72	14%	93.4	73	14%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	2	<1%	*	2	<1%	*	3	1%	*	4	1%	127.8	5	1%
Black/African American	468.5	136	26%	496.0	144	28%	492.8	146	28%	513.5	152	29%	537.6	159	29%
Hispanic/Latino	145.4	12	2%	188.3	16	3%	154.2	14	3%	219.9	21	4%	180.9	21	4%
White	105.8	338	66%	103.6	329	64%	107.8	345	65%	105.4	335	63%	105.5	333	62%
Multi-Race	264.4	25	5%	248.2	25	5%	181.2	19	4%	165.2	18	3%	187.0	21	4%
Unknown	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	<1%	*	1	<1%	*	1	<1%	*	2	<1%	*	3	1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	2	<1%	*	2	<1%	*	2	<1%
Black/African American Males	735.0	104	20%	776.5	110	21%	744.5	108	20%	755.3	109	21%	802.1	115	21%
Black/African American Females	215.0	32	6%	228.7	34	7%	251.3	38	7%	283.5	43	8%	288.7	44	8%
Hispanic/Latino Males	286.9	12	2%	343.2	15	3%	256.1	12	2%	385.6	19	4%	300.7	18	3%
Hispanic/Latina Females	*	-	-	*	1	<1%	*	2	<1%	*	2	<1%	*	3	1%
White Males	183.2	284	55%	181.1	279	54%	186.1	291	55%	178.0	277	52%	181.1	279	52%
White Females	32.8	54	11%	30.6	50	10%	33.0	54	10%	35.7	58	11%	33.4	54	10%
Multi-Race Males	421.1	20	4%	395.6	20	4%	265.6	14	3%	219.0	12	2%	250.1	14	3%
Multi-Race Females	106.2	5	1%	99.7	5	1%	95.9	5	1%	110.8	6	1%	124.4	7	1%
Unknown	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Total	138.7	514		139.8	517		141.2	528		142.5	531		144.9	540	

Notes:

Living with diagnosed HIV infection by year (2019-2023) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Stark County, 2019-2023

Living with Diagnosed HIV Infection										
Transmission Category ^a	2019		2020		2021		2022		2023	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	303	72%	303	71%	309	73%	296	71%	304	71%
Injection drug use (IDU)	16	4%	12	3%	12	3%	11	3%	10	2%
Male-to-male sexual contact and IDU	24	6%	27	6%	27	6%	29	7%	28	7%
Heterosexual contact	18	4%	20	5%	16	4%	21	5%	21	5%
Other/unknown	59	14%	62	15%	61	14%	60	14%	63	15%
Subtotal	420	100%	424	100%	425	100%	417	100%	426	100%
Female adult or adolescent										
Injection drug use	7	8%	7	8%	7	7%	6	6%	7	7%
Heterosexual contact	74	86%	73	86%	82	86%	95	90%	95	90%
Other/unknown	5	6%	5	6%	6	6%	5	5%	4	4%
Subtotal	86	100%	85	100%	95	100%	106	100%	106	100%
Child (<13 years at diagnosis)										
Perinatal	8	100%	8	100%	8	100%	8	100%	8	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	8	100%	8	100%	8	100%	8	100%	8	100%
Total	514		517		528		531		540	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Stark County, 2019-2023

Living with Diagnosed HIV Infection										
Exposure Category ^b	2019		2020		2021		2022		2023	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	187	36%	185	36%	193	37%	187	35%	192	36%
Injection drug use (IDU) only	5	1%	4	1%	4	1%	4	1%	4	1%
Heterosexual contact only	121	24%	120	23%	124	23%	141	27%	145	27%
Male-to-male sexual contact & IDU	7	1%	9	2%	7	1%	12	2%	12	2%
IDU & Heterosexual contact	18	4%	15	3%	15	3%	13	2%	13	2%
Male-to-male sexual contact & Heterosexual contact	115	22%	117	23%	115	22%	108	20%	111	21%
Male-to-male sexual contact & IDU & Heterosexual contact	17	3%	18	3%	20	4%	17	3%	16	3%
Perinatal exposure	8	2%	8	2%	8	2%	8	2%	8	1%
Other/unknown	36	7%	41	8%	42	8%	41	8%	39	7%
Total	514		517		528		531		540	

Notes:

Living with diagnosed HIV infection by year (2019-2023) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.