

New Diagnoses of HIV Infection Reported in Ohio

As of June 30, 2021

Reported new diagnoses of HIV infection in 2020 by disease status and selected characteristics, Ohio

| Characteristic | 2020 diagnosis of HIV infection | | | Disease Status | | | | | |
|----------------------------------------------------|---------------------------------|------------|-----|----------------|-----|------------------|-----|-----------|-----|
| | Rate ^a | No. | % | HIV (not AIDS) | | HIV & later AIDS | | AIDS | |
| | | | | No. | % | No. | % | No. | % |
| Sex at birth | | | | | | | | | |
| Males | 12.9 | 737 | 82% | 568 | 82% | 136 | 82% | 33 | 83% |
| Females | 2.7 | 160 | 18% | 124 | 18% | 29 | 18% | 7 | 18% |
| Age at diagnosis (yr) | | | | | | | | | |
| <13 | * | 2 | <1% | 1 | <1% | - | - | 1 | 3% |
| 13-14 | * | - | - | - | - | - | - | - | - |
| 15-19 | 5.5 | 41 | 5% | 39 | 6% | 2 | 1% | - | - |
| 20-24 | 21.8 | 163 | 18% | 146 | 21% | 14 | 8% | 3 | 8% |
| 25-29 | 21.4 | 170 | 19% | 144 | 21% | 20 | 12% | 6 | 15% |
| 30-34 | 19.0 | 146 | 16% | 112 | 16% | 25 | 15% | 9 | 23% |
| 35-39 | 13.0 | 94 | 10% | 69 | 10% | 21 | 13% | 4 | 10% |
| 40-44 | 9.2 | 63 | 7% | 51 | 7% | 8 | 5% | 4 | 10% |
| 45-49 | 9.4 | 65 | 7% | 46 | 7% | 18 | 11% | 1 | 3% |
| 50-54 | 8.5 | 62 | 7% | 34 | 5% | 20 | 12% | 8 | 20% |
| 55-64 | 4.6 | 73 | 8% | 42 | 6% | 28 | 17% | 3 | 8% |
| 65+ | 0.9 | 18 | 2% | 8 | 1% | 9 | 5% | 1 | 3% |
| Race/Ethnicity^b | | | | | | | | | |
| American Indian/Alaska Native | * | 2 | <1% | 2 | <1% | - | - | - | - |
| Asian/Pacific Islander | 2.3 | 7 | 1% | 5 | 1% | 1 | 1% | 1 | 3% |
| Black/African-American | 31.6 | 470 | 52% | 391 | 57% | 63 | 38% | 16 | 40% |
| Hispanic/Latinx | 6.8 | 33 | 4% | 24 | 3% | 8 | 5% | 1 | 3% |
| White | 4.0 | 361 | 40% | 256 | 37% | 84 | 51% | 21 | 53% |
| Multi-Race | 9.1 | 24 | 3% | 14 | 2% | 9 | 5% | 1 | 3% |
| Race/Ethnicity^b and Sex at birth | | | | | | | | | |
| American Indian/Alaska Native Males | * | 2 | <1% | 2 | <1% | - | - | - | - |
| American Indian/Alaska Native Females | * | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander Males | 4.1 | 6 | 1% | 4 | 1% | 1 | 1% | 1 | 3% |
| Asian/Pacific Islander Females | * | 1 | <1% | 1 | <1% | - | - | - | - |
| Black/African-American Males | 52.9 | 377 | 42% | 317 | 46% | 48 | 29% | 12 | 30% |
| Black/African-American Females | 12.0 | 93 | 10% | 74 | 11% | 15 | 9% | 4 | 10% |
| Hispanic/Latino Males | 12.4 | 31 | 3% | 22 | 3% | 8 | 5% | 1 | 3% |
| Hispanic/Latina Females | * | 2 | <1% | 2 | <1% | - | - | - | - |
| White Males | 6.8 | 303 | 34% | 213 | 31% | 72 | 44% | 18 | 45% |
| White Females | 1.2 | 58 | 6% | 43 | 6% | 12 | 7% | 3 | 8% |
| Multi-Race Males | 13.7 | 18 | 2% | 10 | 1% | 7 | 4% | 1 | 3% |
| Multi-Race Females | 4.5 | 6 | 1% | 4 | 1% | 2 | 1% | - | - |
| Total | 7.7 | 897 | | 692 | | 165 | | 40 | |

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by disease status and transmission category, Ohio

| Transmission Category ^a | 2020 diagnosis of HIV infection | | Disease Status | | | | | |
|----------------------------------------|---------------------------------|------|----------------|------|------------------|------|-----------|------|
| | | | HIV (not AIDS) | | HIV & later AIDS | | AIDS | |
| | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | |
| Male-to-male sexual contact | 400 | 54% | 317 | 56% | 69 | 51% | 14 | 44% |
| Injection drug use (IDU) | 34 | 5% | 29 | 5% | 4 | 3% | 1 | 3% |
| Male-to-male sexual contact and IDU | 31 | 4% | 27 | 5% | 2 | 1% | 2 | 6% |
| Heterosexual contact | 26 | 4% | 21 | 4% | 4 | 3% | 1 | 3% |
| Other/unknown | 244 | 33% | 173 | 31% | 57 | 42% | 14 | 44% |
| Subtotal | 735 | 100% | 567 | 100% | 136 | 100% | 32 | 100% |
| Female adult or adolescent | | | | | | | | |
| Injection drug use | 21 | 13% | 18 | 15% | 2 | 7% | 1 | 14% |
| Heterosexual contact | 110 | 69% | 85 | 69% | 21 | 72% | 4 | 57% |
| Other/unknown | 29 | 18% | 21 | 17% | 6 | 21% | 2 | 29% |
| Subtotal | 160 | 100% | 124 | 100% | 29 | 100% | 7 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | |
| Perinatal | 2 | 100% | 1 | 100% | - | - | 1 | 100% |
| Other/unknown | - | - | - | - | - | - | - | - |
| Subtotal | 2 | 100% | 1 | 100% | - | - | 1 | 100% |
| Total | 897 | | 692 | | 165 | | 40 | |

Reported new diagnoses of HIV infection in 2020 by disease status and exposure category, Ohio

| Exposure Category ^b | 2020 diagnosis of HIV infection | | Disease Status | | | | | |
|----------------------------------------------------------|---------------------------------|-----|----------------|-----|------------------|-----|-----------|-----|
| | | | HIV (not AIDS) | | HIV & later AIDS | | AIDS | |
| | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sexual contact only | 391 | 44% | 310 | 45% | 68 | 41% | 13 | 33% |
| Injection drug use (IDU) only | 29 | 3% | 23 | 3% | 4 | 2% | 2 | 5% |
| Heterosexual contact only | 136 | 15% | 106 | 15% | 25 | 15% | 5 | 13% |
| Male-to-male sexual contact & IDU | 26 | 3% | 23 | 3% | 2 | 1% | 1 | 3% |
| IDU & Heterosexual contact | 26 | 3% | 24 | 3% | 2 | 1% | - | - |
| Male-to-male sexual contact & Heterosexual contact | 9 | 1% | 7 | 1% | 1 | 1% | 1 | 3% |
| Male-to-male sexual contact & IDU & Heterosexual contact | 5 | 1% | 4 | 1% | - | - | 1 | 3% |
| Perinatal exposure | 2 | <1% | 1 | <1% | - | - | 1 | 3% |
| Other/unknown | 273 | 30% | 194 | 28% | 63 | 38% | 16 | 40% |
| Total | 897 | | 692 | | 165 | | 40 | |

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and age at diagnosis, Ohio

2020 diagnosis of HIV infection

| Age at diagnosis (yr) | American Indian/Alaska Native | | | Asian/Pacific Islander | | | Black/African-American | | | Hispanic/ Latinx ^a | | | White | | | Multi-Race | | |
|-----------------------|-------------------------------|----------|-----|------------------------|----------|-----|------------------------|------------|-----|-------------------------------|-----------|-----|-------------------|------------|-----|-------------------|-----------|-----|
| | Rate ^b | No. | % | Rate ^b | No. | % | Rate ^b | No. | % | Rate ^b | No. | % | Rate ^b | No. | % | Rate ^b | No. | % |
| <13 | * | - | - | * | - | - | * | 1 | <1% | * | 1 | 3% | * | - | - | * | - | - |
| 13-14 | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - |
| 15-19 | * | - | - | * | 1 | 14% | 28.6 | 30 | 6% | * | 2 | 6% | 1.5 | 8 | 2% | * | - | - |
| 20-24 | * | - | - | * | - | - | 109.7 | 118 | 25% | 19.9 | 8 | 24% | 6.0 | 33 | 9% | * | 4 | 17% |
| 25-29 | * | - | - | * | 2 | 29% | 76.1 | 95 | 20% | 15.2 | 6 | 18% | 10.0 | 58 | 16% | 42.4 | 9 | 38% |
| 30-34 | * | 1 | 50% | * | 2 | 29% | 77.6 | 85 | 18% | * | 3 | 9% | 8.9 | 51 | 14% | * | 4 | 17% |
| 35-39 | * | - | - | * | - | - | 46.9 | 44 | 9% | * | 3 | 9% | 8.1 | 45 | 12% | * | 2 | 8% |
| 40-44 | * | - | - | * | - | - | 24.3 | 21 | 4% | 18.7 | 6 | 18% | 6.4 | 34 | 9% | * | 2 | 8% |
| 45-49 | * | 1 | 50% | * | 1 | 14% | 29.3 | 25 | 5% | * | 1 | 3% | 6.7 | 37 | 10% | * | - | - |
| 50-54 | * | - | - | * | - | - | 29.8 | 25 | 5% | * | 2 | 6% | 5.9 | 35 | 10% | * | - | - |
| 55-64 | * | - | - | * | - | - | 11.5 | 20 | 4% | * | 1 | 3% | 3.7 | 50 | 14% | * | 2 | 8% |
| 65+ | * | - | - | * | 1 | 14% | 3.1 | 6 | 1% | * | - | - | 0.5 | 10 | 3% | * | 1 | 4% |
| Total | * | 2 | | 2.3 | 7 | | 31.6 | 470 | | 6.8 | 33 | | 4.0 | 361 | | 9.1 | 24 | |

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and transmission category, Ohio

2020 diagnosis of HIV infection

| Transmission Category ^c | American Indian/Alaska Native | | Asian/Pacific Islander | | Black/African-American | | Hispanic/ Latinx ^a | | White | | Multi-Race | |
|----------------------------------------|-------------------------------|------|------------------------|------|------------------------|------|-------------------------------|------|------------|------|------------|------|
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 1 | 50% | 3 | 50% | 216 | 57% | 17 | 57% | 150 | 50% | 13 | 72% |
| Injection drug use (IDU) | - | - | - | - | 3 | 1% | - | - | 30 | 10% | 1 | 6% |
| Male-to-male sexual contact and IDU | - | - | - | - | 4 | 1% | 1 | 3% | 25 | 8% | 1 | 6% |
| Heterosexual contact | - | - | - | - | 18 | 5% | 1 | 3% | 7 | 2% | - | - |
| Other/unknown | 1 | 50% | 3 | 50% | 135 | 36% | 11 | 37% | 91 | 30% | 3 | 17% |
| Subtotal | 2 | 100% | 6 | 100% | 376 | 100% | 30 | 100% | 303 | 100% | 18 | 100% |
| Female adult or adolescent | | | | | | | | | | | | |
| Injection drug use | - | - | - | - | 3 | 3% | - | - | 16 | 28% | 2 | 33% |
| Heterosexual contact | - | - | - | - | 77 | 83% | 1 | 50% | 29 | 50% | 3 | 50% |
| Other/unknown | - | - | 1 | 100% | 13 | 14% | 1 | 50% | 13 | 22% | 1 | 17% |
| Subtotal | - | - | 1 | 100% | 93 | 100% | 2 | 100% | 58 | 100% | 6 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Perinatal | - | - | - | - | 1 | 100% | 1 | 100% | - | - | - | - |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Subtotal | - | - | - | - | 1 | 100% | 1 | 100% | - | - | - | - |
| Total | 2 | | 7 | | 470 | | 33 | | 361 | | 24 | |

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and exposure category, Ohio

2020 diagnosis of HIV infection

| Exposure Category ^d | American Indian/Alaska Native | | Asian/Pacific Islander | | Black/African-American | | Hispanic/ Latinx ^a | | White | | Multi-Race | |
|----------------------------------------------------------|-------------------------------|-----|------------------------|-----|------------------------|-----|-------------------------------|-----|------------|-----|------------|-----|
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sexual contact only | 1 | 50% | 3 | 43% | 210 | 45% | 17 | 52% | 148 | 41% | 12 | 50% |
| Injection drug use (IDU) only | - | - | - | - | 5 | 1% | - | - | 23 | 6% | 1 | 4% |
| Heterosexual contact only | - | - | - | - | 95 | 20% | 2 | 6% | 36 | 10% | 3 | 13% |
| Male-to-male sexual contact & IDU | - | - | - | - | 2 | <1% | 1 | 3% | 22 | 6% | 1 | 4% |
| IDU & Heterosexual contact | - | - | - | - | 1 | <1% | - | - | 23 | 6% | 2 | 8% |
| Male-to-male sexual contact & Heterosexual contact | - | - | - | - | 6 | 1% | - | - | 2 | 1% | 1 | 4% |
| Male-to-male sexual contact & IDU & Heterosexual contact | - | - | - | - | 2 | <1% | - | - | 3 | 1% | - | - |
| Perinatal exposure | - | - | - | - | 1 | <1% | 1 | 3% | - | - | - | - |
| Other/unknown | 1 | 50% | 4 | 57% | 148 | 31% | 12 | 36% | 104 | 29% | 4 | 17% |
| Total | 2 | | 7 | | 470 | | 33 | | 361 | | 24 | |

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by disease status and county, Ohio

| County ^a | 2020 Diagnosis of HIV Infection | | Disease Status | | | County ^a | 2020 Diagnosis of HIV Infection | | Disease Status | | |
|---------------------|---------------------------------|-----|----------------|------------------|------|---------------------|---------------------------------|------------|----------------|------------------|-----------|
| | Rate ^b | No. | HIV (not AIDS) | HIV & later AIDS | AIDS | | Rate ^b | No. | HIV (not AIDS) | HIV & later AIDS | AIDS |
| | | | No. | No. | No. | | | | No. | No. | |
| Adams | * | 1 | 1 | - | - | Logan | * | 1 | 1 | - | - |
| Allen | 5.9 | 6 | 4 | 1 | 1 | Lorain | 6.7 | 21 | 14 | 6 | 1 |
| Ashland | * | 1 | - | 1 | - | Lucas | 5.4 | 23 | 15 | 7 | 1 |
| Ashtabula | * | 3 | 2 | - | 1 | Madison | * | 1 | 1 | - | - |
| Athens | * | 3 | 2 | 1 | - | Mahoning | 9.3 | 21 | 15 | 5 | 1 |
| Auglaize | * | 1 | 1 | - | - | Marion | * | 1 | 1 | - | - |
| Belmont | * | 3 | 3 | - | - | Medina | * | 2 | 2 | - | - |
| Brown | * | - | - | - | - | Meigs | * | - | - | - | - |
| Butler | 5.2 | 20 | 17 | 3 | - | Mercer | * | - | - | - | - |
| Carroll | * | - | - | - | - | Miami | * | 2 | 1 | 1 | - |
| Champaign | * | 1 | 1 | - | - | Monroe | * | - | - | - | - |
| Clark | 5.2 | 7 | 2 | 5 | - | Montgomery | 6.6 | 35 | 25 | 5 | 5 |
| Clermont | 2.9 | 6 | 2 | 4 | - | Morgan | * | 1 | - | 1 | - |
| Clinton | * | 1 | 1 | - | - | Morrow | * | 1 | - | 1 | - |
| Columbiana | * | - | - | - | - | Muskingum | * | 3 | 3 | - | - |
| Coshocton | * | - | - | - | - | Noble | * | - | - | - | - |
| Crawford | * | - | - | - | - | Ottawa | * | - | - | - | - |
| Cuyahoga | 15.5 | 190 | 157 | 28 | 5 | Paulding | * | - | - | - | - |
| Darke | * | 2 | - | 1 | 1 | Perry | * | - | - | - | - |
| Defiance | * | 2 | 1 | - | 1 | Pickaway | * | 1 | 1 | - | - |
| Delaware | * | 3 | 2 | 1 | - | Pike | * | - | - | - | - |
| Erie | 10.9 | 8 | 7 | 1 | - | Portage | * | 4 | 2 | 2 | - |
| Fairfield | 6.3 | 10 | 7 | 2 | 1 | Preble | * | 1 | 1 | - | - |
| Fayette | * | - | - | - | - | Putnam | * | - | - | - | - |
| Franklin | 15.4 | 204 | 158 | 37 | 9 | Richland | * | 3 | 1 | 2 | - |
| Fulton | * | - | - | - | - | Ross | * | 3 | 2 | - | 1 |
| Gallia | * | 1 | 1 | - | - | Sandusky | 8.6 | 5 | 4 | 1 | - |
| Geauga | * | 1 | 1 | - | - | Scioto | * | - | - | - | - |
| Greene | 5.9 | 10 | 7 | 2 | 1 | Seneca | * | - | - | - | - |
| Guernsey | * | 1 | 1 | - | - | Shelby | * | 3 | 3 | - | - |
| Hamilton | 16.3 | 133 | 102 | 26 | 5 | Stark | 7.8 | 29 | 25 | 4 | - |
| Hancock | * | - | - | - | - | Summit | 8.7 | 47 | 39 | 7 | 1 |
| Hardin | * | 1 | 1 | - | - | Trumbull | 2.5 | 5 | 4 | - | 1 |
| Harrison | * | - | - | - | - | Tuscarawas | * | 2 | - | 2 | - |
| Henry | * | - | - | - | - | Union | * | 1 | 1 | - | - |
| Highland | * | 1 | 1 | - | - | Van Wert | * | - | - | - | - |
| Hocking | * | - | - | - | - | Vinton | * | - | - | - | - |
| Holmes | * | - | - | - | - | Warren | 3.8 | 9 | 6 | 1 | 2 |
| Huron | * | 2 | 1 | 1 | - | Washington | * | 1 | 1 | - | - |
| Jackson | * | 1 | 1 | - | - | Wayne | * | 3 | 3 | - | - |
| Jefferson | * | 3 | 1 | 1 | 1 | Williams | * | - | - | - | - |
| Knox | * | 2 | - | 1 | 1 | Wood | * | 1 | - | 1 | - |
| Lake | 2.6 | 6 | 3 | 3 | - | Wyandot | * | - | - | - | - |
| Lawrence | 10.2 | 6 | 6 | - | - | No County | * | 18 | 18 | - | - |
| Licking | 5.1 | 9 | 9 | - | - | Ohio | 7.7 | 897 | 692 | 165 | 40 |

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
 Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a County reflects county of residence at time of initial diagnosis. Cases diagnosed while in a state or federal correctional facility or whose county is unknown are included in No County.
^bThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by selected characteristics, Ohio

| Characteristic | Diagnoses of HIV Infection by Year | | | | | | | | | | | | | | | Cumulative Diagnoses | |
|----------------------------------------------------|------------------------------------|------------|-----|-------------------|------------|-----|-------------------|------------|-----|-------------------|------------|-----|-------------------|------------|-----|----------------------|-----|
| | 2016 | | | 2017 | | | 2018 | | | 2019 | | | 2020 | | | No. | % |
| | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % | | |
| Sex at birth | | | | | | | | | | | | | | | | | |
| Males | 13.6 | 772 | 80% | 14.0 | 801 | 82% | 14.1 | 807 | 82% | 13.5 | 775 | 80% | 12.9 | 737 | 82% | 31,426 | 82% |
| Females | 3.2 | 192 | 20% | 3.0 | 181 | 18% | 2.9 | 174 | 18% | 3.3 | 199 | 20% | 2.7 | 160 | 18% | 6,711 | 18% |
| Age at diagnosis (yr) | | | | | | | | | | | | | | | | | |
| <13 | * | 4 | <1% | * | 2 | <1% | * | 4 | <1% | * | 1 | <1% | * | 2 | <1% | 350 | 1% |
| 13-14 | * | 1 | <1% | * | 2 | <1% | * | - | - | * | - | - | * | - | - | 45 | <1% |
| 15-19 | 6.8 | 52 | 5% | 7.7 | 59 | 6% | 7.1 | 54 | 6% | 5.6 | 42 | 4% | 5.5 | 41 | 5% | 1,416 | 4% |
| 20-24 | 26.3 | 203 | 21% | 23.4 | 179 | 18% | 23.1 | 175 | 18% | 25.2 | 189 | 19% | 21.8 | 163 | 18% | 5,099 | 13% |
| 25-29 | 25.3 | 198 | 21% | 29.3 | 233 | 24% | 24.5 | 198 | 20% | 23.3 | 187 | 19% | 21.4 | 170 | 19% | 6,688 | 18% |
| 30-34 | 18.3 | 133 | 14% | 17.7 | 129 | 13% | 20.3 | 150 | 15% | 19.8 | 149 | 15% | 19.0 | 146 | 16% | 6,951 | 18% |
| 35-39 | 12.9 | 91 | 9% | 12.6 | 90 | 9% | 15.6 | 113 | 12% | 13.9 | 101 | 10% | 13.0 | 94 | 10% | 6,069 | 16% |
| 40-44 | 11.1 | 75 | 8% | 10.8 | 72 | 7% | 11.5 | 77 | 8% | 12.6 | 85 | 9% | 9.2 | 63 | 7% | 4,495 | 12% |
| 45-49 | 8.4 | 63 | 7% | 9.0 | 67 | 7% | 10.6 | 78 | 8% | 8.7 | 62 | 6% | 9.4 | 65 | 7% | 3,087 | 8% |
| 50-54 | 7.9 | 63 | 7% | 8.8 | 68 | 7% | 7.3 | 55 | 6% | 8.3 | 61 | 6% | 8.5 | 62 | 7% | 1,932 | 5% |
| 55-64 | 4.2 | 67 | 7% | 4.0 | 64 | 7% | 3.9 | 63 | 6% | 4.5 | 72 | 7% | 4.6 | 73 | 8% | 1,595 | 4% |
| 65+ | 0.7 | 14 | 1% | 0.9 | 17 | 2% | 0.7 | 14 | 1% | 1.2 | 25 | 3% | 0.9 | 18 | 2% | 410 | 1% |
| Race/Ethnicity^b | | | | | | | | | | | | | | | | | |
| American Indian/Alaska Native | * | - | - | * | 1 | <1% | * | - | - | * | - | - | * | 2 | <1% | 26 | <1% |
| Asian/Pacific Islander | 3.9 | 10 | 1% | 4.8 | 13 | 1% | 2.8 | 8 | 1% | 2.4 | 7 | 1% | 2.3 | 7 | 1% | 163 | <1% |
| Black/African-American | 33.0 | 477 | 49% | 29.5 | 431 | 44% | 31.1 | 458 | 47% | 31.1 | 460 | 47% | 31.6 | 470 | 52% | 15,990 | 42% |
| Hispanic/Latinx | 12.5 | 53 | 5% | 15.2 | 67 | 7% | 12.8 | 59 | 6% | 13.0 | 61 | 6% | 6.8 | 33 | 4% | 1,943 | 5% |
| White | 4.1 | 376 | 39% | 4.6 | 421 | 43% | 4.6 | 422 | 43% | 4.4 | 405 | 42% | 4.0 | 361 | 40% | 18,814 | 49% |
| Multi-Race | 20.5 | 48 | 5% | 20.4 | 49 | 5% | 13.7 | 34 | 3% | 16.2 | 41 | 4% | 9.1 | 24 | 3% | 1,075 | 3% |
| Unknown | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 126 | <1% |
| Race/Ethnicity^b and Sex at birth | | | | | | | | | | | | | | | | | |
| American Indian/Alaska Native Males | * | - | - | * | 1 | <1% | * | - | - | * | - | - | * | 2 | <1% | 19 | <1% |
| American Indian/Alaska Native Females | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 7 | <1% |
| Asian/Pacific Islander Males | 7.3 | 9 | 1% | 5.4 | 7 | 1% | 5.0 | 7 | 1% | 4.9 | 7 | 1% | 4.1 | 6 | 1% | 129 | <1% |
| Asian/Pacific Islander Females | * | 1 | <1% | 4.2 | 6 | 1% | * | 1 | <1% | * | - | - | * | 1 | <1% | 34 | <1% |
| Black/African-American Males | 54.4 | 377 | 39% | 49.3 | 345 | 35% | 53.5 | 378 | 39% | 51.6 | 366 | 38% | 52.9 | 377 | 42% | 12,223 | 32% |
| Black/African-American Females | 13.3 | 100 | 10% | 11.3 | 86 | 9% | 10.4 | 80 | 8% | 12.2 | 94 | 10% | 12.0 | 93 | 10% | 3,767 | 10% |
| Hispanic/Latino Males | 21.1 | 46 | 5% | 27.0 | 61 | 6% | 23.2 | 55 | 6% | 21.6 | 52 | 5% | 12.4 | 31 | 3% | 1,540 | 4% |
| Hispanic/Latina Females | 3.4 | 7 | 1% | 2.8 | 6 | 1% | * | 4 | <1% | 3.9 | 9 | 1% | * | 2 | <1% | 403 | 1% |
| White Males | 6.8 | 306 | 32% | 7.6 | 345 | 35% | 7.5 | 339 | 35% | 7.1 | 320 | 33% | 6.8 | 303 | 34% | 16,558 | 43% |
| White Females | 1.5 | 70 | 7% | 1.6 | 76 | 8% | 1.8 | 83 | 8% | 1.8 | 85 | 9% | 1.2 | 58 | 6% | 2,256 | 6% |
| Multi-Race Males | 29.4 | 34 | 4% | 35.3 | 42 | 4% | 22.8 | 28 | 3% | 24.0 | 30 | 3% | 13.7 | 18 | 2% | 864 | 2% |
| Multi-Race Females | 11.8 | 14 | 1% | 5.7 | 7 | 1% | 4.8 | 6 | 1% | 8.6 | 11 | 1% | 4.5 | 6 | 1% | 211 | 1% |
| Unknown | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 126 | <1% |
| Total | 8.3 | 964 | | 8.4 | 982 | | 8.4 | 981 | | 8.3 | 974 | | 7.7 | 897 | | 38,137 | |

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by transmission category, Ohio

| Transmission Category ^a | Diagnoses of HIV Infection by Year | | | | | | | | | | Cumulative Diagnoses | |
|----------------------------------------|------------------------------------|------|------------|------|------------|------|------------|------|------------|------|----------------------|------|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Male adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 514 | 67% | 528 | 66% | 512 | 64% | 450 | 58% | 400 | 54% | 20,735 | 66% |
| Injection drug use (IDU) | 30 | 4% | 50 | 6% | 67 | 8% | 53 | 7% | 34 | 5% | 1,900 | 6% |
| Male-to-male sexual contact and IDU | 37 | 5% | 36 | 4% | 37 | 5% | 53 | 7% | 31 | 4% | 1,794 | 6% |
| Heterosexual contact | 33 | 4% | 20 | 2% | 26 | 3% | 34 | 4% | 26 | 4% | 1,641 | 5% |
| Other/unknown | 157 | 20% | 167 | 21% | 163 | 20% | 185 | 24% | 244 | 33% | 5,179 | 17% |
| Subtotal | 771 | 100% | 801 | 100% | 805 | 100% | 775 | 100% | 735 | 100% | 31,249 | 100% |
| Female adult or adolescent | | | | | | | | | | | | |
| Injection drug use | 32 | 17% | 43 | 24% | 48 | 28% | 50 | 25% | 21 | 13% | 1,057 | 16% |
| Heterosexual contact | 147 | 78% | 129 | 72% | 114 | 66% | 135 | 68% | 110 | 69% | 4,845 | 74% |
| Other/unknown | 10 | 5% | 7 | 4% | 10 | 6% | 13 | 7% | 29 | 18% | 638 | 10% |
| Subtotal | 189 | 100% | 179 | 100% | 172 | 100% | 198 | 100% | 160 | 100% | 6,540 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Perinatal | 4 | 100% | 1 | 50% | 4 | 100% | 1 | 100% | 2 | 100% | 282 | 81% |
| Other/unknown | - | - | 1 | 50% | - | - | - | - | - | - | 66 | 19% |
| Subtotal | 4 | 100% | 2 | 100% | 4 | 100% | 1 | 100% | 2 | 100% | 348 | 100% |
| Total | 964 | | 982 | | 981 | | 974 | | 897 | | 38,137 | |

Trends in reported diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by exposure category, Ohio

| Exposure Category ^b | Diagnoses of HIV Infection by Year | | | | | | | | | | Cumulative Diagnoses | |
|----------------------------------------------------------|------------------------------------|-----|------------|-----|------------|-----|------------|-----|------------|-----|----------------------|-----|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Male-to-male sexual contact only | 507 | 53% | 514 | 52% | 494 | 50% | 437 | 45% | 391 | 44% | 19,741 | 52% |
| Injection drug use (IDU) only | 30 | 3% | 41 | 4% | 49 | 5% | 48 | 5% | 29 | 3% | 1,665 | 4% |
| Heterosexual contact only | 180 | 19% | 149 | 15% | 140 | 14% | 169 | 17% | 136 | 15% | 6,494 | 17% |
| Male-to-male sexual contact & IDU | 29 | 3% | 26 | 3% | 31 | 3% | 44 | 5% | 26 | 3% | 1,450 | 4% |
| IDU & Heterosexual contact | 32 | 3% | 52 | 5% | 66 | 7% | 55 | 6% | 26 | 3% | 1,292 | 3% |
| Male-to-male sexual contact & Heterosexual contact | 7 | 1% | 14 | 1% | 18 | 2% | 13 | 1% | 9 | 1% | 994 | 3% |
| Male-to-male sexual contact & IDU & Heterosexual contact | 8 | 1% | 10 | 1% | 6 | 1% | 9 | 1% | 5 | 1% | 344 | 1% |
| Perinatal exposure | 6 | 1% | 1 | <1% | 4 | <1% | 1 | <1% | 2 | <1% | 298 | 1% |
| Other/unknown | 165 | 17% | 175 | 18% | 173 | 18% | 198 | 20% | 273 | 30% | 5,859 | 15% |
| Total | 964 | | 982 | | 981 | | 974 | | 897 | | 38,137 | |

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by county, Ohio

| County ^a | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | Cumulative Diagnoses | County ^a | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | Cumulative Diagnoses |
|---------------------|-------------------|-----|-------------------|-----|-------------------|-----|-------------------|-----|-------------------|-----|----------------------|---------------------|-------------------|------------|-------------------|------------|-------------------|------------|-------------------|------------|-------------------|------------|----------------------|
| | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | No. | | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | No. |
| Adams | * | 1 | * | - | * | 1 | * | 1 | * | 1 | 30 | Logan | * | - | * | 1 | * | 3 | * | 2 | * | 1 | 47 |
| Allen | 8.7 | 9 | * | 4 | 6.8 | 7 | 6.8 | 7 | 5.9 | 6 | 280 | Lorain | 2.0 | 6 | 6.2 | 19 | 4.5 | 14 | 3.9 | 12 | 6.7 | 21 | 549 |
| Ashland | * | - | * | 1 | * | 1 | * | - | * | 1 | 34 | Lucas | 9.9 | 43 | 9.1 | 39 | 10.0 | 43 | 9.3 | 40 | 5.4 | 23 | 1,717 |
| Ashtabula | 5.1 | 5 | * | 3 | * | 4 | * | 4 | * | 3 | 124 | Madison | * | - | * | 4 | * | 3 | * | 1 | * | 1 | 51 |
| Athens | * | 3 | * | 1 | * | 1 | * | 2 | * | 3 | 81 | Mahoning | 8.3 | 19 | 6.1 | 14 | 7.4 | 17 | 2.6 | 6 | 9.3 | 21 | 768 |
| Auglaize | * | 2 | * | - | * | - | * | - | * | 1 | 31 | Marion | * | 2 | * | 3 | * | 3 | * | 4 | * | 1 | 113 |
| Belmont | * | 4 | 7.3 | 5 | * | 3 | * | 1 | * | 3 | 84 | Medina | 3.4 | 6 | 5.0 | 9 | 2.8 | 5 | * | 2 | * | 2 | 112 |
| Brown | * | - | * | 1 | * | 1 | * | 1 | * | - | 36 | Meigs | * | - | * | - | * | - | * | - | * | - | 16 |
| Butler | 7.9 | 30 | 6.8 | 26 | 8.4 | 32 | 6.3 | 24 | 5.2 | 20 | 614 | Mercer | * | - | * | - | * | 1 | * | - | * | - | 25 |
| Carroll | * | 3 | * | - | * | - | * | - | * | - | 22 | Miami | * | 3 | * | 3 | * | 1 | 4.7 | 5 | * | 2 | 137 |
| Champaign | * | - | * | 1 | * | 1 | * | 2 | * | 1 | 57 | Monroe | * | - | * | 1 | * | - | * | - | * | - | 6 |
| Clark | 5.2 | 7 | * | 2 | 7.4 | 10 | 4.5 | 6 | 5.2 | 7 | 301 | Montgomery | 10.0 | 53 | 11.9 | 63 | 12.0 | 64 | 17.7 | 94 | 6.6 | 35 | 2,454 |
| Clermont | 4.4 | 9 | * | 1 | 5.8 | 12 | 5.8 | 12 | 2.9 | 6 | 194 | Morgan | * | - | * | 1 | * | - | * | 1 | * | 1 | 14 |
| Clinton | * | 1 | * | 1 | 11.9 | 5 | * | 3 | * | 1 | 62 | Morrow | * | 1 | * | 2 | * | - | * | 2 | * | 1 | 37 |
| Columbiana | 5.8 | 6 | * | 3 | * | 3 | * | 1 | * | - | 123 | Muskingum | * | 2 | * | 1 | * | 2 | * | 3 | * | 3 | 125 |
| Coshocton | * | - | * | - | * | 2 | * | - | * | - | 25 | Noble | * | - | * | - | * | - | * | 1 | * | - | 2 |
| Crawford | * | - | * | 2 | * | 2 | * | - | * | - | 43 | Ottawa | * | - | * | - | * | - | * | - | * | - | 37 |
| Cuyahoga | 15.3 | 191 | 12.0 | 150 | 12.0 | 149 | 13.0 | 160 | 15.5 | 190 | 8,143 | Paulding | * | - | * | - | * | - | * | - | * | - | 19 |
| Darke | * | - | * | 1 | * | - | * | 4 | * | 2 | 67 | Perry | * | - | * | 1 | * | - | * | - | * | - | 31 |
| Defiance | * | - | * | 2 | * | - | * | - | * | 2 | 48 | Pickaway | * | 2 | * | 1 | * | 1 | * | - | * | 1 | 89 |
| Delaware | 3.1 | 6 | * | 2 | * | 4 | * | 4 | * | 3 | 180 | Pike | * | - | * | 1 | * | 2 | * | - | * | - | 32 |
| Erie | * | 3 | * | 4 | 6.7 | 5 | * | 4 | 10.9 | 8 | 156 | Portage | 4.3 | 7 | 4.3 | 7 | 3.1 | 5 | 3.7 | 6 | * | 4 | 186 |
| Fairfield | 4.6 | 7 | * | 4 | * | 3 | * | 3 | 6.3 | 10 | 196 | Preble | * | - | * | - | * | 1 | * | - | * | 1 | 31 |
| Fayette | * | - | * | 2 | * | 2 | * | - | * | - | 34 | Putnam | * | 1 | * | - | * | 1 | * | 1 | * | - | 18 |
| Franklin | 15.6 | 197 | 17.3 | 223 | 15.2 | 199 | 16.3 | 215 | 15.4 | 204 | 7,873 | Richland | 5.0 | 6 | * | 3 | * | 2 | * | 2 | * | 3 | 208 |
| Fulton | * | 1 | * | - | * | 1 | * | - | * | - | 37 | Ross | * | 3 | * | 1 | * | - | * | 3 | * | 3 | 128 |
| Gallia | * | 2 | * | - | * | - | * | 3 | * | 1 | 31 | Sandusky | * | 3 | * | 4 | * | - | * | - | 8.6 | 5 | 81 |
| Geauga | * | 1 | * | 1 | * | - | * | - | * | 1 | 61 | Scioto | * | 2 | * | 2 | * | 2 | * | 3 | * | - | 93 |
| Greene | 3.6 | 6 | 7.8 | 13 | 7.1 | 12 | 7.1 | 12 | 5.9 | 10 | 283 | Seneca | * | 1 | * | 2 | * | 3 | * | 1 | * | - | 49 |
| Guernsey | * | - | * | - | * | - | * | - | * | 1 | 31 | Shelby | * | 4 | * | 4 | * | 1 | 10.3 | 5 | * | 3 | 67 |
| Hamilton | 17.2 | 139 | 22.9 | 186 | 22.7 | 185 | 20.8 | 170 | 16.3 | 133 | 5,136 | Stark | 8.6 | 32 | 7.2 | 27 | 8.3 | 31 | 4.0 | 15 | 7.8 | 29 | 807 |
| Hancock | * | - | 6.6 | 5 | * | 4 | * | 3 | * | - | 83 | Summit | 10.4 | 56 | 8.3 | 45 | 9.0 | 49 | 9.1 | 49 | 8.7 | 47 | 1,589 |
| Hardin | * | - | * | 1 | * | 1 | * | - | * | 1 | 21 | Trumbull | 4.0 | 8 | 4.0 | 8 | 5.5 | 11 | 5.1 | 10 | 2.5 | 5 | 307 |
| Harrison | * | - | * | 1 | * | 1 | * | - | * | - | 15 | Tuscarawas | * | 1 | * | 3 | * | 2 | * | 1 | * | 2 | 51 |
| Henry | * | - | * | 1 | * | - | * | 1 | * | - | 19 | Union | * | 3 | * | 2 | * | 1 | * | 2 | * | 1 | 49 |
| Highland | * | 1 | * | 3 | * | 2 | * | 3 | * | 1 | 43 | Van Wert | * | 1 | * | - | * | - | * | - | * | - | 13 |
| Hocking | * | - | * | 1 | * | 1 | * | - | * | - | 32 | Vinton | * | - | * | - | * | - | * | - | * | - | 6 |
| Holmes | * | - | * | - | * | - | * | - | * | - | 14 | Warren | * | 4 | 2.2 | 5 | 2.6 | 6 | 2.6 | 6 | 3.8 | 9 | 186 |
| Huron | * | 1 | * | - | * | - | * | - | * | 2 | 44 | Washington | * | - | * | 3 | * | 1 | * | 4 | * | 1 | 97 |
| Jackson | * | 3 | * | 1 | * | 2 | * | - | * | 1 | 33 | Wayne | * | 2 | * | 2 | * | 1 | 4.3 | 5 | * | 3 | 93 |
| Jefferson | 7.5 | 5 | * | 1 | * | - | * | 1 | * | 3 | 114 | Williams | * | - | * | 1 | * | 1 | * | - | * | - | 46 |
| Knox | * | 1 | * | - | * | 2 | * | 1 | * | 2 | 45 | Wood | * | 3 | * | 2 | 6.9 | 9 | 4.6 | 6 | * | 1 | 132 |
| Lake | 7.4 | 17 | 6.1 | 14 | * | 3 | 2.6 | 6 | 2.6 | 6 | 288 | Wyandot | * | - | * | - | * | - | * | 1 | * | - | 9 |
| Lawrence | * | 2 | * | 1 | 10.0 | 6 | * | 3 | 10.2 | 6 | 89 | No County | * | 23 | * | 24 | * | 16 | * | 18 | * | 18 | 2,007 |
| Licking | * | 4 | 3.5 | 6 | 6.8 | 12 | 3.4 | 6 | 5.1 | 9 | 246 | Ohio | 8.3 | 964 | 8.4 | 982 | 8.4 | 981 | 8.3 | 974 | 7.7 | 897 | 38,137 |

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a County reflects county of residence at time of initial diagnosis. Cases diagnosed while in a state or federal correctional facility or whose county is unknown are included in No County.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection among transgender people by year of diagnosis (2016-2020) and cumulative diagnoses by selected characteristics, Ohio

| Characteristic | Diagnoses of HIV Infection by Year | | | | | | | | | | Cumulative Diagnoses | |
|------------------------------------------------------|------------------------------------|------|-----------|------|-----------|------|-----------|-----|-----------|------|----------------------|-----|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Current Gender | | | | | | | | | | | | |
| Transgender Women | 16 | 100% | 15 | 100% | 19 | 100% | 19 | 86% | 24 | 100% | 265 | 97% |
| Transgender Men | - | - | - | - | - | - | 3 | 14% | - | - | 7 | 3% |
| Age at diagnosis (yr) | | | | | | | | | | | | |
| <13 | - | - | - | - | - | - | - | - | - | - | 1 | <1% |
| 13-14 | - | - | - | - | - | - | - | - | - | - | - | - |
| 15-19 | 3 | 19% | 1 | 7% | 2 | 11% | 2 | 9% | 1 | 4% | 41 | 15% |
| 20-24 | 3 | 19% | 3 | 20% | 8 | 42% | 3 | 14% | 7 | 29% | 70 | 26% |
| 25-29 | 6 | 38% | 4 | 27% | 5 | 26% | 10 | 45% | 5 | 21% | 60 | 22% |
| 30-34 | 2 | 13% | 5 | 33% | 1 | 5% | 5 | 23% | 5 | 21% | 46 | 17% |
| 35-39 | 2 | 13% | 1 | 7% | 2 | 11% | - | - | 3 | 13% | 24 | 9% |
| 40-44 | - | - | - | - | - | - | - | - | - | - | 6 | 2% |
| 45-49 | - | - | - | - | 1 | 5% | 1 | 5% | 1 | 4% | 13 | 5% |
| 50-54 | - | - | 1 | 7% | - | - | - | - | 1 | 4% | 6 | 2% |
| 55-64 | - | - | - | - | - | - | 1 | 5% | 1 | 4% | 5 | 2% |
| 65+ | - | - | - | - | - | - | - | - | - | - | - | - |
| Race/Ethnicity^a | | | | | | | | | | | | |
| American Indian/Alaska Native | - | - | - | - | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander | - | - | - | - | - | - | - | - | - | - | - | - |
| Black/African-American | 15 | 94% | 10 | 67% | 11 | 58% | 17 | 77% | 16 | 67% | 199 | 73% |
| Hispanic/Latinx | - | - | - | - | 2 | 11% | 1 | 5% | 1 | 4% | 11 | 4% |
| White | 1 | 6% | 4 | 27% | 4 | 21% | 3 | 14% | 6 | 25% | 48 | 18% |
| Multi-Race | - | - | 1 | 7% | 2 | 11% | 1 | 5% | 1 | 4% | 14 | 5% |
| Unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Race/Ethnicity^a and Current Gender | | | | | | | | | | | | |
| American Indian/Alaska Native Transgender Women | - | - | - | - | - | - | - | - | - | - | - | - |
| American Indian/Alaska Native Transgender Men | - | - | - | - | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander Transgender Women | - | - | - | - | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander Transgender Men | - | - | - | - | - | - | - | - | - | - | - | - |
| Black/African-American Transgender Women | 15 | 94% | 10 | 67% | 11 | 58% | 15 | 68% | 16 | 67% | 193 | 71% |
| Black/African-American Transgender Men | - | - | - | - | - | - | 2 | 9% | - | - | 6 | 2% |
| Hispanic/Latina Transgender Women | - | - | - | - | 2 | 11% | 1 | 5% | 1 | 4% | 11 | 4% |
| Hispanic/Latino Transgender Men | - | - | - | - | - | - | - | - | - | - | - | - |
| White Transgender Women | 1 | 6% | 4 | 27% | 4 | 21% | 3 | 14% | 6 | 25% | 48 | 18% |
| White Transgender Men | - | - | - | - | - | - | - | - | - | - | - | - |
| Multi-Race Transgender Women | - | - | 1 | 7% | 2 | 11% | - | - | 1 | 4% | 13 | 5% |
| Multi-Race Transgender Men | - | - | - | - | - | - | 1 | 5% | - | - | 1 | <1% |
| Total | 16 | | 15 | | 19 | | 22 | | 24 | | 272 | |

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Transgender is a term used to describe persons whose current gender identity is different than their sex (male or female) assigned at birth. Transgender women identify as female, but were assigned male at birth and transgender men identify as male, but were assigned female at birth.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection among transgender people by year of diagnosis (2016-2020) and cumulative diagnoses by transmission category, Ohio

Diagnoses of HIV Infection by Year

| Transmission Category ^a | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | Cumulative Diagnoses | |
|------------------------------------|-----------|------|-----------|-----|-----------|-----|-----------|-----|-----------|-----|----------------------|-----|
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Transgender Women | | | | | | | | | | | | |
| Sexual contact | 16 | 100% | 12 | 80% | 16 | 84% | 14 | 74% | 23 | 96% | 236 | 89% |
| Injection drug use (IDU) | - | - | - | - | - | - | - | - | - | - | - | - |
| Sexual contact and IDU | - | - | 1 | 7% | 3 | 16% | 3 | 16% | - | - | 18 | 7% |
| Other/unknown | - | - | 2 | 13% | - | - | 2 | 11% | 1 | 4% | 11 | 4% |
| Subtotal | 16 | | 15 | | 19 | | 19 | | 24 | | 265 | |
| Transgender Men | | | | | | | | | | | | |
| Injection drug use | - | - | - | - | - | - | - | - | - | - | - | - |
| Sexual contact | - | - | - | - | - | - | 1 | 33% | - | - | 5 | 71% |
| Other/unknown | - | - | - | - | - | - | 2 | 67% | - | - | 2 | 29% |
| Subtotal | - | - | - | - | - | - | 3 | | - | | 7 | |
| Total | 16 | | 15 | | 19 | | 22 | | 24 | | 272 | |

Trends in reported diagnoses of HIV infection among transgender people by year of diagnosis (2016-2020) and cumulative diagnoses by region, Ohio

Diagnoses of HIV Infection by Year

| Region ^b | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | Cumulative Diagnoses | |
|---------------------------|-----------|-----|-----------|-----|-----------|-----|-----------|-----|-----------|-----|----------------------|-----|
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Region 1 | 1 | 6% | - | - | 2 | 11% | - | - | 1 | 4% | 11 | 4% |
| Region 2 | - | - | - | - | - | - | - | - | - | - | 2 | 1% |
| Part A-Cleveland/Region 3 | 7 | 44% | 4 | 27% | 3 | 16% | 5 | 23% | 7 | 29% | 68 | 25% |
| Region 4 | 1 | 6% | 3 | 20% | - | - | 3 | 14% | 1 | 4% | 13 | 5% |
| Region 5 | 1 | 6% | - | - | 1 | 5% | - | - | 1 | 4% | 8 | 3% |
| Region 6 | - | - | - | - | - | - | - | - | - | - | 2 | 1% |
| Region 7 | - | - | - | - | - | - | - | - | 1 | 4% | 2 | 1% |
| Region 8 | 2 | 13% | 2 | 13% | 3 | 16% | 4 | 18% | 5 | 21% | 58 | 21% |
| Region 9 | - | - | - | - | 2 | 11% | 2 | 9% | 6 | 25% | 29 | 11% |
| Region 10 | - | - | - | - | - | - | 1 | 5% | - | - | 1 | <1% |
| Part A-Columbus/Region 11 | 4 | 25% | 4 | 27% | 8 | 42% | 5 | 23% | 2 | 8% | 60 | 22% |
| No County | - | - | 2 | 13% | - | - | 2 | 9% | - | - | 18 | 7% |
| Total | 16 | | 15 | | 19 | | 22 | | 24 | | 272 | |

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Transgender is a term used to describe persons whose current gender identity is different than their sex (male or female) assigned at birth. Transgender women identify as female, but were assigned male at birth and transgender men identify as male, but were assigned female at birth.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. For the purposes of displaying transmission categories by current gender, the CDC-defined categories have been modified slightly.

^b Region reflects HIV Prevention Planning Region of residence at time of initial diagnosis. Cases diagnosed while in a state or federal correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and selected characteristics, Ohio

| Characteristic | AIDS Diagnoses | | | | | |
|----------------------------------------------------|---------------------|-----|--------------------------------|-----|---------------------------------------------|-----|
| | 2020 AIDS diagnosis | | ≥12 Months after HIV Diagnosis | | <12 Months after HIV Diagnosis ^a | |
| | No. | % | No. | % | No. | % |
| Sex at birth | | | | | | |
| Males | 297 | 82% | 123 | 81% | 174 | 82% |
| Females | 66 | 18% | 29 | 19% | 37 | 18% |
| Age at AIDS diagnosis (yr) | | | | | | |
| <13 | 1 | <1% | - | - | 1 | <1% |
| 13-14 | 1 | <1% | 1 | 1% | - | - |
| 15-19 | 2 | 1% | - | - | 2 | 1% |
| 20-24 | 28 | 8% | 10 | 7% | 18 | 9% |
| 25-29 | 52 | 14% | 21 | 14% | 31 | 15% |
| 30-34 | 51 | 14% | 17 | 11% | 34 | 16% |
| 35-39 | 46 | 13% | 21 | 14% | 25 | 12% |
| 40-44 | 30 | 8% | 18 | 12% | 12 | 6% |
| 45-49 | 31 | 9% | 13 | 9% | 18 | 9% |
| 50-54 | 43 | 12% | 13 | 9% | 30 | 14% |
| 55-64 | 56 | 15% | 25 | 16% | 31 | 15% |
| 65+ | 22 | 6% | 13 | 9% | 9 | 4% |
| Race/Ethnicity^b | | | | | | |
| American Indian/Alaska Native | - | - | - | - | - | - |
| Asian/Pacific Islander | 4 | 1% | 2 | 1% | 2 | 1% |
| Black/African-American | 152 | 42% | 68 | 45% | 84 | 40% |
| Hispanic/Latinx | 21 | 6% | 10 | 7% | 11 | 5% |
| White | 164 | 45% | 60 | 39% | 104 | 49% |
| Multi-Race | 22 | 6% | 12 | 8% | 10 | 5% |
| Race/Ethnicity^b and Sex at birth | | | | | | |
| American Indian/Alaska Native Males | - | - | - | - | - | - |
| American Indian/Alaska Native Females | - | - | - | - | - | - |
| Asian/Pacific Islander Males | 4 | 1% | 2 | 1% | 2 | 1% |
| Asian/Pacific Islander Females | - | - | - | - | - | - |
| Black/African-American Males | 115 | 32% | 51 | 34% | 64 | 30% |
| Black/African-American Females | 37 | 10% | 17 | 11% | 20 | 9% |
| Hispanic/Latino Males | 20 | 6% | 9 | 6% | 11 | 5% |
| Hispanic/Latina Females | 1 | <1% | 1 | 1% | - | - |
| White Males | 139 | 38% | 50 | 33% | 89 | 42% |
| White Females | 25 | 7% | 10 | 7% | 15 | 7% |
| Multi-Race Males | 19 | 5% | 11 | 7% | 8 | 4% |
| Multi-Race Females | 3 | 1% | 1 | 1% | 2 | 1% |
| Total | 363 | | 152 | | 211 | |

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and transmission category, Ohio

| Transmission Category ^b | AIDS Diagnoses | | | | | |
|----------------------------------------|---------------------|------|--------------------------------|------|---------------------------------------------|------|
| | 2020 AIDS diagnosis | | ≥12 Months after HIV Diagnosis | | <12 Months after HIV Diagnosis ^a | |
| | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | |
| Male-to-male sexual contact | 162 | 55% | 74 | 62% | 88 | 51% |
| Injection drug use (IDU) | 12 | 4% | 8 | 7% | 4 | 2% |
| Male-to-male sexual contact and IDU | 11 | 4% | 7 | 6% | 4 | 2% |
| Heterosexual contact | 17 | 6% | 11 | 9% | 6 | 3% |
| Other/unknown | 91 | 31% | 20 | 17% | 71 | 41% |
| Subtotal | 293 | 100% | 120 | 100% | 173 | 100% |
| Female adult or adolescent | | | | | | |
| Injection drug use | 10 | 15% | 7 | 24% | 3 | 8% |
| Heterosexual contact | 45 | 68% | 19 | 66% | 26 | 70% |
| Other/unknown | 11 | 17% | 3 | 10% | 8 | 22% |
| Subtotal | 66 | 100% | 29 | 100% | 37 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | |
| Perinatal | 4 | 100% | 3 | 100% | 1 | 100% |
| Other/unknown | - | - | - | - | - | - |
| Subtotal | 4 | 100% | 3 | 100% | 1 | 100% |
| Total | 363 | | 152 | | 211 | |

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and exposure category, Ohio

| Exposure Category ^c | AIDS Diagnoses | | | | | |
|----------------------------------------------------------|---------------------|-----|-------------------------------|-----|---------------------------------------------|-----|
| | 2020 AIDS diagnosis | | ≥12Months after HIV Diagnosis | | <12 Months after HIV Diagnosis ^a | |
| | No. | % | No. | % | No. | % |
| Male-to-male sexual contact only | 159 | 44% | 73 | 48% | 86 | 41% |
| Injection drug use (IDU) only | 11 | 3% | 6 | 4% | 5 | 2% |
| Heterosexual contact only | 62 | 17% | 30 | 20% | 32 | 15% |
| Male-to-male sexual contact & IDU | 8 | 2% | 5 | 3% | 3 | 1% |
| IDU & Heterosexual contact | 11 | 3% | 9 | 6% | 2 | 1% |
| Male-to-male sexual contact & Heterosexual contact | 3 | 1% | 1 | 1% | 2 | 1% |
| Male-to-male sexual contact & IDU & Heterosexual contact | 3 | 1% | 2 | 1% | 1 | <1% |
| Perinatal exposure | 4 | 1% | 3 | 2% | 1 | <1% |
| Other/unknown | 102 | 28% | 23 | 15% | 79 | 37% |
| Total | 363 | | 152 | | 211 | |

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and county, Ohio

| County ^b | AIDS Diagnoses | | | County ^b | AIDS Diagnoses | | |
|---------------------|---------------------|--------------------------------|---------------------------------------------|---------------------|---------------------|--------------------------------|---------------------------------------------|
| | 2020 AIDS Diagnosis | ≥12 Months after HIV Diagnosis | <12 Months after HIV Diagnosis ^a | | 2020 AIDS Diagnosis | ≥12 Months after HIV Diagnosis | <12 Months after HIV Diagnosis ^a |
| | No. | No. | No. | | No. | No. | No. |
| Adams | 1 | 1 | - | Logan | 1 | 1 | - |
| Allen | 2 | - | 2 | Lorain | 7 | 1 | 6 |
| Ashland | 1 | - | 1 | Lucas | 11 | 3 | 8 |
| Ashtabula | 2 | 1 | 1 | Madison | - | - | - |
| Athens | - | - | - | Mahoning | 7 | 1 | 6 |
| Auglaize | - | - | - | Marion | - | - | - |
| Belmont | 1 | - | 1 | Medina | 1 | 1 | - |
| Brown | - | - | - | Meigs | - | - | - |
| Butler | 6 | 3 | 3 | Mercer | - | - | - |
| Carroll | - | - | - | Miami | 2 | - | 2 |
| Champaign | - | - | - | Monroe | - | - | - |
| Clark | 6 | 1 | 5 | Montgomery | 19 | 6 | 13 |
| Clermont | 6 | 2 | 4 | Morgan | - | - | - |
| Clinton | 1 | 1 | - | Morrow | 1 | - | 1 |
| Columbiana | 1 | - | 1 | Muskingum | - | - | - |
| Coshocton | - | - | - | Noble | - | - | - |
| Crawford | - | - | - | Ottawa | - | - | - |
| Cuyahoga | 75 | 39 | 36 | Paulding | - | - | - |
| Darke | 1 | - | 1 | Perry | - | - | - |
| Defiance | 2 | 1 | 1 | Pickaway | 1 | 1 | - |
| Delaware | 2 | 1 | 1 | Pike | - | - | - |
| Erie | 3 | 2 | 1 | Portage | 3 | 1 | 2 |
| Fairfield | 4 | 1 | 3 | Preble | - | - | - |
| Fayette | - | - | - | Putnam | - | - | - |
| Franklin | 78 | 32 | 46 | Richland | 2 | - | 2 |
| Fulton | - | - | - | Ross | 1 | - | 1 |
| Gallia | - | - | - | Sandusky | 1 | - | 1 |
| Geauga | - | - | - | Scioto | - | - | - |
| Greene | 3 | 1 | 2 | Seneca | - | - | - |
| Guernsey | - | - | - | Shelby | - | - | - |
| Hamilton | 60 | 29 | 31 | Stark | 6 | - | 6 |
| Hancock | - | - | - | Summit | 19 | 11 | 8 |
| Hardin | - | - | - | Trumbull | 2 | 1 | 1 |
| Harrison | - | - | - | Tuscarawas | 2 | - | 2 |
| Henry | - | - | - | Union | - | - | - |
| Highland | - | - | - | Van Wert | - | - | - |
| Hocking | - | - | - | Vinton | - | - | - |
| Holmes | - | - | - | Warren | 3 | - | 3 |
| Huron | 1 | - | 1 | Washington | 1 | 1 | - |
| Jackson | 1 | 1 | - | Wayne | - | - | - |
| Jefferson | 1 | - | 1 | Williams | - | - | - |
| Knox | 2 | - | 2 | Wood | 1 | - | 1 |
| Lake | 5 | 2 | 3 | Wyandot | - | - | - |
| Lawrence | - | - | - | No County | 4 | 3 | 1 |
| Licking | 3 | 3 | - | Ohio | 363 | 152 | 211 |

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b County reflects county of residence at time of AIDS diagnosis. Cases diagnosed while in a state or federal correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by selected characteristics, Ohio

| Characteristic | Year of AIDS Diagnosis | | | | | | | | | | Cumulative AIDS | |
|----------------------------------------------------|------------------------|-----|------------|-----|------------|-----|------------|-----|------------|-----|-----------------|-----|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | AIDS | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Sex at birth | | | | | | | | | | | | |
| Males | 378 | 78% | 325 | 80% | 338 | 78% | 358 | 80% | 297 | 82% | 18,358 | 84% |
| Females | 104 | 22% | 81 | 20% | 94 | 22% | 87 | 20% | 66 | 18% | 3,502 | 16% |
| Age at AIDS diagnosis (yr) | | | | | | | | | | | | |
| <13 | 1 | <1% | - | - | 3 | 1% | - | - | 1 | <1% | 151 | 1% |
| 13-14 | - | - | - | - | - | - | - | - | 1 | <1% | 19 | <1% |
| 15-19 | 7 | 1% | 7 | 2% | 3 | 1% | 6 | 1% | 2 | 1% | 207 | 1% |
| 20-24 | 47 | 10% | 39 | 10% | 37 | 9% | 34 | 8% | 28 | 8% | 1,246 | 6% |
| 25-29 | 85 | 18% | 48 | 12% | 54 | 13% | 68 | 15% | 52 | 14% | 2,915 | 13% |
| 30-34 | 49 | 10% | 63 | 16% | 64 | 15% | 71 | 16% | 51 | 14% | 4,134 | 19% |
| 35-39 | 59 | 12% | 51 | 13% | 61 | 14% | 47 | 11% | 46 | 13% | 4,145 | 19% |
| 40-44 | 55 | 11% | 50 | 12% | 42 | 10% | 44 | 10% | 30 | 8% | 3,451 | 16% |
| 45-49 | 43 | 9% | 39 | 10% | 54 | 13% | 40 | 9% | 31 | 9% | 2,431 | 11% |
| 50-54 | 69 | 14% | 43 | 11% | 45 | 10% | 50 | 11% | 43 | 12% | 1,521 | 7% |
| 55-64 | 48 | 10% | 51 | 13% | 55 | 13% | 59 | 13% | 56 | 15% | 1,304 | 6% |
| 65+ | 19 | 4% | 15 | 4% | 14 | 3% | 26 | 6% | 22 | 6% | 336 | 2% |
| Race/Ethnicity^a | | | | | | | | | | | | |
| American Indian/Alaska Native | - | - | - | - | - | - | - | - | - | - | 11 | <1% |
| Asian/Pacific Islander | 3 | 1% | 2 | <1% | 3 | 1% | 2 | <1% | 4 | 1% | 64 | <1% |
| Black/African-American | 256 | 53% | 190 | 47% | 213 | 49% | 205 | 46% | 152 | 42% | 8,743 | 40% |
| Hispanic/Latinx | 27 | 6% | 33 | 8% | 24 | 6% | 31 | 7% | 21 | 6% | 1,021 | 5% |
| White | 182 | 38% | 160 | 39% | 180 | 42% | 192 | 43% | 164 | 45% | 11,492 | 53% |
| Multi-Race | 14 | 3% | 21 | 5% | 12 | 3% | 15 | 3% | 22 | 6% | 529 | 2% |
| Race/Ethnicity^a and Sex at birth | | | | | | | | | | | | |
| American Indian/Alaska Native Males | - | - | - | - | - | - | - | - | - | - | 6 | <1% |
| American Indian/Alaska Native Females | - | - | - | - | - | - | - | - | - | - | 5 | <1% |
| Asian/Pacific Islander Males | 3 | 1% | - | - | 2 | <1% | 2 | <1% | 4 | 1% | 52 | <1% |
| Asian/Pacific Islander Females | - | - | 2 | <1% | 1 | <1% | - | - | - | - | 12 | <1% |
| Black/African-American Males | 190 | 39% | 147 | 36% | 163 | 38% | 162 | 36% | 115 | 32% | 6,730 | 31% |
| Black/African-American Females | 66 | 14% | 43 | 11% | 50 | 12% | 43 | 10% | 37 | 10% | 2,013 | 9% |
| Hispanic/Latino Males | 22 | 5% | 29 | 7% | 22 | 5% | 24 | 5% | 20 | 6% | 799 | 4% |
| Hispanic/Latina Females | 5 | 1% | 4 | 1% | 2 | <1% | 7 | 2% | 1 | <1% | 222 | 1% |
| White Males | 153 | 32% | 131 | 32% | 142 | 33% | 159 | 36% | 139 | 38% | 10,346 | 47% |
| White Females | 29 | 6% | 29 | 7% | 38 | 9% | 33 | 7% | 25 | 7% | 1,146 | 5% |
| Multi-Race Males | 10 | 2% | 18 | 4% | 9 | 2% | 11 | 2% | 19 | 5% | 425 | 2% |
| Multi-Race Females | 4 | 1% | 3 | 1% | 3 | 1% | 4 | 1% | 3 | 1% | 104 | <1% |
| Total | 482 | | 406 | | 432 | | 445 | | 363 | | 21,860 | |

Notes:
 AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.
 Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by transmission category, Ohio

| Transmission Category ^a | Year of AIDS Diagnosis | | | | | | | | | | Cumulative AIDS | |
|----------------------------------------|------------------------|------|------------|------|------------|------|------------|------|------------|------|-----------------|------|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Male adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 234 | 62% | 212 | 65% | 199 | 59% | 200 | 56% | 162 | 55% | 12,383 | 68% |
| Injection drug use (IDU) | 16 | 4% | 11 | 3% | 18 | 5% | 27 | 8% | 12 | 4% | 1,239 | 7% |
| Male-to-male sexual contact and IDU | 13 | 3% | 13 | 4% | 13 | 4% | 17 | 5% | 11 | 4% | 1,155 | 6% |
| Heterosexual contact | 29 | 8% | 14 | 4% | 20 | 6% | 16 | 4% | 17 | 6% | 1,027 | 6% |
| Other/unknown | 84 | 22% | 75 | 23% | 86 | 26% | 96 | 27% | 91 | 31% | 2,444 | 13% |
| Subtotal | 376 | 100% | 325 | 100% | 336 | 100% | 356 | 100% | 293 | 100% | 18,248 | 100% |
| Female adult or adolescent | | | | | | | | | | | | |
| Injection drug use | 16 | 16% | 18 | 23% | 23 | 25% | 22 | 25% | 10 | 15% | 638 | 19% |
| Heterosexual contact | 85 | 83% | 55 | 69% | 61 | 66% | 61 | 70% | 45 | 68% | 2571 | 76% |
| Other/unknown | 2 | 2% | 7 | 9% | 8 | 9% | 4 | 5% | 11 | 17% | 193 | 6% |
| Subtotal | 103 | 100% | 80 | 100% | 92 | 100% | 87 | 100% | 66 | 100% | 3,402 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Perinatal | 3 | 100% | 1 | 100% | 4 | 100% | 2 | 100% | 4 | 100% | 167 | 80% |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | 43 | 20% |
| Subtotal | 3 | 100% | 1 | 100% | 4 | 100% | 2 | 100% | 4 | 100% | 210 | 100% |
| Total | 482 | | 406 | | 432 | | 445 | | 363 | | 21,860 | |

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by exposure category, Ohio

| Exposure Category ^b | Year of AIDS Diagnosis | | | | | | | | | | Cumulative AIDS | |
|----------------------------------------------------------|------------------------|-----|------------|-----|------------|-----|------------|-----|------------|-----|-----------------|-----|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Male-to-male sexual contact only | 225 | 47% | 202 | 50% | 193 | 45% | 197 | 44% | 159 | 44% | 11,758 | 54% |
| Injection drug use (IDU) only | 12 | 2% | 15 | 4% | 14 | 3% | 27 | 6% | 11 | 3% | 1,098 | 5% |
| Heterosexual contact only | 114 | 24% | 69 | 17% | 81 | 19% | 77 | 17% | 62 | 17% | 3,604 | 16% |
| Male-to-male sexual contact & IDU | 9 | 2% | 10 | 2% | 12 | 3% | 11 | 2% | 8 | 2% | 931 | 4% |
| IDU & Heterosexual contact | 20 | 4% | 14 | 3% | 27 | 6% | 22 | 5% | 11 | 3% | 779 | 4% |
| Male-to-male sexual contact & Heterosexual contact | 9 | 2% | 10 | 2% | 6 | 1% | 3 | 1% | 3 | 1% | 625 | 3% |
| Male-to-male sexual contact & IDU & Heterosexual contact | 4 | 1% | 3 | 1% | 1 | <1% | 6 | 1% | 3 | 1% | 224 | 1% |
| Perinatal exposure | 4 | 1% | 1 | <1% | 5 | 1% | 2 | <1% | 4 | 1% | 172 | 1% |
| Other/unknown | 85 | 18% | 82 | 20% | 93 | 22% | 100 | 22% | 102 | 28% | 2,669 | 12% |
| Total | 482 | | 406 | | 432 | | 445 | | 363 | | 21,860 | |

Notes:

AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by county, Ohio

| County ^a | 2016 | 2017 | 2018 | 2019 | 2020 | Cumulative AIDS | County ^a | 2016 | 2017 | 2018 | 2019 | 2020 | Cumulative AIDS |
|---------------------|------|------|------|------|------|-----------------|---------------------|------------|------------|------------|------------|------------|-----------------|
| | No. | No. | No. | No. | No. | No. | | No. | No. | No. | No. | No. | No. |
| Adams | 2 | - | - | 1 | 1 | 18 | Logan | - | 1 | 1 | 1 | 1 | 34 |
| Allen | 3 | 4 | 4 | 3 | 2 | 149 | Lorain | 4 | 9 | 8 | 8 | 7 | 357 |
| Ashland | - | - | 1 | - | 1 | 24 | Lucas | 38 | 18 | 27 | 24 | 11 | 1,021 |
| Ashtabula | 1 | 3 | 1 | 2 | 2 | 80 | Madison | - | 2 | 1 | 1 | - | 28 |
| Athens | 1 | 1 | 1 | - | - | 49 | Mahoning | 12 | 3 | 7 | 5 | 7 | 440 |
| Auglaize | 2 | - | - | 1 | - | 29 | Marion | 1 | - | 1 | 1 | - | 72 |
| Belmont | - | 2 | 2 | - | 1 | 50 | Medina | 1 | 2 | 1 | 3 | 1 | 56 |
| Brown | - | 1 | - | 1 | - | 27 | Meigs | - | - | - | 1 | - | 6 |
| Butler | 20 | 13 | 11 | 11 | 6 | 351 | Mercer | - | - | - | - | - | 21 |
| Carroll | - | 1 | - | - | - | 15 | Miami | 1 | - | 2 | 1 | 2 | 86 |
| Champaign | 1 | 1 | - | 2 | - | 45 | Monroe | - | - | - | - | - | 5 |
| Clark | 5 | 4 | 3 | 2 | 6 | 199 | Montgomery | 25 | 26 | 29 | 43 | 19 | 1,481 |
| Clermont | 2 | 2 | 5 | 2 | 6 | 116 | Morgan | - | 1 | - | - | - | 7 |
| Clinton | 1 | - | 1 | - | 1 | 35 | Morrow | 1 | - | - | 1 | 1 | 23 |
| Columbiana | 1 | - | - | - | 1 | 71 | Muskingum | 1 | 1 | 1 | 1 | - | 74 |
| Coshocton | - | - | 2 | - | - | 18 | Noble | - | - | - | - | - | 3 |
| Crawford | - | - | 1 | - | - | 32 | Ottawa | - | - | - | - | - | 17 |
| Cuyahoga | 88 | 69 | 91 | 88 | 75 | 4,907 | Paulding | 1 | - | - | - | - | 14 |
| Darke | - | 1 | - | 1 | 1 | 41 | Perry | - | 1 | - | - | - | 22 |
| Defiance | - | - | - | - | 2 | 29 | Pickaway | 2 | - | - | - | 1 | 41 |
| Delaware | 4 | 2 | 3 | 1 | 2 | 89 | Pike | - | 1 | 2 | - | - | 26 |
| Erie | 3 | 2 | 1 | 2 | 3 | 103 | Portage | 2 | 1 | 1 | 3 | 3 | 92 |
| Fairfield | 3 | - | 2 | 3 | 4 | 129 | Preble | 1 | - | - | - | - | 24 |
| Fayette | - | 2 | 1 | - | - | 25 | Putnam | - | - | - | 1 | - | 11 |
| Franklin | 99 | 98 | 85 | 94 | 78 | 4,294 | Richland | 3 | 1 | - | 2 | 2 | 144 |
| Fulton | - | - | - | - | - | 28 | Ross | 1 | 1 | - | 1 | 1 | 88 |
| Gallia | 2 | 1 | - | - | - | 21 | Sandusky | 1 | 1 | - | 1 | 1 | 58 |
| Geauga | - | 1 | - | - | - | 44 | Scioto | 1 | - | 3 | 1 | - | 61 |
| Greene | 4 | 1 | 3 | 5 | 3 | 160 | Seneca | - | 1 | - | 1 | - | 29 |
| Guernsey | 1 | - | - | - | - | 21 | Shelby | 1 | 1 | 2 | 1 | - | 38 |
| Hamilton | 69 | 71 | 69 | 67 | 60 | 2,924 | Stark | 15 | 7 | 12 | 9 | 6 | 423 |
| Hancock | 2 | 3 | 1 | 1 | - | 55 | Summit | 24 | 16 | 15 | 16 | 19 | 918 |
| Hardin | 2 | 1 | - | - | - | 16 | Trumbull | 3 | 3 | 2 | 8 | 2 | 176 |
| Harrison | - | 1 | 1 | 1 | - | 11 | Tuscarawas | 1 | 1 | 1 | - | 2 | 30 |
| Henry | - | - | - | - | - | 13 | Union | - | - | 1 | 2 | - | 29 |
| Highland | - | 2 | 1 | 2 | - | 33 | Van Wert | - | - | - | - | - | 9 |
| Hocking | - | - | 1 | - | - | 27 | Vinton | - | - | - | - | - | 4 |
| Holmes | - | - | - | - | - | 10 | Warren | 1 | 3 | 4 | 1 | 3 | 103 |
| Huron | 1 | - | 1 | - | 1 | 29 | Washington | - | 1 | - | 2 | 1 | 60 |
| Jackson | - | - | 1 | - | 1 | 15 | Wayne | 2 | 2 | 1 | 1 | - | 48 |
| Jefferson | 2 | - | 1 | 1 | 1 | 78 | Williams | - | - | 1 | 1 | - | 27 |
| Knox | - | - | 1 | 1 | 2 | 30 | Wood | 1 | 1 | - | - | 1 | 73 |
| Lake | 5 | 4 | 3 | 3 | 5 | 185 | Wyandot | - | - | - | 1 | - | 6 |
| Lawrence | 1 | - | 1 | - | - | 45 | No County | 11 | 6 | 5 | 4 | 4 | 743 |
| Licking | 2 | 4 | 5 | 4 | 3 | 162 | Ohio | 482 | 406 | 432 | 445 | 363 | 21,860 |

Notes:

AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a County reflects county of residence at time of initial diagnosis. Cases diagnosed while in a state or federal correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Persons Living with Diagnosed HIV Infection Reported in Ohio

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Ohio

| Characteristic | Living with diagnosed HIV infection in 2020 | | | Current Disease Status | | | |
|----------------------------------------------------|---------------------------------------------|---------------|-----|------------------------|-----|---------------|-----|
| | Rate ^a | No. | % | HIV (not AIDS) | | AIDS | |
| | | | | No. | % | No. | % |
| Sex at birth | | | | | | | |
| Males | 346.4 | 19,856 | 79% | 10,434 | 78% | 9,422 | 80% |
| Females | 87.9 | 5,240 | 21% | 2,889 | 22% | 2,351 | 20% |
| Age at end of year | | | | | | | |
| <13 | 2.7 | 49 | <1% | 40 | <1% | 9 | <1% |
| 13-14 | 7.1 | 21 | <1% | 15 | <1% | 6 | <1% |
| 15-19 | 15.3 | 114 | <1% | 94 | 1% | 20 | <1% |
| 20-24 | 97.1 | 726 | 3% | 605 | 5% | 121 | 1% |
| 25-29 | 243.0 | 1,927 | 8% | 1,477 | 11% | 450 | 4% |
| 30-34 | 350.2 | 2,693 | 11% | 1,840 | 14% | 853 | 7% |
| 35-39 | 322.4 | 2,339 | 9% | 1,444 | 11% | 895 | 8% |
| 40-44 | 346.9 | 2,377 | 9% | 1,325 | 10% | 1,052 | 9% |
| 45-49 | 380.4 | 2,634 | 10% | 1,295 | 10% | 1,339 | 11% |
| 50-54 | 474.5 | 3,452 | 14% | 1,530 | 11% | 1,922 | 16% |
| 55-64 | 400.9 | 6,362 | 25% | 2,676 | 20% | 3,686 | 31% |
| 65+ | 114.5 | 2,402 | 10% | 982 | 7% | 1,420 | 12% |
| Race/Ethnicity^b | | | | | | | |
| American Indian/Alaska Native | 66.3 | 15 | <1% | 9 | <1% | 6 | <1% |
| Asian/Pacific Islander | 56.7 | 173 | 1% | 113 | 1% | 60 | 1% |
| Black/African-American | 745.4 | 11,095 | 44% | 5,970 | 45% | 5,125 | 44% |
| Hispanic/Latinx | 373.7 | 1,824 | 7% | 912 | 7% | 912 | 8% |
| White | 118.7 | 10,828 | 43% | 5,694 | 43% | 5,134 | 44% |
| Multi-Race | 398.9 | 1,056 | 4% | 520 | 4% | 536 | 5% |
| Unknown | * | 105 | <1% | 105 | 1% | - | - |
| Race/Ethnicity^b and Sex at birth | | | | | | | |
| American Indian/Alaska Native Males | 96.8 | 11 | <1% | 8 | <1% | 3 | <1% |
| American Indian/Alaska Native Females | * | 4 | <1% | 1 | <1% | 3 | <1% |
| Asian/Pacific Islander Males | 90.5 | 134 | 1% | 89 | 1% | 45 | <1% |
| Asian/Pacific Islander Females | 24.8 | 39 | <1% | 24 | <1% | 15 | <1% |
| Black/African-American Males | 1,142.8 | 8,145 | 32% | 4,367 | 33% | 3,778 | 32% |
| Black/African-American Females | 380.3 | 2,950 | 12% | 1,603 | 12% | 1,347 | 11% |
| Hispanic/Latino Males | 570.8 | 1,423 | 6% | 734 | 6% | 689 | 6% |
| Hispanic/Latina Females | 167.9 | 401 | 2% | 178 | 1% | 223 | 2% |
| White Males | 206.0 | 9,229 | 37% | 4,753 | 36% | 4,476 | 38% |
| White Females | 34.4 | 1,599 | 6% | 941 | 7% | 658 | 6% |
| Multi-Race Males | 641.0 | 841 | 3% | 410 | 3% | 431 | 4% |
| Multi-Race Females | 161.0 | 215 | 1% | 110 | 1% | 105 | 1% |
| Unknown | * | 105 | <1% | 105 | 1% | - | - |
| Total | 214.6 | 25,096 | | 13,323 | | 11,773 | |

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Ohio

| Transmission Category ^a | Living with diagnosed HIV infection | | Current Disease Status | | | |
|----------------------------------------|-------------------------------------|------|------------------------|------|---------------|------|
| | in 2020 | | HIV (not AIDS) | | AIDS | |
| | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | |
| Male-to-male sexual contact | 13,409 | 68% | 7,020 | 68% | 6,389 | 68% |
| Injection drug use (IDU) | 803 | 4% | 358 | 3% | 445 | 5% |
| Male-to-male sexual contact and IDU | 1,037 | 5% | 459 | 4% | 578 | 6% |
| Heterosexual contact | 1,180 | 6% | 508 | 5% | 672 | 7% |
| Other/unknown | 3,300 | 17% | 2,021 | 19% | 1,279 | 14% |
| Subtotal | 19,729 | 100% | 10,366 | 100% | 9,363 | 100% |
| Female adult or adolescent | | | | | | |
| Injection drug use | 649 | 13% | 317 | 11% | 332 | 15% |
| Heterosexual contact | 3,984 | 78% | 2,117 | 76% | 1,867 | 82% |
| Other/unknown | 456 | 9% | 368 | 13% | 88 | 4% |
| Subtotal | 5,089 | 100% | 2,802 | 100% | 2,287 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | |
| Perinatal | 233 | 84% | 125 | 81% | 108 | 88% |
| Other/unknown | 45 | 16% | 30 | 19% | 15 | 12% |
| Subtotal | 278 | 100% | 155 | 100% | 123 | 100% |
| Total | 25,096 | | 13,323 | | 11,773 | |

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Ohio

| Exposure Category ^b | Living with diagnosed HIV infection | | Current Disease Status | | | |
|----------------------------------------------------------|-------------------------------------|-----|------------------------|-----|---------------|-----|
| | in 2020 | | HIV (not AIDS) | | AIDS | |
| | No. | % | No. | % | No. | % |
| Male-to-male sexual contact only | 12,725 | 51% | 6,730 | 51% | 5,995 | 51% |
| Injection drug use (IDU) only | 687 | 3% | 328 | 2% | 359 | 3% |
| Heterosexual contact only | 5,168 | 21% | 2,627 | 20% | 2,541 | 22% |
| Male-to-male sexual contact & IDU | 811 | 3% | 375 | 3% | 436 | 4% |
| IDU & Heterosexual contact | 765 | 3% | 347 | 3% | 418 | 4% |
| Male-to-male sexual contact & Heterosexual contact | 684 | 3% | 290 | 2% | 394 | 3% |
| Male-to-male sexual contact & IDU & Heterosexual contact | 226 | 1% | 84 | 1% | 142 | 1% |
| Perinatal exposure | 246 | 1% | 135 | 1% | 111 | 1% |
| Other/unknown | 3,784 | 15% | 2,407 | 18% | 1,377 | 12% |
| Total | 25,096 | | 13,323 | | 11,773 | |

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Ohio

Living with diagnosed HIV infection in 2020

| Age at end of year | American Indian/Alaska Native | | | Asian/Pacific Islander | | | Black/African-American | | | Hispanic/ Latinx ^a | | | White | | | Multi-Race | | | Unknown | |
|--------------------|-------------------------------|-----------|-----|------------------------|------------|-----|------------------------|---------------|-----|-------------------------------|--------------|-----|-------------------|---------------|-----|-------------------|--------------|-----|------------|-----|
| | Rate ^b | No. | % | Rate ^b | No. | % | Rate ^b | No. | % | Rate ^b | No. | % | Rate ^b | No. | % | Rate ^b | No. | % | No. | % |
| <13 | * | - | - | 13.3 | 7 | 4% | 8.8 | 25 | <1% | 4.7 | 6 | <1% | 0.9 | 11 | <1% | * | - | - | - | - |
| 13-14 | * | - | - | * | - | - | 31.8 | 14 | <1% | * | 1 | <1% | * | 4 | <1% | * | 2 | <1% | - | - |
| 15-19 | * | - | - | * | 4 | 2% | 81.0 | 85 | 1% | 11.2 | 5 | <1% | 3.1 | 17 | <1% | * | 3 | <1% | - | - |
| 20-24 | * | - | - | 37.6 | 9 | 5% | 437.9 | 471 | 4% | 141.6 | 57 | 3% | 28.3 | 155 | 1% | 130.2 | 34 | 3% | - | - |
| 25-29 | * | 1 | 7% | 59.0 | 15 | 9% | 915.4 | 1,142 | 10% | 346.7 | 137 | 8% | 91.2 | 529 | 5% | 484.9 | 103 | 10% | - | - |
| 30-34 | * | 2 | 13% | 65.9 | 20 | 12% | 1,343.3 | 1,472 | 13% | 511.3 | 192 | 11% | 149.5 | 860 | 8% | 997.5 | 146 | 14% | 1 | 1% |
| 35-39 | * | - | - | 65.9 | 19 | 11% | 1,231.3 | 1,154 | 10% | 484.9 | 176 | 10% | 159.4 | 884 | 8% | 973.0 | 106 | 10% | - | - |
| 40-44 | * | - | - | 77.4 | 18 | 10% | 1,162.3 | 1,004 | 9% | 670.4 | 215 | 12% | 193.6 | 1,031 | 10% | 1,130.6 | 107 | 10% | 2 | 2% |
| 45-49 | * | 4 | 27% | 154.8 | 33 | 19% | 1,274.7 | 1,086 | 10% | 873.2 | 233 | 13% | 210.9 | 1,159 | 11% | 1,316.3 | 108 | 10% | 11 | 10% |
| 50-54 | * | 3 | 20% | 127.0 | 22 | 13% | 1,585.7 | 1,330 | 12% | 1,046.1 | 227 | 12% | 291.6 | 1,739 | 16% | 1,731.3 | 116 | 11% | 15 | 14% |
| 55-64 | * | 2 | 13% | 64.1 | 17 | 10% | 1,394.9 | 2,429 | 22% | 1,283.4 | 427 | 23% | 238.0 | 3,182 | 29% | 2,087.9 | 257 | 24% | 48 | 46% |
| 65+ | * | 3 | 20% | 31.7 | 9 | 5% | 461.7 | 883 | 8% | 488.3 | 148 | 8% | 68.7 | 1,257 | 12% | 600.0 | 74 | 7% | 28 | 27% |
| Total | 66.3 | 15 | | 56.7 | 173 | | 745.4 | 11,095 | | 373.7 | 1,824 | | 118.7 | 10,828 | | 398.9 | 1,056 | | 105 | |

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Ohio

Living with diagnosed HIV infection in 2020

| Transmission Category ^c | American Indian/Alaska Native | | Asian/Pacific Islander | | Black/African-American | | Hispanic/ Latinx ^a | | White | | Multi-Race | | Unknown | |
|----------------------------------------|-------------------------------|------|------------------------|------|------------------------|------|-------------------------------|------|---------------|------|--------------|------|------------|------|
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | | | | | | | |
| Male-to-male sexual contact | 9 | 82% | 70 | 55% | 5,024 | 62% | 835 | 59% | 6,873 | 75% | 597 | 71% | 1 | 1% |
| Injection drug use (IDU) | 1 | 9% | 1 | 1% | 307 | 4% | 140 | 10% | 324 | 4% | 30 | 4% | - | - |
| Male-to-male sexual contact and IDU | - | - | 5 | 4% | 303 | 4% | 86 | 6% | 587 | 6% | 56 | 7% | - | - |
| Heterosexual contact | - | - | 8 | 6% | 707 | 9% | 127 | 9% | 284 | 3% | 53 | 6% | 1 | 1% |
| Other/unknown | 1 | 9% | 43 | 34% | 1,728 | 21% | 220 | 16% | 1,136 | 12% | 101 | 12% | 71 | 97% |
| Subtotal | 11 | 100% | 127 | 100% | 8,069 | 100% | 1,408 | 100% | 9,204 | 100% | 837 | 100% | 73 | 100% |
| Female adult or adolescent | | | | | | | | | | | | | | |
| Injection drug use | - | - | - | - | 211 | 7% | 53 | 14% | 357 | 23% | 28 | 13% | - | - |
| Heterosexual contact | 4 | 100% | 27 | 77% | 2,420 | 85% | 303 | 79% | 1,065 | 68% | 164 | 79% | 1 | 3% |
| Other/unknown | - | - | 8 | 23% | 232 | 8% | 28 | 7% | 142 | 9% | 16 | 8% | 30 | 97% |
| Subtotal | 4 | 100% | 35 | 100% | 2,863 | 100% | 384 | 100% | 1,564 | 100% | 208 | 100% | 31 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | | | |
| Perinatal | - | - | 7 | 64% | 141 | 87% | 30 | 94% | 45 | 75% | 10 | 91% | - | - |
| Other/unknown | - | - | 4 | 36% | 22 | 13% | 2 | 6% | 15 | 25% | 1 | 9% | 1 | 100% |
| Subtotal | - | - | 11 | 100% | 163 | 100% | 32 | 100% | 60 | 100% | 11 | 100% | 1 | 100% |
| Total | 15 | | 173 | | 11,095 | | 1,824 | | 10,828 | | 1,056 | | 105 | |

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Ohio

Living with diagnosed HIV infection in 2020

| Exposure Category ^d | American Indian/Alaska Native | | Asian/Pacific Islander | | Black/African-American | | Hispanic/ Latinx ^a | | White | | Multi-Race | | Unknown | |
|----------------------------------------------------------|-------------------------------|-----|------------------------|-----|------------------------|-----|-------------------------------|-----|---------------|-----|--------------|-----|------------|-----|
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sexual contact only | 9 | 60% | 67 | 39% | 4,683 | 42% | 783 | 43% | 6,623 | 61% | 559 | 53% | 1 | 1% |
| Injection drug use (IDU) only | 1 | 7% | 1 | 1% | 264 | 2% | 90 | 5% | 310 | 3% | 21 | 2% | - | - |
| Heterosexual contact only | 4 | 27% | 35 | 20% | 3,128 | 28% | 430 | 24% | 1,352 | 12% | 217 | 21% | 2 | 2% |
| Male-to-male sexual contact & IDU | - | - | 5 | 3% | 222 | 2% | 66 | 4% | 476 | 4% | 42 | 4% | - | - |
| IDU & Heterosexual contact | - | - | - | - | 254 | 2% | 103 | 6% | 371 | 3% | 37 | 4% | - | - |
| Male-to-male sexual contact & Heterosexual contact | - | - | 3 | 2% | 341 | 3% | 52 | 3% | 250 | 2% | 38 | 4% | - | - |
| Male-to-male sexual contact & IDU & Heterosexual contact | - | - | - | - | 81 | 1% | 20 | 1% | 111 | 1% | 14 | 1% | - | - |
| Perinatal exposure | - | - | 8 | 5% | 148 | 1% | 33 | 2% | 45 | 0% | 12 | 1% | - | - |
| Other/unknown | 1 | 7% | 54 | 31% | 1,974 | 18% | 247 | 14% | 1,290 | 12% | 116 | 11% | 102 | 97% |
| Total | 15 | | 173 | | 11,095 | | 1,824 | | 10,828 | | 1,056 | | 105 | |

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and county, Ohio

| County ^a | Living with diagnosed HIV infection in 2020 | | | Current Disease Status | | County ^a | Living with diagnosed HIV infection in 2020 | | | Current Disease Status | |
|---------------------|---------------------------------------------|-------|----------------|------------------------|-------------|---------------------|---------------------------------------------|---------------|----------------|------------------------|------|
| | Rate ^b | No. | HIV (not AIDS) | | AIDS | | Rate ^b | No. | HIV (not AIDS) | | AIDS |
| | | | No. | No. | | | | | No. | No. | |
| Adams | 79.9 | 22 | 7 | 15 | Logan | 77.2 | 35 | 20 | 15 | | |
| Allen | 154.0 | 157 | 83 | 74 | Lorain | 122.4 | 382 | 190 | 192 | | |
| Ashland | 39.4 | 21 | 7 | 14 | Lucas | 259.6 | 1,112 | 547 | 565 | | |
| Ashtabula | 121.2 | 117 | 56 | 61 | Madison | 89.8 | 40 | 22 | 18 | | |
| Athens | 91.6 | 60 | 32 | 28 | Mahoning | 219.4 | 496 | 272 | 224 | | |
| Auglaize | 46.0 | 21 | 11 | 10 | Marion | 111.1 | 72 | 29 | 43 | | |
| Belmont | 83.4 | 55 | 26 | 29 | Medina | 60.8 | 110 | 61 | 49 | | |
| Brown | 69.1 | 30 | 11 | 19 | Meigs | 44.1 | 10 | 5 | 5 | | |
| Butler | 126.5 | 488 | 238 | 250 | Mercer | 43.6 | 18 | 8 | 10 | | |
| Carroll | 40.9 | 11 | 7 | 4 | Miami | 73.5 | 79 | 35 | 44 | | |
| Champaign | 82.1 | 32 | 9 | 23 | Monroe | 73.6 | 10 | 4 | 6 | | |
| Clark | 122.8 | 164 | 77 | 87 | Montgomery | 297.2 | 1,580 | 796 | 784 | | |
| Clermont | 84.4 | 175 | 89 | 86 | Morgan | 104.9 | 15 | 6 | 9 | | |
| Clinton | 83.5 | 35 | 14 | 21 | Morrow | 59.3 | 21 | 13 | 8 | | |
| Columbiana | 59.3 | 60 | 35 | 25 | Muskingum | 88.4 | 76 | 41 | 35 | | |
| Coshocton | 57.6 | 21 | 7 | 14 | Noble | * | 3 | 2 | 1 | | |
| Crawford | 67.7 | 28 | 11 | 17 | Ottawa | 44.7 | 18 | 14 | 4 | | |
| Cuyahoga | 421.9 | 5,180 | 2,721 | 2,459 | Paulding | 42.9 | 8 | 3 | 5 | | |
| Darke | 58.6 | 30 | 19 | 11 | Perry | 104.9 | 38 | 19 | 19 | | |
| Defiance | 92.6 | 35 | 16 | 19 | Pickaway | 49.4 | 29 | 15 | 14 | | |
| Delaware | 63.2 | 135 | 74 | 61 | Pike | 79.4 | 22 | 9 | 13 | | |
| Erie | 131.6 | 97 | 51 | 46 | Portage | 73.8 | 120 | 74 | 46 | | |
| Fairfield | 119.6 | 191 | 108 | 83 | Preble | 46.5 | 19 | 9 | 10 | | |
| Fayette | 63.0 | 18 | 7 | 11 | Putnam | 35.7 | 12 | 6 | 6 | | |
| Franklin | 418.2 | 5,539 | 3,090 | 2,449 | Richland | 97.6 | 118 | 56 | 62 | | |
| Fulton | 54.9 | 23 | 11 | 12 | Ross | 89.0 | 68 | 35 | 33 | | |
| Gallia | 67.1 | 20 | 13 | 7 | Sandusky | 84.0 | 49 | 18 | 31 | | |
| Geauga | 40.7 | 38 | 16 | 22 | Scioto | 102.2 | 76 | 34 | 42 | | |
| Greene | 128.7 | 219 | 121 | 98 | Seneca | 45.5 | 25 | 12 | 13 | | |
| Guernsey | 82.5 | 32 | 17 | 15 | Shelby | 78.6 | 38 | 22 | 16 | | |
| Hamilton | 401.7 | 3,286 | 1,726 | 1,560 | Stark | 138.5 | 512 | 292 | 220 | | |
| Hancock | 88.9 | 67 | 35 | 32 | Summit | 198.4 | 1,069 | 596 | 473 | | |
| Hardin | 60.4 | 19 | 8 | 11 | Trumbull | 119.9 | 236 | 119 | 117 | | |
| Harrison | 59.9 | 9 | 4 | 5 | Tuscarawas | 39.2 | 36 | 17 | 19 | | |
| Henry | 29.7 | 8 | 5 | 3 | Union | 50.0 | 30 | 16 | 14 | | |
| Highland | 73.9 | 32 | 12 | 20 | Van Wert | 32.0 | 9 | 2 | 7 | | |
| Hocking | 99.7 | 28 | 13 | 15 | Vinton | 46.3 | 6 | 2 | 4 | | |
| Holmes | 13.6 | 6 | 2 | 4 | Warren | 79.3 | 189 | 103 | 86 | | |
| Huron | 75.9 | 44 | 24 | 20 | Washington | 92.2 | 55 | 31 | 24 | | |
| Jackson | 83.1 | 27 | 11 | 16 | Wayne | 69.1 | 80 | 35 | 45 | | |
| Jefferson | 109.3 | 71 | 32 | 39 | Williams | 98.5 | 36 | 12 | 24 | | |
| Knox | 67.3 | 42 | 18 | 24 | Wood | 67.9 | 89 | 53 | 36 | | |
| Lake | 95.0 | 218 | 112 | 106 | Wyandot | 27.6 | 6 | 3 | 3 | | |
| Lawrence | 143.8 | 85 | 55 | 30 | No County | * | 848 | 530 | 318 | | |
| Licking | 112.3 | 200 | 94 | 106 | Ohio | 214.6 | 25,096 | 13,323 | 11,773 | | |

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a County reflects current county of residence. Cases in a state or federal correctional facility or whose county is unknown are included in No County.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Ohio, 2016-2020

| Characteristic | Living with Diagnosed HIV Infection | | | | | | | | | | | | | | |
|----------------------------------------------------|-------------------------------------|---------------|-----|-------------------|---------------|-----|-------------------|---------------|-----|-------------------|---------------|-----|-------------------|---------------|-----|
| | 2016 | | | 2017 | | | 2018 | | | 2019 | | | 2020 | | |
| | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % |
| Sex at birth | | | | | | | | | | | | | | | |
| Males | 315.8 | 17,976 | 80% | 324.6 | 18,543 | 79% | 331.0 | 18,969 | 79% | 338.2 | 19,382 | 79% | 346.4 | 19,856 | 79% |
| Females | 78.2 | 4,634 | 20% | 80.6 | 4,792 | 21% | 82.6 | 4,924 | 21% | 85.5 | 5,093 | 21% | 87.9 | 5,240 | 21% |
| Age at end of year | | | | | | | | | | | | | | | |
| <13 | 3.0 | 56 | <1% | 3.0 | 55 | <1% | 3.1 | 58 | <1% | 2.8 | 51 | <1% | 2.7 | 49 | <1% |
| 13-14 | 6.4 | 19 | <1% | 4.7 | 14 | <1% | 3.7 | 11 | <1% | 6.4 | 19 | <1% | 7.1 | 21 | <1% |
| 15-19 | 16.9 | 130 | 1% | 16.4 | 126 | 1% | 16.4 | 125 | 1% | 15.1 | 114 | <1% | 15.3 | 114 | <1% |
| 20-24 | 113.8 | 878 | 4% | 109.5 | 838 | 4% | 105.5 | 801 | 3% | 104.1 | 782 | 3% | 97.1 | 726 | 3% |
| 25-29 | 257.6 | 2,014 | 9% | 268.4 | 2,138 | 9% | 262.1 | 2,115 | 9% | 250.6 | 2,009 | 8% | 243.0 | 1,927 | 8% |
| 30-34 | 267.3 | 1,938 | 9% | 286.1 | 2,081 | 9% | 309.0 | 2,282 | 10% | 330.8 | 2,490 | 10% | 350.2 | 2,693 | 11% |
| 35-39 | 299.5 | 2,106 | 9% | 301.9 | 2,159 | 9% | 301.4 | 2,183 | 9% | 311.3 | 2,255 | 9% | 322.4 | 2,339 | 9% |
| 40-44 | 329.9 | 2,233 | 10% | 334.8 | 2,230 | 10% | 338.7 | 2,259 | 9% | 342.4 | 2,306 | 9% | 346.9 | 2,377 | 9% |
| 45-49 | 430.5 | 3,225 | 14% | 412.8 | 3,072 | 13% | 398.5 | 2,919 | 12% | 391.8 | 2,796 | 11% | 380.4 | 2,634 | 10% |
| 50-54 | 485.2 | 3,875 | 17% | 487.5 | 3,779 | 16% | 488.1 | 3,672 | 15% | 483.3 | 3,547 | 14% | 474.5 | 3,452 | 14% |
| 55-64 | 300.8 | 4,829 | 21% | 327.2 | 5,276 | 23% | 351.6 | 5,666 | 24% | 374.7 | 6,012 | 25% | 400.9 | 6,362 | 25% |
| 65+ | 69.3 | 1,307 | 6% | 80.6 | 1,567 | 7% | 90.3 | 1,802 | 8% | 102.3 | 2,094 | 9% | 114.5 | 2,402 | 10% |
| Race/Ethnicity^b | | | | | | | | | | | | | | | |
| American Indian/Alaska Native | 53.6 | 12 | <1% | 53.2 | 12 | <1% | 57.1 | 13 | <1% | 61.0 | 14 | <1% | 66.3 | 15 | <1% |
| Asian/Pacific Islander | 47.3 | 121 | 1% | 51.1 | 139 | 1% | 51.8 | 149 | 1% | 54.4 | 159 | 1% | 56.7 | 173 | 1% |
| Black/African-American | 678.5 | 9,818 | 43% | 693.1 | 10,140 | 43% | 706.2 | 10,412 | 44% | 725.1 | 10,741 | 44% | 745.4 | 11,095 | 44% |
| Hispanic/Latinx | 370.2 | 1,572 | 7% | 376.7 | 1,661 | 7% | 374.7 | 1,727 | 7% | 379.4 | 1,785 | 7% | 373.7 | 1,824 | 7% |
| White | 108.5 | 10,015 | 44% | 111.5 | 10,277 | 44% | 113.7 | 10,460 | 44% | 115.8 | 10,615 | 43% | 118.7 | 10,828 | 43% |
| Multi-Race | 414.5 | 971 | 4% | 417.5 | 1,005 | 4% | 416.2 | 1,031 | 4% | 418.7 | 1,060 | 4% | 398.9 | 1,056 | 4% |
| Unknown | * | 101 | <1% | * | 101 | <1% | * | 101 | <1% | * | 101 | <1% | * | 105 | <1% |
| Race/Ethnicity^b and Sex at birth | | | | | | | | | | | | | | | |
| American Indian/Alaska Native Males | 71.1 | 8 | <1% | 70.7 | 8 | <1% | 78.9 | 9 | <1% | 87.0 | 10 | <1% | 96.8 | 11 | <1% |
| American Indian/Alaska Native Females | * | 4 | <1% | * | 4 | <1% | * | 4 | <1% | * | 4 | <1% | * | 4 | <1% |
| Asian/Pacific Islander Males | 76.2 | 94 | <1% | 81.2 | 106 | <1% | 82.0 | 114 | <1% | 85.5 | 121 | <1% | 90.5 | 134 | 1% |
| Asian/Pacific Islander Females | 20.4 | 27 | <1% | 23.3 | 33 | <1% | 23.5 | 35 | <1% | 25.2 | 38 | <1% | 24.8 | 39 | <1% |
| Black/African-American Males | 1,040.5 | 7,205 | 32% | 1,065.4 | 7,458 | 32% | 1,085.2 | 7,664 | 32% | 1,113.1 | 7,896 | 32% | 1,142.8 | 8,145 | 32% |
| Black/African-American Females | 346.3 | 2,613 | 12% | 351.6 | 2,682 | 11% | 357.7 | 2,748 | 12% | 368.5 | 2,845 | 12% | 380.3 | 2,950 | 12% |
| Hispanic/Latino Males | 549.4 | 1,197 | 5% | 561.5 | 1,270 | 5% | 560.6 | 1,328 | 6% | 575.3 | 1,386 | 6% | 570.8 | 1,423 | 6% |
| Hispanic/Latina Females | 181.4 | 375 | 2% | 182.1 | 391 | 2% | 178.1 | 399 | 2% | 173.8 | 399 | 2% | 167.9 | 401 | 2% |
| White Males | 190.4 | 8,629 | 38% | 195.0 | 8,827 | 38% | 198.4 | 8,958 | 37% | 201.1 | 9,055 | 37% | 206.0 | 9,229 | 37% |
| White Females | 29.5 | 1,386 | 6% | 30.9 | 1,450 | 6% | 32.1 | 1,502 | 6% | 33.4 | 1,560 | 6% | 34.4 | 1,599 | 6% |
| Multi-Race Males | 667.4 | 771 | 3% | 674.6 | 802 | 3% | 671.7 | 824 | 3% | 672.5 | 842 | 3% | 641.0 | 841 | 3% |
| Multi-Race Females | 168.5 | 200 | 1% | 166.6 | 203 | 1% | 165.6 | 207 | 1% | 170.4 | 218 | 1% | 161.0 | 215 | 1% |
| Unknown | * | 101 | <1% | * | 101 | <1% | * | 101 | <1% | * | 101 | <1% | * | 105 | <1% |
| Total | 194.7 | 22,610 | | 200.2 | 23,335 | | 204.4 | 23,893 | | 209.4 | 24,475 | | 214.6 | 25,096 | |

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Ohio, 2016-2020

| Transmission Category ^a | Living with Diagnosed HIV Infection | | | | | | | | | |
|----------------------------------------|-------------------------------------|------|---------------|------|---------------|------|---------------|------|---------------|------|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | | | | | |
| Male-to-male sexual contact | 12,354 | 69% | 12,754 | 69% | 13,023 | 69% | 13,184 | 68% | 13,409 | 68% |
| Injection drug use (IDU) | 723 | 4% | 750 | 4% | 775 | 4% | 797 | 4% | 803 | 4% |
| Male-to-male sexual contact and IDU | 951 | 5% | 979 | 5% | 984 | 5% | 1,006 | 5% | 1,037 | 5% |
| Heterosexual contact | 1,128 | 6% | 1,132 | 6% | 1,154 | 6% | 1,175 | 6% | 1,180 | 6% |
| Other/unknown | 2,709 | 15% | 2,819 | 15% | 2,920 | 15% | 3,101 | 16% | 3,300 | 17% |
| Subtotal | 17,865 | 100% | 18,434 | 100% | 18,856 | 100% | 19,263 | 100% | 19,729 | 100% |
| Female adult or adolescent | | | | | | | | | | |
| Injection drug use | 552 | 12% | 574 | 12% | 587 | 12% | 630 | 13% | 649 | 13% |
| Heterosexual contact | 3,562 | 79% | 3,689 | 79% | 3,796 | 79% | 3,898 | 79% | 3,984 | 78% |
| Other/unknown | 390 | 9% | 394 | 8% | 403 | 8% | 420 | 8% | 456 | 9% |
| Subtotal | 4,504 | 100% | 4,657 | 100% | 4,786 | 100% | 4,948 | 100% | 5,089 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | |
| Perinatal | 209 | 87% | 210 | 86% | 216 | 86% | 225 | 85% | 233 | 84% |
| Other/unknown | 32 | 13% | 34 | 14% | 35 | 14% | 39 | 15% | 45 | 16% |
| Subtotal | 241 | 100% | 244 | 100% | 251 | 100% | 264 | 100% | 278 | 100% |
| Total | 22,610 | | 23,335 | | 23,893 | | 24,475 | | 25,096 | |

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Ohio, 2016-2020

| Exposure Category ^b | Living with Diagnosed HIV Infection | | | | | | | | | |
|----------------------------------------------------------|-------------------------------------|-----|---------------|-----|---------------|-----|---------------|-----|---------------|-----|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sexual contact only | 11,636 | 51% | 12,047 | 52% | 12,315 | 52% | 12,485 | 51% | 12,725 | 51% |
| Injection drug use (IDU) only | 616 | 3% | 633 | 3% | 646 | 3% | 674 | 3% | 687 | 3% |
| Heterosexual contact only | 4,696 | 21% | 4,827 | 21% | 4,955 | 21% | 5,077 | 21% | 5,168 | 21% |
| Male-to-male sexual contact & IDU | 732 | 3% | 752 | 3% | 763 | 3% | 783 | 3% | 811 | 3% |
| IDU & Heterosexual contact | 659 | 3% | 691 | 3% | 716 | 3% | 753 | 3% | 765 | 3% |
| Male-to-male sexual contact & Heterosexual contact | 718 | 3% | 707 | 3% | 708 | 3% | 699 | 3% | 684 | 3% |
| Male-to-male sexual contact & IDU & Heterosexual contact | 219 | 1% | 227 | 1% | 221 | 1% | 223 | 1% | 226 | 1% |
| Perinatal exposure | 223 | 1% | 221 | 1% | 227 | 1% | 238 | 1% | 246 | 1% |
| Other/unknown | 3,111 | 14% | 3,230 | 14% | 3,342 | 14% | 3,543 | 14% | 3,784 | 15% |
| Total | 22,610 | | 23,335 | | 23,893 | | 24,475 | | 25,096 | |

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and county, Ohio, 2016-2020

| County ^a | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | County ^a | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | |
|---------------------|-------------------|-------|-------------------|-------|-------------------|-------|-------------------|-------|-------------------|-------|---------------------|-------------------|---------------|-------------------|---------------|-------------------|---------------|-------------------|---------------|-------------------|---------------|
| | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. |
| Adams | 75.2 | 21 | 72.1 | 20 | 72.1 | 20 | 68.6 | 19 | 79.9 | 22 | Logan | 53.1 | 24 | 57.4 | 26 | 63.9 | 29 | 70.1 | 32 | 77.2 | 35 |
| Allen | 150.4 | 156 | 150.2 | 155 | 149.0 | 153 | 153.4 | 157 | 154.0 | 157 | Lorain | 109.7 | 336 | 109.4 | 337 | 115.0 | 356 | 116.2 | 360 | 122.4 | 382 |
| Ashland | 48.5 | 26 | 48.5 | 26 | 46.5 | 25 | 37.4 | 20 | 39.4 | 21 | Lucas | 235.2 | 1,017 | 241.8 | 1,042 | 246.6 | 1,060 | 254.0 | 1,088 | 259.6 | 1,112 |
| Ashtabula | 101.8 | 100 | 109.4 | 107 | 113.9 | 111 | 113.1 | 110 | 121.2 | 117 | Madison | 64.5 | 28 | 65.9 | 29 | 76.6 | 34 | 89.4 | 40 | 89.8 | 40 |
| Athens | 78.6 | 52 | 73.6 | 49 | 79.0 | 52 | 79.6 | 52 | 91.6 | 60 | Mahoning | 196.1 | 451 | 204.1 | 469 | 205.1 | 471 | 209.0 | 478 | 219.4 | 496 |
| Auglaize | 50.1 | 23 | 52.4 | 24 | 43.7 | 20 | 48.2 | 22 | 46.0 | 21 | Marion | 75.3 | 49 | 81.6 | 53 | 87.3 | 57 | 90.6 | 59 | 111.1 | 72 |
| Belmont | 62.6 | 43 | 69.1 | 47 | 74.1 | 50 | 79.1 | 53 | 83.4 | 55 | Medina | 50.2 | 89 | 52.1 | 93 | 57.5 | 103 | 56.2 | 101 | 60.8 | 110 |
| Brown | 57.1 | 25 | 57.4 | 25 | 64.2 | 28 | 69.1 | 30 | 69.1 | 30 | Meigs | 47.6 | 11 | 47.7 | 11 | 47.6 | 11 | 52.4 | 12 | 44.1 | 10 |
| Butler | 110.5 | 417 | 114.0 | 434 | 120.0 | 459 | 125.3 | 480 | 126.5 | 488 | Mercer | 24.4 | 10 | 29.4 | 12 | 41.5 | 17 | 34.0 | 14 | 43.6 | 18 |
| Carroll | 39.8 | 11 | 32.9 | 9 | 33.2 | 9 | 29.7 | 8 | 40.9 | 11 | Miami | 64.0 | 67 | 62.8 | 66 | 66.8 | 71 | 73.8 | 79 | 73.5 | 79 |
| Champaign | 80.0 | 31 | 79.8 | 31 | 74.8 | 29 | 77.2 | 30 | 82.1 | 32 | Monroe | 56.3 | 8 | 50.2 | 7 | 65.3 | 9 | 73.2 | 10 | 73.6 | 10 |
| Clark | 102.4 | 138 | 101.1 | 136 | 106.3 | 143 | 114.9 | 154 | 122.8 | 164 | Montgomery | 269.6 | 1,432 | 278.8 | 1,482 | 284.2 | 1,513 | 297.5 | 1,582 | 297.2 | 1,580 |
| Clermont | 67.0 | 136 | 69.0 | 141 | 78.4 | 161 | 78.5 | 162 | 84.4 | 175 | Morgan | 94.6 | 14 | 74.8 | 11 | 82.2 | 12 | 89.6 | 13 | 104.9 | 15 |
| Clinton | 83.5 | 35 | 83.3 | 35 | 88.0 | 37 | 88.2 | 37 | 83.5 | 35 | Morrow | 51.4 | 18 | 48.6 | 17 | 54.1 | 19 | 56.6 | 20 | 59.3 | 21 |
| Columbiana | 58.8 | 61 | 61.1 | 63 | 57.5 | 59 | 55.0 | 56 | 59.3 | 60 | Muskingum | 81.3 | 70 | 78.9 | 68 | 88.2 | 76 | 90.5 | 78 | 88.4 | 76 |
| Coshocton | 60.1 | 22 | 54.7 | 20 | 57.3 | 21 | 51.9 | 19 | 57.6 | 21 | Noble | * | 1 | * | 3 | * | 2 | * | 4 | * | 3 |
| Crawford | 47.5 | 20 | 57.5 | 24 | 55.4 | 23 | 60.2 | 25 | 67.7 | 28 | Ottawa | 34.5 | 14 | 39.4 | 16 | 41.7 | 17 | 32.1 | 13 | 44.7 | 18 |
| Cuyahoga | 383.4 | 4,790 | 394.1 | 4,921 | 403.4 | 5,018 | 409.9 | 5,062 | 421.9 | 5,180 | Paulding | 47.7 | 9 | 47.8 | 9 | 42.6 | 8 | 37.5 | 7 | 42.9 | 8 |
| Darke | 71.5 | 37 | 64.0 | 33 | 58.5 | 30 | 64.6 | 33 | 58.6 | 30 | Perry | 89.1 | 32 | 91.6 | 33 | 105.5 | 38 | 113.5 | 41 | 104.9 | 38 |
| Defiance | 76.0 | 29 | 76.0 | 29 | 81.2 | 31 | 89.3 | 34 | 92.6 | 35 | Pickaway | 55.6 | 32 | 55.3 | 32 | 49.9 | 29 | 49.6 | 29 | 49.4 | 29 |
| Delaware | 71.8 | 141 | 69.3 | 139 | 63.5 | 130 | 68.4 | 143 | 63.2 | 135 | Pike | 81.7 | 23 | 74.3 | 21 | 81.9 | 23 | 86.4 | 24 | 79.4 | 22 |
| Erie | 111.8 | 84 | 113.6 | 85 | 123.3 | 92 | 119.8 | 89 | 131.6 | 97 | Portage | 60.5 | 98 | 61.0 | 99 | 63.8 | 104 | 68.9 | 112 | 73.8 | 120 |
| Fairfield | 100.3 | 153 | 109.2 | 169 | 115.5 | 180 | 121.2 | 191 | 119.6 | 191 | Preble | 55.8 | 23 | 55.9 | 23 | 53.7 | 22 | 58.7 | 24 | 46.5 | 19 |
| Fayette | 66.3 | 19 | 83.5 | 24 | 87.2 | 25 | 80.6 | 23 | 63.0 | 18 | Putnam | 23.5 | 8 | 23.6 | 8 | 32.6 | 11 | 32.5 | 11 | 35.7 | 12 |
| Franklin | 390.4 | 4,937 | 395.1 | 5,105 | 395.9 | 5,187 | 404.6 | 5,327 | 418.2 | 5,539 | Richland | 87.5 | 106 | 92.9 | 112 | 95.0 | 115 | 94.9 | 115 | 97.6 | 118 |
| Fulton | 51.7 | 22 | 52.0 | 22 | 54.4 | 23 | 52.2 | 22 | 54.9 | 23 | Ross | 83.1 | 64 | 85.4 | 66 | 78.0 | 60 | 80.9 | 62 | 89.0 | 68 |
| Gallia | 63.3 | 19 | 60.1 | 18 | 56.7 | 17 | 63.5 | 19 | 67.1 | 20 | Sandusky | 75.8 | 45 | 81.1 | 48 | 78.2 | 46 | 83.7 | 49 | 84.0 | 49 |
| Geauga | 44.7 | 42 | 46.8 | 44 | 39.3 | 37 | 41.6 | 39 | 40.7 | 38 | Scioto | 106.5 | 81 | 111.9 | 85 | 108.6 | 82 | 106.2 | 80 | 102.2 | 76 |
| Greene | 108.6 | 179 | 110.3 | 184 | 116.7 | 196 | 121.3 | 205 | 128.7 | 219 | Seneca | 36.1 | 20 | 43.4 | 24 | 48.9 | 27 | 47.1 | 26 | 45.5 | 25 |
| Guernsey | 71.7 | 28 | 66.5 | 26 | 71.8 | 28 | 72.0 | 28 | 82.5 | 32 | Shelby | 72.0 | 35 | 77.9 | 38 | 74.0 | 36 | 67.9 | 33 | 78.6 | 38 |
| Hamilton | 369.7 | 2,991 | 387.9 | 3,157 | 388.5 | 3,173 | 399.0 | 3,262 | 401.7 | 3,286 | Stark | 125.0 | 467 | 128.3 | 478 | 137.0 | 509 | 140.3 | 520 | 138.5 | 512 |
| Hancock | 63.3 | 48 | 71.3 | 54 | 79.0 | 60 | 87.1 | 66 | 88.9 | 67 | Summit | 169.9 | 918 | 177.0 | 958 | 184.3 | 999 | 184.8 | 1,000 | 198.4 | 1,069 |
| Hardin | 60.4 | 19 | 63.8 | 20 | 63.5 | 20 | 60.6 | 19 | 60.4 | 19 | Trumbull | 104.5 | 211 | 108.8 | 218 | 113.3 | 225 | 112.6 | 223 | 119.9 | 236 |
| Harrison | * | 4 | 52.6 | 8 | 79.1 | 12 | 66.5 | 10 | 59.9 | 9 | Tuscarawas | 24.9 | 23 | 33.6 | 31 | 32.5 | 30 | 33.7 | 31 | 39.2 | 36 |
| Henry | 36.2 | 10 | 40.5 | 11 | 40.6 | 11 | 29.6 | 8 | 29.7 | 8 | Union | 59.5 | 33 | 52.9 | 30 | 48.4 | 28 | 45.8 | 27 | 50.0 | 30 |
| Highland | 48.8 | 21 | 60.5 | 26 | 62.7 | 27 | 67.2 | 29 | 73.9 | 32 | Van Wert | 42.3 | 12 | 49.6 | 14 | 46.0 | 13 | 42.4 | 12 | 32.0 | 9 |
| Hocking | 95.3 | 27 | 98.3 | 28 | 109.2 | 31 | 106.1 | 30 | 99.7 | 28 | Vinton | 54.2 | 7 | * | 4 | 38.1 | 5 | 45.9 | 6 | 46.3 | 6 |
| Holmes | 13.7 | 6 | 15.9 | 7 | 11.4 | 5 | 13.6 | 6 | 13.6 | 6 | Warren | 58.6 | 133 | 61.2 | 140 | 66.8 | 155 | 75.0 | 176 | 79.3 | 189 |
| Huron | 61.6 | 36 | 70.1 | 41 | 70.1 | 41 | 72.1 | 42 | 75.9 | 44 | Washington | 67.6 | 41 | 79.4 | 48 | 79.8 | 48 | 88.5 | 53 | 92.2 | 55 |
| Jackson | 70.8 | 23 | 74.0 | 24 | 92.6 | 30 | 89.5 | 29 | 83.1 | 27 | Wayne | 64.4 | 75 | 65.5 | 76 | 63.8 | 74 | 65.7 | 76 | 69.1 | 80 |
| Jefferson | 92.9 | 62 | 91.9 | 61 | 94.3 | 62 | 101.0 | 66 | 109.3 | 71 | Williams | 72.9 | 27 | 92.4 | 34 | 89.7 | 33 | 92.7 | 34 | 98.5 | 36 |
| Knox | 49.3 | 30 | 50.6 | 31 | 63.0 | 39 | 64.2 | 40 | 67.3 | 42 | Wood | 59.1 | 77 | 58.2 | 76 | 63.5 | 83 | 66.5 | 87 | 67.9 | 89 |
| Lake | 75.7 | 173 | 85.2 | 196 | 86.3 | 199 | 88.6 | 204 | 95.0 | 218 | Wyandot | 22.6 | 5 | * | 4 | 22.8 | 5 | 32.2 | 7 | 27.6 | 6 |
| Lawrence | 95.3 | 58 | 96.3 | 58 | 101.9 | 61 | 114.4 | 68 | 143.8 | 85 | No County | * | 882 | * | 833 | * | 874 | * | 906 | * | 848 |
| Licking | 104.0 | 179 | 106.1 | 184 | 107.5 | 189 | 109.1 | 193 | 112.3 | 200 | Ohio | 194.7 | 22,610 | 200.2 | 23,335 | 204.4 | 23,893 | 209.4 | 24,475 | 214.6 | 25,096 |

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a County reflects current county of residence. Cases in a state or federal correctional facility or whose county is unknown are included in No County.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported transgender persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Ohio

| Characteristic | Living with diagnosed HIV infection in 2020 | | Current Disease Status | | | |
|------------------------------------------------------|---------------------------------------------|-----|------------------------|-----|-----------|-----|
| | | | HIV (not AIDS) | | AIDS | |
| | No. | % | No. | % | No. | % |
| Current Gender | | | | | | |
| Transgender Women | 256 | 96% | 165 | 96% | 91 | 95% |
| Transgender Men | 11 | 4% | 6 | 4% | 5 | 5% |
| Age at diagnosis (yr) | | | | | | |
| <13 | - | - | - | - | - | - |
| 13-14 | - | - | - | - | - | - |
| 15-19 | 4 | 1% | 4 | 2% | - | - |
| 20-24 | 20 | 7% | 16 | 9% | 4 | 4% |
| 25-29 | 52 | 19% | 43 | 25% | 9 | 9% |
| 30-34 | 75 | 28% | 53 | 31% | 22 | 23% |
| 35-39 | 37 | 14% | 24 | 14% | 13 | 14% |
| 40-44 | 18 | 7% | 11 | 6% | 7 | 7% |
| 45-49 | 17 | 6% | 7 | 4% | 10 | 10% |
| 50-54 | 15 | 6% | 4 | 2% | 11 | 11% |
| 55-64 | 21 | 8% | 6 | 4% | 15 | 16% |
| 65+ | 8 | 3% | 3 | 2% | 5 | 5% |
| Race/Ethnicity^a | | | | | | |
| American Indian/Alaska Native | - | - | - | - | - | - |
| Asian/Pacific Islander | - | - | - | - | - | - |
| Black/African-American | 187 | 70% | 127 | 74% | 60 | 63% |
| Hispanic/Latinx | 17 | 6% | 11 | 6% | 6 | 6% |
| White | 47 | 18% | 25 | 15% | 22 | 23% |
| Multi-Race | 16 | 6% | 8 | 5% | 8 | 8% |
| Unknown | - | - | - | - | - | - |
| Race/Ethnicity^a and Current Gender | | | | | | |
| American Indian/Alaska Native Transgender Women | - | - | - | - | - | - |
| American Indian/Alaska Native Transgender Men | - | - | - | - | - | - |
| Asian/Pacific Islander Transgender Women | - | - | - | - | - | - |
| Asian/Pacific Islander Transgender Men | - | - | - | - | - | - |
| Black/African-American Transgender Women | 178 | 67% | 123 | 72% | 55 | 57% |
| Black/African-American Transgender Men | 9 | 3% | 4 | 2% | 5 | 5% |
| Hispanic/Latina Transgender Women | 17 | 6% | 11 | 6% | 6 | 6% |
| Hispanic/Latino Transgender Men | - | - | - | - | - | - |
| White Transgender Women | 46 | 17% | 24 | 14% | 22 | 23% |
| White Transgender Men | 1 | <1% | 1 | 1% | - | - |
| Multi-Race Transgender Women | 15 | 6% | 7 | 4% | 8 | 8% |
| Multi-Race Transgender Men | 1 | <1% | 1 | 1% | - | - |
| Unknown | - | - | - | - | - | - |
| Total | 267 | | 171 | | 96 | |

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020.

Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Transgender is a term used to describe persons whose current gender identity is different than their sex (male or female) assigned at birth. Transgender women identify as female, but were assigned male at birth and transgender men identify as male, but were assigned female at birth.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported transgender persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Ohio

| Transmission Category ^a | Living with diagnosed HIV infection in 2020 | | Current Disease Status | | | |
|------------------------------------|---------------------------------------------|-----|------------------------|-----|-----------|-----|
| | | | HIV (not AIDS) | | AIDS | |
| | No. | % | No. | % | No. | % |
| Transgender Women | | | | | | |
| Sexual contact | 229 | 89% | 149 | 90% | 80 | 88% |
| Injection drug use (IDU) | 1 | <1% | 1 | 1% | - | - |
| Sexual contact and IDU | 20 | 8% | 11 | 7% | 9 | 10% |
| Other/unknown | 6 | 2% | 4 | 2% | 2 | 2% |
| Subtotal | 256 | | 165 | | 91 | |
| Transgender Men | | | | | | |
| Injection drug use | 1 | 9% | - | - | 1 | 20% |
| Sexual contact | 8 | 73% | 4 | 67% | 4 | 80% |
| Other/unknown | 2 | 18% | 2 | 33% | - | - |
| Subtotal | 11 | | 6 | | 5 | |
| Total | 267 | | 171 | | 96 | |

Reported transgender persons living with diagnosed HIV infection in 2020 by current disease status and region, Ohio

| Region ^b | Living with diagnosed HIV infection in 2020 | | Current Disease Status | | | |
|---------------------------|---------------------------------------------|-----|------------------------|-----|-----------|-----|
| | | | HIV (not AIDS) | | AIDS | |
| | No. | % | No. | % | No. | % |
| Region 1 | 10 | 4% | 5 | 3% | 5 | 5% |
| Region 2 | 2 | 1% | - | - | 2 | 2% |
| Part A-Cleveland/Region 3 | 73 | 27% | 52 | 30% | 21 | 22% |
| Region 4 | 15 | 6% | 10 | 6% | 5 | 5% |
| Region 5 | 5 | 2% | 2 | 1% | 3 | 3% |
| Region 6 | 4 | 1% | 1 | 1% | 3 | 3% |
| Region 7 | 2 | 1% | 2 | 1% | - | - |
| Region 8 | 52 | 19% | 31 | 18% | 21 | 22% |
| Region 9 | 25 | 9% | 16 | 9% | 9 | 9% |
| Region 10 | 4 | 1% | 3 | 2% | 1 | 1% |
| Part A-Columbus/Region 11 | 65 | 24% | 43 | 25% | 22 | 23% |
| No County | 10 | 4% | 6 | 4% | 4 | 4% |
| Total | 267 | | 171 | | 96 | |

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020.

Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Transgender is a term used to describe persons whose current gender identity is different than their sex (male or female) assigned at birth. Transgender women identify as female, but were assigned male at birth and transgender men identify as male, but were assigned female at birth.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. For the purposes of displaying transmission categories by current gender, the CDC-defined categories have been modified slightly.

^b Region reflects HIV Prevention Planning Region of residence at time of initial diagnosis. Cases diagnosed while in a state or federal correctional facility or whose county is unknown are included in No County.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Deaths among Reported Persons with Diagnosed HIV Infection in Ohio

As of June 30, 2021

Trends in deaths among reported persons with diagnosed HIV infection by year of death (2016-2020) and cumulative deaths by selected characteristics, Ohio

| Characteristic | Year of Death among Persons with Diagnosed HIV Infection | | | | | | | | | | | | | | | Cumulative Deaths | |
|----------------------------------------------------|----------------------------------------------------------|------------|-----|-------------------|------------|-----|-------------------|------------|-----|-------------------|------------|-----|-------------------|------------|-----|-------------------|-----|
| | 2016 | | | 2017 | | | 2018 | | | 2019 | | | 2020 | | | | |
| | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % | Rate ^a | No. | % | No. | % |
| Sex at birth | | | | | | | | | | | | | | | | | |
| Males | 4.7 | 270 | 81% | 5.1 | 292 | 80% | 5.0 | 289 | 84% | 5.1 | 293 | 79% | 5.7 | 324 | 81% | 12,626 | 87% |
| Females | 1.1 | 63 | 19% | 1.2 | 71 | 20% | 0.9 | 55 | 16% | 1.3 | 80 | 21% | 1.3 | 77 | 19% | 1,955 | 13% |
| Age at death (yr) | | | | | | | | | | | | | | | | | |
| <13 | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 73 | 1% |
| 13-14 | * | - | - | * | - | - | * | - | - | * | 1 | <1% | * | - | - | 3 | <1% |
| 15-19 | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 32 | <1% |
| 20-24 | 0.6 | 5 | 2% | 0.8 | 6 | 2% | * | 4 | 1% | 0.8 | 6 | 2% | * | 4 | 1% | 224 | 2% |
| 25-29 | 1.4 | 11 | 3% | 2.1 | 17 | 5% | 1.9 | 15 | 4% | 1.7 | 14 | 4% | 1.9 | 15 | 4% | 1,089 | 7% |
| 30-34 | 3.3 | 24 | 7% | 2.9 | 21 | 6% | 2.0 | 15 | 4% | 2.5 | 19 | 5% | 2.7 | 21 | 5% | 2,107 | 14% |
| 35-39 | 3.0 | 21 | 6% | 4.1 | 29 | 8% | 2.2 | 16 | 5% | 3.2 | 23 | 6% | 3.4 | 25 | 6% | 2,502 | 17% |
| 40-44 | 3.4 | 23 | 7% | 3.3 | 22 | 6% | 4.3 | 29 | 8% | 2.5 | 17 | 5% | 3.8 | 26 | 6% | 2,359 | 16% |
| 45-49 | 5.2 | 39 | 12% | 5.2 | 39 | 11% | 6.0 | 44 | 13% | 6.3 | 45 | 12% | 4.8 | 33 | 8% | 1,996 | 14% |
| 50-54 | 9.3 | 74 | 22% | 7.4 | 57 | 16% | 9.0 | 68 | 20% | 8.0 | 59 | 16% | 6.9 | 50 | 12% | 1,581 | 11% |
| 55-64 | 5.4 | 87 | 26% | 7.1 | 115 | 32% | 6.1 | 99 | 29% | 7.0 | 112 | 30% | 8.8 | 139 | 35% | 1,810 | 12% |
| 65+ | 2.6 | 49 | 15% | 2.9 | 57 | 16% | 2.7 | 54 | 16% | 3.8 | 77 | 21% | 4.2 | 88 | 22% | 805 | 6% |
| Race/Ethnicity^b | | | | | | | | | | | | | | | | | |
| American Indian/Alaska Native | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 13 | <1% |
| Asian/Pacific Islander | * | - | - | * | - | - | * | 1 | <1% | * | - | - | * | 1 | <1% | 22 | <1% |
| Black/African-American | 8.1 | 117 | 35% | 10.3 | 150 | 41% | 10.2 | 150 | 44% | 10.1 | 150 | 40% | 9.9 | 148 | 37% | 5,290 | 36% |
| Hispanic/Latinx | 2.4 | 10 | 3% | 4.1 | 18 | 5% | 2.4 | 11 | 3% | 5.1 | 24 | 6% | 3.1 | 15 | 4% | 495 | 3% |
| White | 2.1 | 193 | 58% | 1.9 | 174 | 48% | 1.8 | 169 | 49% | 1.9 | 176 | 47% | 2.3 | 208 | 52% | 8,592 | 59% |
| Multi-Race | 5.5 | 13 | 4% | 8.7 | 21 | 6% | 5.2 | 13 | 4% | 9.1 | 23 | 6% | 11.0 | 29 | 7% | 167 | 1% |
| Unknown | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 2 | <1% |
| Race/Ethnicity^b and Sex at birth | | | | | | | | | | | | | | | | | |
| American Indian/Alaska Native Males | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 9 | <1% |
| American Indian/Alaska Native Females | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 4 | <1% |
| Asian/Pacific Islander Males | * | - | - | * | - | - | * | 1 | <1% | * | - | - | * | 1 | <1% | 21 | <1% |
| Asian/Pacific Islander Females | * | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 1 | <1% |
| Black/African-American Males | 12.1 | 84 | 25% | 16.0 | 112 | 31% | 17.6 | 124 | 36% | 15.2 | 108 | 29% | 16.3 | 116 | 29% | 4,226 | 29% |
| Black/African-American Females | 4.4 | 33 | 10% | 5.0 | 38 | 10% | 3.4 | 26 | 8% | 5.4 | 42 | 11% | 4.1 | 32 | 8% | 1,064 | 7% |
| Hispanic/Latino Males | 3.7 | 8 | 2% | 6.6 | 15 | 4% | 4.2 | 10 | 3% | 7.5 | 18 | 5% | 4.8 | 12 | 3% | 387 | 3% |
| Hispanic/Latina Females | * | 2 | 1% | * | 3 | 1% | * | 1 | <1% | 2.6 | 6 | 2% | * | 3 | 1% | 108 | 1% |
| White Males | 3.7 | 167 | 50% | 3.3 | 149 | 41% | 3.2 | 143 | 42% | 3.3 | 150 | 40% | 3.9 | 175 | 44% | 7,853 | 54% |
| White Females | 0.6 | 26 | 8% | 0.5 | 25 | 7% | 0.6 | 26 | 8% | 0.6 | 26 | 7% | 0.7 | 33 | 8% | 739 | 5% |
| Multi-Race Males | 9.5 | 11 | 3% | 13.5 | 16 | 4% | 9.0 | 11 | 3% | 13.6 | 17 | 5% | 15.2 | 20 | 5% | 128 | 1% |
| Multi-Race Females | * | 2 | 1% | 4.1 | 5 | 1% | * | 2 | 1% | 4.7 | 6 | 2% | 6.7 | 9 | 2% | 39 | <1% |
| Unknown | - | - | - | * | - | - | * | - | - | * | - | - | * | - | - | 2 | <1% |
| Total | 2.9 | 333 | | 3.1 | 363 | | 2.9 | 344 | | 3.2 | 373 | | 3.4 | 401 | | 14,581 | |

Notes:
 Deaths among persons with HIV/AIDS represent deaths attributed to any cause among persons reported with a diagnosis of HIV and/or AIDS who were living in Ohio at the time of death. Deaths by year (2016-2020) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2020 among persons reported with a diagnosis of HIV and/or AIDS who were living in Ohio at the time of death.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of deaths among persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, Bureau of Vital Statistics; analysis conducted by HIV Surveillance Program. Data reported through June 30, 2021.

Trends in deaths among reported persons with diagnosed HIV infection by year of death and cumulative deaths by transmission category, Ohio, 2016-2020

| Transmission Category ^a | Year of Death among Persons with Diagnosed HIV Infection | | | | | | | | | | Cumulative Deaths | |
|----------------------------------------|----------------------------------------------------------|------|------------|------|------------|------|------------|------|------------|------|-------------------|------|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Male adult or adolescent | | | | | | | | | | | | |
| Male-to-male sexual contact | 146 | 54% | 176 | 60% | 156 | 54% | 171 | 59% | 194 | 60% | 8,060 | 64% |
| Injection drug use (IDU) | 20 | 7% | 21 | 7% | 28 | 10% | 25 | 9% | 29 | 9% | 1,128 | 9% |
| Male-to-male sexual contact and IDU | 25 | 9% | 23 | 8% | 26 | 9% | 25 | 9% | 19 | 6% | 1,013 | 8% |
| Heterosexual contact | 21 | 8% | 19 | 7% | 20 | 7% | 28 | 10% | 22 | 7% | 539 | 4% |
| Other/unknown | 58 | 21% | 52 | 18% | 59 | 20% | 43 | 15% | 60 | 19% | 1,823 | 15% |
| Subtotal | 270 | 100% | 291 | 100% | 289 | 100% | 292 | 100% | 324 | 100% | 12,563 | 100% |
| Female adult or adolescent | | | | | | | | | | | | |
| Injection drug use | 14 | 23% | 24 | 34% | 19 | 35% | 18 | 23% | 14 | 18% | 512 | 27% |
| Heterosexual contact | 41 | 66% | 37 | 53% | 31 | 56% | 55 | 70% | 57 | 74% | 1,192 | 63% |
| Other/unknown | 7 | 11% | 9 | 13% | 5 | 9% | 6 | 8% | 6 | 8% | 199 | 10% |
| Subtotal | 62 | 100% | 70 | 100% | 55 | 100% | 79 | 100% | 77 | 100% | 1,903 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | | | | | | | |
| Perinatal | 1 | 100% | 2 | 100% | - | - | 2 | 100% | - | - | 86 | 75% |
| Other/unknown | - | - | - | - | - | - | - | - | - | - | 29 | 25% |
| Subtotal | 1 | 100% | 2 | 100% | - | - | 2 | 100% | - | - | 115 | 100% |
| Total | 333 | | 363 | | 344 | | 373 | | 401 | | 14,581 | |

Trends in deaths among reported persons with diagnosed HIV infection by year of death and cumulative deaths by exposure category, Ohio, 2016-2020

| Exposure Category ^b | Year of Death among Persons with Diagnosed HIV Infection | | | | | | | | | | Cumulative Deaths | |
|----------------------------------------------------------|----------------------------------------------------------|-----|------------|-----|------------|-----|------------|-----|------------|-----|-------------------|-----|
| | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | No. | % |
| | No. | % | No. | % | No. | % | No. | % | No. | % | | |
| Male-to-male sexual contact only | 135 | 41% | 162 | 45% | 142 | 41% | 156 | 42% | 184 | 46% | 7,752 | 53% |
| Injection drug use (IDU) only | 15 | 5% | 25 | 7% | 21 | 6% | 21 | 6% | 20 | 5% | 1,010 | 7% |
| Heterosexual contact only | 62 | 19% | 56 | 15% | 51 | 15% | 84 | 23% | 79 | 20% | 1,734 | 12% |
| Male-to-male sexual contact & IDU | 19 | 6% | 17 | 5% | 18 | 5% | 16 | 4% | 16 | 4% | 829 | 6% |
| IDU & Heterosexual contact | 19 | 6% | 20 | 6% | 26 | 8% | 22 | 6% | 23 | 6% | 630 | 4% |
| Male-to-male sexual contact & Heterosexual contact | 11 | 3% | 14 | 4% | 14 | 4% | 15 | 4% | 10 | 2% | 308 | 2% |
| Male-to-male sexual contact & IDU & Heterosexual contact | 6 | 2% | 6 | 2% | 8 | 2% | 9 | 2% | 3 | 1% | 184 | 1% |
| Perinatal exposure | 1 | <1% | 3 | 1% | 1 | <1% | 2 | 1% | - | - | 88 | 1% |
| Other/unknown | 65 | 20% | 60 | 17% | 63 | 18% | 48 | 13% | 66 | 16% | 2,046 | 14% |
| Total | 333 | | 363 | | 344 | | 373 | | 401 | | 14,581 | |

Notes:

Deaths among persons with HIV/AIDS represent deaths attributed to any cause among persons reported with a diagnosis of HIV and/or AIDS who were living in Ohio at the time of death. Deaths by year (2016-2020) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2020 among persons reported with a diagnosis of HIV and/or AIDS who were living in Ohio at the time of death.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, Bureau of Vital Statistics; analysis conducted by HIV Surveillance Program. Data reported through June 30, 2021.

Trends in deaths among reported persons with diagnosed HIV infection by year of death (2016-2020) and cumulative deaths by county, Ohio

| County ^a | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | Cumulative Deaths No. | County ^a | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | | Cumulative Deaths No. |
|---------------------|-------------------|-----|-------------------|-----|-------------------|-----|-------------------|-----|-------------------|-----|--------------------------|---------------------|-------------------|------------|-------------------|------------|-------------------|------------|-------------------|------------|-------------------|------------|--------------------------|
| | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | | | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | Rate ^b | No. | |
| Adams | * | - | * | - | * | 1 | * | 1 | * | - | 10 | Logan | * | 1 | * | 2 | * | 1 | * | - | * | 1 | 27 |
| Allen | * | 1 | * | 4 | * | 3 | * | 1 | 6.9 | 7 | 97 | Lorain | * | 1 | 2.9 | 9 | 1.9 | 6 | * | 4 | 2.9 | 9 | 229 |
| Ashland | * | - | * | - | * | - | * | 2 | * | 1 | 15 | Lucas | 6.2 | 27 | 4.9 | 21 | 4.4 | 19 | 4.4 | 19 | 2.8 | 12 | 662 |
| Ashtabula | * | 1 | * | 2 | * | - | * | 4 | * | 1 | 65 | Madison | * | 3 | * | 1 | * | - | * | - | * | - | 19 |
| Athens | * | 1 | * | - | * | 1 | * | 2 | * | 1 | 32 | Mahoning | 3.5 | 8 | 6.1 | 14 | 2.6 | 6 | 6.1 | 14 | 5.3 | 12 | 324 |
| Auglaize | * | - | * | 2 | * | 3 | * | - | * | 1 | 22 | Marion | * | 1 | * | - | * | 2 | * | 1 | * | 1 | 53 |
| Belmont | * | 1 | * | 1 | * | - | * | - | * | 1 | 28 | Medina | * | 4 | * | 4 | * | 2 | * | 1 | * | - | 48 |
| Brown | * | - | * | 1 | * | 1 | * | 1 | * | 1 | 18 | Meigs | * | 1 | * | - | * | 1 | * | - | * | - | 9 |
| Butler | * | 3 | 2.9 | 11 | 2.1 | 8 | 2.9 | 11 | 1.8 | 7 | 203 | Mercer | * | - | * | - | * | - | * | - | * | - | 15 |
| Carroll | * | - | * | - | * | - | * | 2 | * | - | 14 | Miami | * | 2 | * | 1 | * | 1 | * | - | * | 2 | 55 |
| Champaign | * | 1 | * | - | * | - | * | - | * | - | 24 | Monroe | * | - | * | - | * | - | * | - | * | - | 7 |
| Clark | * | 4 | 5.2 | 7 | 3.7 | 5 | * | 4 | * | 4 | 163 | Montgomery | 4.7 | 25 | 4.1 | 22 | 5.8 | 31 | 5.3 | 28 | 2.6 | 14 | 876 |
| Clermont | * | 2 | * | 4 | * | 3 | * | 1 | * | 1 | 81 | Morgan | * | - | * | - | * | - | * | - | * | - | 4 |
| Clinton | * | - | * | 1 | * | 1 | * | - | * | 1 | 28 | Morrow | * | 1 | * | - | * | - | * | - | * | - | 19 |
| Columbiana | * | - | * | 1 | * | - | * | - | * | 1 | 44 | Muskingum | * | 2 | * | 3 | * | - | * | 2 | * | 2 | 55 |
| Coshocton | * | - | * | - | * | 1 | * | - | * | - | 8 | Noble | * | 1 | * | - | * | 1 | * | - | * | - | 3 |
| Crawford | * | - | * | - | * | 1 | * | 1 | * | - | 29 | Ottawa | * | 1 | * | - | * | - | * | 1 | * | - | 17 |
| Cuyahoga | 4.5 | 56 | 4.3 | 54 | 4.5 | 56 | 7.1 | 88 | 7.7 | 95 | 2,775 | Paulding | * | 2 | * | - | * | - | * | - | * | - | 9 |
| Darke | * | 2 | * | 1 | * | 2 | * | 2 | * | 2 | 30 | Perry | * | 2 | * | - | * | - | * | - | * | - | 15 |
| Defiance | * | 1 | * | 1 | * | - | * | - | * | 1 | 18 | Pickaway | * | 2 | * | 1 | * | - | * | - | * | 2 | 55 |
| Delaware | * | 4 | * | 2 | * | 1 | * | 1 | * | 2 | 55 | Pike | * | - | * | - | * | 2 | * | - | * | 1 | 13 |
| Erie | * | 1 | * | - | * | 2 | * | 4 | * | 3 | 76 | Portage | * | - | * | 1 | * | 2 | * | 1 | * | 3 | 65 |
| Fairfield | * | - | * | 1 | * | 3 | 3.2 | 5 | * | 2 | 73 | Preble | * | 1 | * | 1 | * | 1 | * | - | * | 1 | 21 |
| Fayette | * | - | * | - | * | - | * | - | * | 1 | 24 | Putnam | * | - | * | - | * | - | * | - | * | - | 6 |
| Franklin | 4.6 | 58 | 5.5 | 71 | 5.0 | 66 | 3.8 | 50 | 5.7 | 75 | 2,342 | Richland | * | - | * | 2 | * | - | * | 2 | * | 2 | 106 |
| Fulton | * | 1 | * | - | * | - | * | - | * | 1 | 20 | Ross | * | 1 | * | 1 | * | 1 | * | 1 | * | 2 | 63 |
| Gallia | * | - | * | - | * | - | * | 2 | * | - | 8 | Sandusky | * | 1 | * | 1 | * | - | * | 2 | * | - | 36 |
| Geauga | * | 1 | * | - | * | - | * | 1 | * | - | 31 | Scioto | * | - | * | 1 | * | 4 | * | 2 | * | 3 | 56 |
| Greene | * | 1 | * | 2 | * | 2 | 3.0 | 5 | 2.9 | 5 | 102 | Seneca | * | - | * | 1 | * | - | * | - | * | 1 | 27 |
| Guernsey | * | - | * | 2 | * | - | * | - | * | - | 19 | Shelby | * | 1 | * | - | * | - | * | 2 | * | - | 26 |
| Hamilton | 4.8 | 39 | 6.1 | 50 | 6.5 | 53 | 5.3 | 43 | 6.6 | 54 | 1,518 | Stark | 2.9 | 11 | * | 2 | 2.2 | 8 | 1.6 | 6 | 2.4 | 9 | 316 |
| Hancock | * | - | * | - | * | - | * | - | * | 1 | 33 | Summit | 4.3 | 23 | 4.6 | 25 | 3.0 | 16 | 3.9 | 21 | 2.4 | 13 | 655 |
| Hardin | * | - | * | - | * | - | * | - | * | - | 12 | Trumbull | 4.5 | 9 | * | 2 | * | 2 | 5.1 | 10 | 3.0 | 6 | 159 |
| Harrison | * | - | * | - | * | 1 | * | - | * | 2 | 7 | Tuscarawas | * | 4 | * | - | * | - | * | - | * | 2 | 25 |
| Henry | * | - | * | - | * | - | * | - | * | 1 | 9 | Union | * | - | * | 1 | * | 2 | * | 2 | * | - | 27 |
| Highland | * | - | * | - | * | - | * | - | * | 1 | 25 | Van Wert | * | - | * | 1 | * | - | * | - | * | - | 8 |
| Hocking | * | - | * | 1 | * | - | * | 2 | * | - | 18 | Vinton | * | - | * | - | * | - | * | - | * | - | 7 |
| Holmes | * | 1 | * | - | * | 1 | * | - | * | - | 6 | Warren | * | 2 | * | 1 | 2.2 | 5 | * | 2 | * | 4 | 57 |
| Huron | * | 1 | * | - | * | - | * | 2 | * | 2 | 22 | Washington | * | 1 | * | - | * | - | * | - | * | - | 41 |
| Jackson | * | 2 | * | - | * | - | * | - | * | 1 | 16 | Wayne | * | 1 | * | 3 | * | - | * | 2 | * | - | 33 |
| Jefferson | * | - | * | 1 | * | 1 | * | - | * | 1 | 57 | Williams | * | 1 | * | - | * | 2 | * | - | * | 2 | 21 |
| Knox | * | 2 | * | 2 | * | - | * | - | * | - | 24 | Wood | * | 1 | * | 3 | * | 1 | * | 2 | * | 1 | 50 |
| Lake | 2.2 | 5 | 2.2 | 5 | * | 2 | 3.0 | 7 | 2.2 | 5 | 107 | Wyandot | * | - | * | - | * | - | * | - | * | - | 10 |
| Lawrence | * | - | * | 1 | * | - | * | 2 | * | - | 26 | No County | * | 2 | * | 4 | * | 4 | * | 2 | * | 5 | 1,818 |
| Licking | * | 1 | 4.0 | 7 | 3.4 | 6 | * | 2 | 3.4 | 6 | 100 | Ohio | 2.9 | 333 | 3.1 | 363 | 2.9 | 344 | 3.2 | 373 | 3.4 | 401 | 14,581 |

Notes:
 Deaths among persons with HIV/AIDS represent deaths attributed to any cause among persons reported with a diagnosis of HIV and/or AIDS who were living in Ohio at the time of death. Deaths by year (2016-2020) represent the number of deaths per year; cumulative deaths represent all reported deaths since the beginning of the epidemic through 2020 among persons reported with a diagnosis of HIV and/or AIDS who were living in Ohio at the time of death.
 Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a County reflects county of residence at time of death. Persons whose county of residence at time of death is unknown are included in No County.
^b The rate is the number of deaths among persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
 Source: Ohio Department of Health, Bureau of Vital Statistics; analysis conducted by HIV Surveillance Program. Data reported through June 30, 2021.