



Department of  
Health

# New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 10

**Counties:** Allen, Auglaize, Champaign, Hancock, Hardin, Logan,  
Mercer, Paulding, Putnam, Shelby, Van Wert

\*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2025

## Reported new diagnoses of HIV infection in 2024 by disease status and selected characteristics, HIV Prevention Planning Region 10

Characteristic	2024 diagnosis of HIV infection			Disease Status					
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex									
Males	7.8	20	56%	17	59%	3	50%	-	-
Females	6.3	16	44%	12	41%	3	50%	1	100%
Age at diagnosis (years)									
0-9	*	-	-	-	-	-	-	-	-
10-14	*	-	-	-	-	-	-	-	-
15-19	*	1	3%	1	3%	-	-	-	-
20-24	*	1	3%	1	3%	-	-	-	-
25-29	*	4	11%	4	14%	-	-	-	-
30-34	*	4	11%	4	14%	-	-	-	-
35-39	36.9	11	31%	9	31%	2	33%	-	-
40-44	20.1	6	17%	4	14%	1	17%	1	100%
45-49	24.3	7	19%	5	17%	2	33%	-	-
50-54	*	1	3%	1	3%	-	-	-	-
55-64	*	1	3%	-	-	1	17%	-	-
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity <sup>b</sup>									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
Black/African American	104.3	19	53%	13	45%	5	83%	1	100%
Hispanic/Latino	*	1	3%	1	3%	-	-	-	-
White	3.5	16	44%	15	52%	1	17%	-	-
Multi-Race	*	-	-	-	-	-	-	-	-
Race/Ethnicity <sup>b</sup> and Sex									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African American Males	93.3	9	25%	6	21%	3	50%	-	-
Black/African American Females	116.8	10	28%	7	24%	2	33%	1	100%
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-
Hispanic/Latina Females	*	1	3%	1	3%	-	-	-	-
White Males	4.9	11	31%	11	38%	-	-	-	-
White Females	2.2	5	14%	4	14%	1	17%	-	-
Multi-Race Males	*	-	-	-	-	-	-	-	-
Multi-Race Females	*	-	-	-	-	-	-	-	-
Total	7.1	36		29		6		1	

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Reported new diagnoses of HIV infection in 2024 by disease status and transmission category, HIV Prevention Planning Region 10

Transmission Category <sup>a</sup>	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	12	60%	12	71%	-	-	-	-
Injection drug use (IDU)	1	5%	1	6%	-	-	-	-
Male-to-male sexual contact and IDU	2	10%	2	12%	-	-	-	-
Heterosexual contact	1	5%	1	6%	-	-	-	-
Other/unknown	4	20%	1	6%	3	100%	-	-
Subtotal	20	100%	17	100%	3	100%	-	-
<b>Female adult or adolescent</b>								
Injection drug use	2	13%	2	17%	-	-	-	-
Heterosexual contact	11	69%	8	67%	2	67%	1	100%
Other/unknown	3	19%	2	17%	1	33%	-	-
Subtotal	16	100%	12	100%	3	100%	1	100%
<b>Child (&lt;13 years at diagnosis)</b>								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
<b>Total</b>	<b>36</b>		<b>29</b>		<b>6</b>		<b>1</b>	

## Reported new diagnoses of HIV infection in 2024 by disease status and exposure category, HIV Prevention Planning Region 10

Exposure Category <sup>b</sup>	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	8	22%	8	28%	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	15	42%	10	34%	4	67%	1	100%
Male-to-male sexual contact & IDU	1	3%	1	3%	-	-	-	-
IDU & Heterosexual contact	3	8%	3	10%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	4	11%	4	14%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	1	3%	1	3%	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	4	11%	2	7%	2	33%	-	-
<b>Total</b>	<b>36</b>		<b>29</b>		<b>6</b>		<b>1</b>	

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 10

2024 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino <sup>a</sup>			White			Multi-Race		
	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
0-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	1	5%	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	1	5%	*	-	-	*	-	-	*	-	-
25-29	*	-	-	*	-	-	*	1	5%	*	-	-	*	3	19%	*	-	-
30-34	*	-	-	*	-	-	*	-	-	*	1	100%	*	3	19%	*	-	-
35-39	*	-	-	*	-	-	602.9	7	37%	*	-	-	*	4	25%	*	-	-
40-44	*	-	-	*	-	-	*	3	16%	*	-	-	*	3	19%	*	-	-
45-49	*	-	-	*	-	-	*	4	21%	*	-	-	*	3	19%	*	-	-
50-54	*	-	-	*	-	-	*	1	5%	*	-	-	*	-	-	*	-	-
55-64	*	-	-	*	-	-	*	1	5%	*	-	-	*	-	-	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	104.3	19	-	*	1	-	3.5	16	-	*	-	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 10

Transmission Category <sup>c</sup>	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	4	44%	-	-	8	73%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	9%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	2	18%	-	-
Heterosexual contact	-	-	-	-	1	11%	-	-	-	-	-	-
Other/unknown	-	-	-	-	4	44%	-	-	-	-	-	-
Subtotal	-	-	-	-	9	100%	-	-	11	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	2	40%	-	-
Heterosexual contact	-	-	-	-	7	70%	1	100%	3	60%	-	-
Other/unknown	-	-	-	-	3	30%	-	-	-	-	-	-
Subtotal	-	-	-	-	10	100%	1	100%	5	100%	-	-
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	19	-	1	-	16	-	-	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 10

Exposure Category <sup>d</sup>	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	3	16%	-	-	5	31%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	11	58%	1	100%	3	19%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	1	6%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	3	19%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	1	5%	-	-	3	19%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	6%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	4	21%	-	-	-	-	-	-
Total	-	-	-	-	19	-	1	-	16	-	-	-

Notes:  
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.  
Asterisk (\*) indicates rate not calculated, for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.  
<sup>b</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.  
<sup>c</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.  
<sup>d</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

**Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 10**

Diagnoses of HIV Infection by Year																	
Characteristic	2020			2021			2022			2023			2024			Cumulative Diagnoses	
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	No.	%
Sex																	
Males	3.5	9	75%	2.0	5	50%	3.5	9	75%	7.8	20	95%	7.8	20	56%	563	77%
Females	*	3	25%	2.0	5	50%	*	3	25%	*	1	5%	6.3	16	44%	170	23%
Age at diagnosis (years)																	
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	7	1%
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
15-19	*	-	-	*	-	-	*	1	8%	*	-	-	*	1	3%	19	3%
20-24	*	2	17%	*	1	10%	*	-	-	*	4	19%	*	1	3%	88	12%
25-29	*	1	8%	*	2	20%	*	4	33%	*	3	14%	*	4	11%	116	16%
30-34	*	3	25%	*	1	10%	*	2	17%	*	2	10%	*	4	11%	119	16%
35-39	*	1	8%	*	-	-	17.1	5	42%	*	3	14%	36.9	11	31%	111	15%
40-44	*	-	-	*	1	10%	*	-	-	*	2	10%	20.1	6	17%	99	14%
45-49	*	1	8%	*	2	20%	*	-	-	*	3	14%	24.3	7	19%	64	9%
50-54	*	2	17%	*	1	10%	*	-	-	*	2	10%	*	1	3%	53	7%
55-64	*	1	8%	*	1	10%	*	-	-	*	2	10%	*	1	3%	44	6%
65+	*	1	8%	*	1	10%	*	-	-	*	-	-	*	-	-	11	2%
Race/Ethnicity <sup>b</sup>																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Black/African American	*	1	8%	*	2	20%	*	2	17%	33.5	6	29%	104.3	19	53%	137	19%
Hispanic/Latino	*	-	-	*	-	-	*	2	17%	*	2	10%	*	1	3%	29	4%
White	2.4	11	92%	1.7	8	80%	1.8	8	67%	2.9	13	62%	3.5	16	44%	543	74%
Multi-Race	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	23	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Race/Ethnicity <sup>b</sup> and Sex																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African American Males	*	1	8%	*	-	-	*	2	17%	63.2	6	29%	93.3	9	25%	80	11%
Black/African American Females	*	-	-	*	2	20%	*	-	-	*	-	-	116.8	10	28%	57	8%
Hispanic/Latino Males	*	-	-	*	-	-	*	2	17%	*	2	10%	*	-	-	20	3%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%	9	1%
White Males	3.5	8	67%	2.2	5	50%	2.2	5	42%	5.3	12	57%	4.9	11	31%	445	61%
White Females	*	3	25%	*	3	30%	*	3	25%	*	1	5%	2.2	5	14%	98	13%
Multi-Race Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	17	2%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	6	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Total	2.4	12		2.0	10		2.4	12		4.1	21		7.1	36		733	

Notes:  
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 10

Transmission Category <sup>a</sup>	Diagnoses of HIV Infection by Year											
	2020		2021		2022		2023		2024		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	7	78%	4	80%	5	56%	11	55%	12	60%	364	65%
Injection drug use (IDU)	-	-	-	-	-	-	1	5%	1	5%	29	5%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	2	10%	2	10%	36	6%
Heterosexual contact	-	-	-	-	-	-	2	10%	1	5%	32	6%
Other/unknown	2	22%	1	20%	4	44%	4	20%	4	20%	102	18%
Subtotal	9	100%	5	100%	9	100%	20	100%	20	100%	563	100%
<b>Female adult or adolescent</b>												
Injection drug use	1	33%	-	-	-	-	-	-	2	13%	28	17%
Heterosexual contact	2	67%	4	80%	3	100%	1	100%	11	69%	122	74%
Other/unknown	-	-	1	20%	-	-	-	-	3	19%	14	9%
Subtotal	3	100%	5	100%	3	100%	1	100%	16	100%	164	100%
<b>Child (&lt;13 years at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	17%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
<b>Total</b>	<b>12</b>		<b>10</b>		<b>12</b>		<b>21</b>		<b>36</b>		<b>733</b>	

## Trends in reported diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 10

Exposure Category <sup>b</sup>	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	6	50%	3	30%	4	33%	8	38%	8	22%	243	33%
Injection drug use (IDU) only	-	-	-	-	-	-	1	5%	-	-	7	1%
Heterosexual contact only	3	25%	5	50%	4	33%	6	29%	15	42%	194	26%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	1	5%	1	3%	20	3%
IDU & Heterosexual contact	1	8%	-	-	-	-	-	-	3	8%	50	7%
Male-to-male sexual contact & Heterosexual contact	1	8%	1	10%	1	8%	3	14%	4	11%	121	17%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	1	5%	1	3%	16	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	1	8%	1	10%	3	25%	1	5%	4	11%	77	11%
Total	12		10		12		21		36		733	

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 10

Characteristic	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
<b>Sex</b>						
Males	6	55%	2	67%	4	50%
Females	5	45%	1	33%	4	50%
<b>Age at AIDS diagnosis (years)</b>						
0-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	1	9%	1	33%	-	-
35-39	3	27%	-	-	3	38%
40-44	2	18%	-	-	2	25%
45-49	4	36%	2	67%	2	25%
50-54	-	-	-	-	-	-
55-64	1	9%	-	-	1	13%
65+	-	-	-	-	-	-
<b>Race/Ethnicity<sup>b</sup></b>						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African American	6	55%	1	33%	5	63%
Hispanic/Latino	-	-	-	-	-	-
White	5	45%	2	67%	3	38%
Multi-Race	-	-	-	-	-	-
<b>Race/Ethnicity<sup>b</sup> and Sex</b>						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African American Males	4	36%	1	33%	3	38%
Black/African American Females	2	18%	-	-	2	25%
Hispanic/Latino Males	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-
White Males	2	18%	1	33%	1	13%
White Females	3	27%	1	33%	2	25%
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
<b>Total</b>	<b>11</b>		<b>3</b>		<b>8</b>	

**Notes:**

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

<sup>a</sup> Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 10

Transmission Category <sup>b</sup>	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	1	17%	-	-	1	25%
Injection drug use (IDU)	1	17%	1	50%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Other/unknown	4	67%	1	50%	3	75%
Subtotal	6	100%	2	100%	4	100%
<b>Female adult or adolescent</b>						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	5	100%	1	100%	4	100%
Other/unknown	-	-	-	-	-	-
Subtotal	5	100%	1	100%	4	100%
<b>Child (&lt;13 years at diagnosis)</b>						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
<b>Total</b>	<b>11</b>		<b>3</b>		<b>8</b>	

## Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 10

Exposure Category <sup>c</sup>	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-
Injection drug use (IDU) only	1	9%	1	33%	-	-
Heterosexual contact only	8	73%	2	67%	6	75%
Male-to-male sexual contact & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	1	9%	-	-	1	13%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	1	9%	-	-	1	13%
<b>Total</b>	<b>11</b>		<b>3</b>		<b>8</b>	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

<sup>b</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>c</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.



# Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 10

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	3	100%	8	89%	7	88%	5	100%	6	55%	386	82%
Females	-	-	1	11%	1	13%	-	-	5	45%	86	18%
Age at AIDS diagnosis (years)												
0-9	-	-	-	-	-	-	-	-	-	-	6	1%
10-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-	4	1%
20-24	-	-	-	-	-	-	1	20%	-	-	18	4%
25-29	-	-	-	-	1	13%	-	-	-	-	66	14%
30-34	1	33%	1	11%	2	25%	-	-	1	9%	74	16%
35-39	-	-	2	22%	1	13%	1	20%	3	27%	71	15%
40-44	-	-	-	-	1	13%	1	20%	2	18%	65	14%
45-49	-	-	1	11%	1	13%	1	20%	4	36%	66	14%
50-54	-	-	-	-	1	13%	1	20%	-	-	44	9%
55-64	1	33%	3	33%	1	13%	-	-	1	9%	46	10%
65+	1	33%	2	22%	-	-	-	-	-	-	12	3%
Race/Ethnicity <sup>a</sup>												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
Black/African American	-	-	1	11%	2	25%	1	20%	6	55%	78	17%
Hispanic/Latino	1	33%	-	-	1	13%	1	20%	-	-	17	4%
White	2	67%	8	89%	5	63%	3	60%	5	45%	361	76%
Multi-Race	-	-	-	-	-	-	-	-	-	-	16	3%
Race/Ethnicity <sup>a</sup> and Sex												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
Black/African American Males	-	-	1	11%	1	13%	1	20%	4	36%	48	10%
Black/African American Females	-	-	-	-	1	13%	-	-	2	18%	30	6%
Hispanic/Latino Males	1	33%	-	-	1	13%	1	20%	-	-	12	3%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	5	1%
White Males	2	67%	7	78%	5	63%	3	60%	2	18%	313	66%
White Females	-	-	1	11%	-	-	-	-	3	27%	48	10%
Multi-Race Males	-	-	-	-	-	-	-	-	-	-	13	3%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	3	1%
Total	3		9		8		5		11		472	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 10

Transmission Category <sup>a</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	2	67%	5	63%	2	29%	2	40%	1	17%	252	66%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	17%	15	4%
Male-to-male sexual contact and IDU	-	-	1	13%	-	-	-	-	-	-	19	5%
Heterosexual contact	-	-	1	13%	1	14%	1	20%	-	-	25	7%
Other/unknown	1	33%	1	13%	4	57%	2	40%	4	67%	73	19%
Subtotal	3	100%	8	100%	7	100%	5	100%	6	100%	384	100%
<b>Female adult or adolescent</b>												
Injection drug use	-	-	-	-	-	-	-	-	-	-	15	18%
Heterosexual contact	-	-	1	100%	1	100%	-	-	5	100%	63	77%
Other/unknown	-	-	-	-	-	-	-	-	-	-	4	5%
Subtotal	-	-	1	100%	1	100%	-	-	5	100%	82	100%
<b>Child (&lt;13 years at diagnosis)</b>												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	17%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
<b>Total</b>	<b>3</b>		<b>9</b>		<b>8</b>		<b>5</b>		<b>11</b>		<b>472</b>	

## Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 10

Exposure Category <sup>b</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	33%	4	44%	2	25%	2	40%	-	-	168	36%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	1	9%	5	1%
Heterosexual contact only	-	-	3	33%	3	38%	3	60%	8	73%	118	25%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	9	2%
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	25	5%
Male-to-male sexual contact & Heterosexual contact	1	33%	1	11%	-	-	-	-	1	9%	84	18%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	11%	-	-	-	-	-	-	10	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	1	33%	-	-	3	38%	-	-	1	9%	48	10%
Total	3		9		8		5		11		472	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

# Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 10

---

**Counties:** Allen, Auglaize, Champaign, Hancock, Hardin, Logan,  
Mercer, Paulding, Putnam, Shelby, Van Wert

## Reported persons living with diagnosed HIV infection in 2024 by current disease status and selected characteristics, HIV Prevention Planning Region 10

Characteristic	Living with diagnosed HIV infection in 2024			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	137.9	352	74%	176	70%	176	79%
Females	47.6	121	26%	75	30%	46	21%
Age at end of year (years)							
0-9	*	-	-	-	-	-	-
10-14	*	2	<1%	2	1%	-	-
15-19	*	1	<1%	-	-	1	<1%
20-24	15.8	5	1%	5	2%	-	-
25-29	70.5	21	4%	18	7%	3	1%
30-34	114.0	36	8%	31	12%	5	2%
35-39	198.1	59	12%	36	14%	23	10%
40-44	137.0	41	9%	29	12%	12	5%
45-49	201.0	58	12%	33	13%	25	11%
50-54	178.4	54	11%	28	11%	26	12%
55-64	194.5	128	27%	46	18%	82	37%
65+	67.4	68	14%	23	9%	45	20%
Race/Ethnicity <sup>b</sup>							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	3	1%	2	1%	1	<1%
Black/African American	631.3	115	24%	69	27%	46	21%
Hispanic/Latino	171.9	34	7%	20	8%	14	6%
White	66.1	300	63%	150	60%	150	68%
Multi-Race	195.5	21	4%	10	4%	11	5%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity <sup>b</sup> and Sex							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	2	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	*	1	<1%	1	<1%	-	-
Black/African American Males	704.6	68	14%	40	16%	28	13%
Black/African American Females	548.7	47	10%	29	12%	18	8%
Hispanic/Latino Males	242.2	25	5%	13	5%	12	5%
Hispanic/Latina Females	95.2	9	2%	7	3%	2	1%
White Males	106.0	240	51%	114	45%	126	57%
White Females	26.4	60	13%	36	14%	24	11%
Multi-Race Males	315.3	17	4%	8	3%	9	4%
Multi-Race Females	*	4	1%	2	1%	2	1%
Unknown	*	-	-	-	-	-	-
Total	92.8	473		251		222	

### Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Reported persons living with diagnosed HIV infection in 2024 by current disease status and transmission category, HIV Prevention Planning Region 10

Transmission Category <sup>a</sup>	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	232	66%	118	67%	114	65%
Injection drug use (IDU)	12	3%	7	4%	5	3%
Male-to-male sexual contact and IDU	20	6%	9	5%	11	6%
Heterosexual contact	28	8%	9	5%	19	11%
Other/unknown	58	17%	32	18%	26	15%
Subtotal	350	100%	175	100%	175	100%
<b>Female adult or adolescent</b>						
Injection drug use	14	12%	9	12%	5	11%
Heterosexual contact	94	80%	56	77%	38	84%
Other/unknown	10	8%	8	11%	2	4%
Subtotal	118	100%	73	100%	45	100%
<b>Child (&lt;13 years at diagnosis)</b>						
Perinatal	4	80%	3	100%	1	50%
Other/unknown	1	20%	-	-	1	50%
Subtotal	5	100%	3	100%	2	100%
<b>Total</b>	<b>473</b>		<b>251</b>		<b>222</b>	

## Reported persons living with diagnosed HIV infection in 2024 by current disease status and exposure category, HIV Prevention Planning Region 10

Exposure Category <sup>b</sup>	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	165	35%	81	32%	84	38%
Injection drug use (IDU) only	1	<1%	1	<1%	-	-
Heterosexual contact only	148	31%	78	31%	70	32%
Male-to-male sexual contact & IDU	8	2%	4	2%	4	2%
IDU & Heterosexual contact	25	5%	15	6%	10	5%
Male-to-male sexual contact & Heterosexual contact	67	14%	37	15%	30	14%
Male-to-male sexual contact & IDU & Heterosexual contact	12	3%	5	2%	7	3%
Perinatal exposure	5	1%	4	2%	1	<1%
Other/unknown	42	9%	26	10%	16	7%
<b>Total</b>	<b>473</b>		<b>251</b>		<b>222</b>	

### Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and age, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2024																				
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino <sup>a</sup>			White			Multi-Race			Unknown	
	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
10-14	*	-	-	*	-	-	*	1	1%	*	1	3%	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	*	4	3%	*	1	3%	*	-	-	*	-	-	-	-
25-29	*	-	-	*	-	-	*	4	3%	*	4	12%	43.1	11	4%	*	2	10%	-	-
30-34	*	-	-	*	-	-	1,005.7	14	12%	*	2	6%	65.5	18	6%	*	2	10%	-	-
35-39	*	-	-	*	-	-	1,808.8	21	18%	387.0	5	15%	117.2	31	10%	*	2	10%	-	-
40-44	*	-	-	*	-	-	869.6	10	9%	*	1	3%	108.7	29	10%	*	1	5%	-	-
45-49	*	-	-	*	-	-	1,410.9	16	14%	650.6	7	21%	127.5	33	11%	*	2	10%	-	-
50-54	*	-	-	*	2	67%	883.2	9	8%	*	4	12%	119.2	33	11%	2,678.6	6	29%	-	-
55-64	*	-	-	*	-	-	1,475.7	30	26%	531.1	7	21%	142.2	87	29%	*	4	19%	-	-
65+	*	-	-	*	1	33%	204.2	5	4%	*	2	6%	60.8	58	19%	*	2	10%	-	-
Total	*	-	-	*	3		631.3	115		171.9	34		66.1	300		195.5	21		-	-

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2024														
Transmission Category <sup>c</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/ Latino <sup>a</sup>		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	39	59%	11	44%	169	70%	13	76%	-	-
Injection drug use (IDU)	-	-	-	-	3	5%	1	4%	7	3%	1	6%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	2	3%	2	8%	15	6%	1	6%	-	-
Heterosexual contact	-	-	1	50%	7	11%	7	28%	12	5%	1	6%	-	-
Other/unknown	-	-	1	50%	15	23%	4	16%	37	15%	1	6%	-	-
Subtotal	-	-	2	100%	66	100%	25	100%	240	100%	17	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	2	4%	1	13%	11	19%	-	-	-	-
Heterosexual contact	-	-	1	100%	38	81%	6	75%	45	78%	4	100%	-	-
Other/unknown	-	-	-	-	7	15%	1	13%	2	3%	-	-	-	-
Subtotal	-	-	1	100%	47	100%	8	100%	58	100%	4	100%	-	-
Child (<13 years at diagnosis)														
Perinatal	-	-	-	-	1	50%	1	100%	2	100%	-	-	-	-
Other/unknown	-	-	-	-	1	50%	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	2	100%	1	100%	2	100%	-	-	-	-
Total	-		3		115		34		300		21		-	-

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2024														
Exposure Category <sup>d</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	27	23%	7	21%	123	41%	8	38%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	1	0%	-	-	-	-
Heterosexual contact only	-	-	2	67%	54	47%	15	44%	71	24%	6	29%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	1	1%	1	3%	6	2%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	5	4%	2	6%	17	6%	1	5%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	12	10%	4	12%	46	15%	5	24%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	1	1%	1	3%	9	3%	1	5%	-	-
Perinatal exposure	-	-	-	-	1	1%	2	6%	2	1%	-	-	-	-
Other/unknown	-	-	1	33%	14	12%	2	6%	25	8%	-	-	-	-
Total	-	-	3	-	115	-	34	-	300	-	21	-	-	-

Notes:  
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.  
Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.  
<sup>b</sup> The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.  
<sup>c</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.  
<sup>d</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 10, 2020-2024

Living with Diagnosed HIV Infection															
Characteristic	2020			2021			2022			2023			2024		
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%
Sex															
Males	124.7	317	77%	120.9	310	76%	123.0	314	76%	122.8	313	75%	137.9	352	74%
Females	38.1	97	23%	38.6	98	24%	38.2	97	24%	40.6	103	25%	47.6	121	26%
Age at end of year (years)															
0-9	*	1	<1%	*	1	<1%	*	2	<1%	*	-	-	*	-	-
10-14	*	2	<1%	*	2	<1%	*	2	<1%	*	2	<1%	*	2	<1%
15-19	*	2	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
20-24	18.9	6	1%	*	4	1%	*	3	1%	19.0	6	1%	15.8	5	1%
25-29	53.5	17	4%	65.5	20	5%	60.5	18	4%	64.7	19	5%	70.5	21	4%
30-34	139.8	41	10%	119.3	36	9%	132.6	41	10%	125.0	39	9%	114.0	36	8%
35-39	125.8	37	9%	114.3	34	8%	130.0	38	9%	129.1	38	9%	198.1	59	12%
40-44	151.9	44	11%	147.1	44	11%	120.2	36	9%	126.4	38	9%	137.0	41	9%
45-49	155.4	46	11%	155.5	45	11%	161.7	46	11%	161.4	46	11%	201.0	58	12%
50-54	202.5	63	15%	180.4	57	14%	192.0	60	15%	194.7	60	14%	178.4	54	11%
55-64	162.8	115	28%	173.2	122	30%	166.1	114	28%	161.4	109	26%	194.5	128	27%
65+	42.3	40	10%	44.8	42	10%	51.9	50	12%	59.0	58	14%	67.4	68	14%
Race/Ethnicity <sup>b</sup>															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	3	1%	*	3	1%	*	3	1%	*	3	1%	*	3	1%
Black/African American	471.3	85	21%	458.6	83	20%	468.2	83	20%	468.9	84	20%	631.3	115	24%
Hispanic/Latino	176.1	30	7%	163.9	29	7%	155.3	28	7%	164.6	30	7%	171.9	34	7%
White	60.0	275	66%	59.0	270	66%	60.9	278	68%	61.3	279	67%	66.1	300	63%
Multi-Race	221.7	21	5%	234.2	23	6%	188.4	19	5%	190.5	20	5%	195.5	21	4%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity <sup>b</sup> and Sex															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	2	<1%	*	2	<1%	*	2	<1%	*	2	<1%	*	2	<1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Black/African American Males	536.3	52	13%	488.3	48	12%	529.0	50	12%	526.4	50	12%	704.6	68	14%
Black/African American Females	395.7	33	8%	423.4	35	9%	398.7	33	8%	403.9	34	8%	548.7	47	10%
Hispanic/Latino Males	255.9	23	6%	236.3	22	5%	221.8	21	5%	231.3	22	5%	242.2	25	5%
Hispanic/Latina Females	87.0	7	2%	83.5	7	2%	81.8	7	2%	91.8	8	2%	95.2	9	2%
White Males	97.9	223	54%	95.6	219	54%	99.0	226	55%	98.5	224	54%	106.0	240	51%
White Females	22.6	52	13%	22.3	51	13%	22.8	52	13%	24.1	55	13%	26.4	60	13%
Multi-Race Males	358.2	17	4%	387.8	19	5%	298.2	15	4%	284.8	15	4%	315.3	17	4%
Multi-Race Females	*	4	1%	*	4	1%	*	4	1%	95.6	5	1%	*	4	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	81.4	414		80.0	408		80.8	411		81.8	416		92.8	473	

Notes:  
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.  
<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

**Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 10, 2020-2024**

Living with Diagnosed HIV Infection										
Transmission Category <sup>a</sup>	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	205	65%	205	67%	207	66%	209	67%	232	66%
Injection drug use (IDU)	10	3%	9	3%	9	3%	11	4%	12	3%
Male-to-male sexual contact and IDU	26	8%	22	7%	21	7%	17	5%	20	6%
Heterosexual contact	29	9%	25	8%	23	7%	22	7%	28	8%
Other/unknown	45	14%	47	15%	52	17%	52	17%	58	17%
Subtotal	315	100%	308	100%	312	100%	311	100%	350	100%
<b>Female adult or adolescent</b>										
Injection drug use	13	14%	13	14%	10	11%	14	14%	14	12%
Heterosexual contact	76	81%	78	81%	79	84%	80	79%	94	80%
Other/unknown	5	5%	5	5%	5	5%	7	7%	10	8%
Subtotal	94	100%	96	100%	94	100%	101	100%	118	100%
<b>Child (&lt;13 years at diagnosis)</b>										
Perinatal	4	80%	3	75%	4	80%	3	75%	4	80%
Other/unknown	1	20%	1	25%	1	20%	1	25%	1	20%
Subtotal	5	100%	4	100%	5	100%	4	100%	5	100%
<b>Total</b>	<b>414</b>		<b>408</b>		<b>411</b>		<b>416</b>		<b>473</b>	

**Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 10, 2020-2024**

Living with Diagnosed HIV Infection										
Exposure Category <sup>b</sup>	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	145	35%	145	36%	143	35%	146	35%	165	35%
Injection drug use (IDU) only	1	<1%	1	<1%	1	<1%	1	<1%	1	<1%
Heterosexual contact only	124	30%	122	30%	123	30%	125	30%	148	31%
Male-to-male sexual contact & IDU	11	3%	9	2%	8	2%	5	1%	8	2%
IDU & Heterosexual contact	22	5%	21	5%	18	4%	24	6%	25	5%
Male-to-male sexual contact & Heterosexual contact	60	14%	60	15%	64	16%	63	15%	67	14%
Male-to-male sexual contact & IDU & Heterosexual contact	15	4%	13	3%	13	3%	12	3%	12	3%
Perinatal exposure	5	1%	4	1%	5	1%	4	1%	5	1%
Other/unknown	31	7%	33	8%	36	9%	36	9%	42	9%
<b>Total</b>	<b>414</b>		<b>408</b>		<b>411</b>		<b>416</b>		<b>473</b>	

Notes:  
 Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.  
 Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.  
<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.