

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 10

Counties: Allen, Auglaize, Champaign, Hancock, Hardin, Logan, Mercer, Paulding, Putnam, Shelby, Van Wert

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2021

Reported new diagnoses of HIV infection in 2020 by disease status and selected characteristics, HIV Prevention Planning Region 10

Characteristic	2020 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	3.9	10	77%	8	73%	1	100%	1	100%
Females	*	3	23%	3	27%	-	-	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-	-	-
20-24	*	2	15%	2	18%	-	-	-	-
25-29	*	1	8%	1	9%	-	-	-	-
30-34	*	3	23%	3	27%	-	-	-	-
35-39	*	1	8%	1	9%	-	-	-	-
40-44	*	-	-	-	-	-	-	-	-
45-49	*	1	8%	1	9%	-	-	-	-
50-54	*	3	23%	3	27%	-	-	-	-
55-64	*	1	8%	-	-	-	-	1	100%
65+	*	1	8%	-	-	1	100%	-	-
Race/Ethnicity^b									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
Black/African-American	*	1	8%	1	9%	-	-	-	-
Hispanic/Latinx	*	-	-	-	-	-	-	-	-
White	2.6	12	92%	10	91%	1	100%	1	100%
Multi-Race	*	-	-	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African-American Males	*	1	8%	1	9%	-	-	-	-
Black/African-American Females	*	-	-	-	-	-	-	-	-
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
White Males	4.0	9	69%	7	64%	1	100%	1	100%
White Females	*	3	23%	3	27%	-	-	-	-
Multi-Race Males	*	-	-	-	-	-	-	-	-
Multi-Race Females	*	-	-	-	-	-	-	-	-
Total	2.6	13		11		1		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by disease status and transmission category, HIV Prevention Planning Region 10

Transmission Category ^a	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	8	80%	7	88%	-	-	1	100%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	2	20%	1	13%	1	100%	-	-
Subtotal	10	100%	8	100%	1	100%	1	100%
Female adult or adolescent								
Injection drug use	1	33%	1	33%	-	-	-	-
Heterosexual contact	2	67%	2	67%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	3	100%	3	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	13		11		1		1	

Reported new diagnoses of HIV infection in 2020 by disease status and exposure category, HIV Prevention Planning Region 10

Exposure Category ^b	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	8	62%	7	64%	-	-	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	2	15%	2	18%	-	-	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	8%	1	9%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	2	15%	1	9%	1	100%	-	-
Total	13		11		1		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 10

2020 diagnosis of HIV infection

Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	-	-	*	-	-	*	2	17%	*	-	-
25-29	*	-	-	*	-	-	*	-	-	*	-	-	*	1	8%	*	-	-
30-34	*	-	-	*	-	-	*	-	-	*	-	-	*	3	25%	*	-	-
35-39	*	-	-	*	-	-	*	-	-	*	-	-	*	1	8%	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
45-49	*	-	-	*	-	-	*	1	100%	*	-	-	*	-	-	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	3	25%	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	1	8%	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	1	8%	*	-	-
Total	*	-	-	*	-	-	*	1	-	*	-	-	2.6	12	-	*	-	-

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and transmission category, HIV Prevention Planning Region 10

2020 diagnosis of HIV infection

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	1	100%	-	-	7	78%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	22%	-	-
Subtotal	-	-	-	-	1	100%	-	-	9	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	1	33%	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	2	67%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	3	100%	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	1	-	-	-	12	-	-	-

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and exposure category, HIV Prevention Planning Region 10

2020 diagnosis of HIV infection

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	1	100%	-	-	7	58%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	-	-	-	-	2	17%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	8%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	17%	-	-
Total	-	-	-	-	1	-	-	-	12	-	-	-

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 10

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2016			2017			2018			2019			2020			No.	%
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%		
Sex at birth																	
Males	3.5	9	53%	4.3	11	69%	5.1	13	68%	6.3	16	80%	3.9	10	77%	513	78%
Females	3.1	8	47%	2.0	5	31%	2.3	6	32%	*	4	20%	*	3	23%	148	22%
Age at diagnosis (yr)																	
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	7	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	17	3%
20-24	*	1	6%	*	2	13%	*	2	11%	15.7	5	25%	*	2	15%	81	12%
25-29	16.5	5	29%	*	1	6%	15.6	5	26%	*	2	10%	*	1	8%	103	16%
30-34	*	3	18%	*	2	13%	*	4	21%	*	2	10%	*	3	23%	111	17%
35-39	*	2	12%	*	1	6%	*	-	-	*	4	20%	*	1	8%	93	14%
40-44	*	-	-	*	3	19%	*	3	16%	*	1	5%	*	-	-	92	14%
45-49	*	1	6%	*	2	13%	*	1	5%	*	-	-	*	1	8%	53	8%
50-54	*	2	12%	14.6	5	31%	*	-	-	*	3	15%	*	3	23%	51	8%
55-64	*	3	18%	*	-	-	*	3	16%	*	3	15%	*	1	8%	41	6%
65+	*	-	-	*	-	-	*	1	5%	*	-	-	*	1	8%	10	2%
Race/Ethnicity^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Black/African-American	*	3	18%	*	-	-	33.4	6	32%	*	4	20%	*	1	8%	110	17%
Hispanic/Latinx	*	1	6%	*	4	25%	*	-	-	*	-	-	*	-	-	24	4%
White	2.8	13	76%	2.4	11	69%	2.8	13	68%	3.5	16	80%	2.6	12	92%	505	76%
Multi-Race	*	-	-	*	1	6%	*	-	-	*	-	-	*	-	-	20	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Race/Ethnicity^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African-American Males	*	1	6%	*	-	-	*	4	21%	*	1	5%	*	1	8%	64	10%
Black/African-American Females	*	2	12%	*	-	-	*	2	11%	*	3	15%	*	-	-	46	7%
Hispanic/Latino Males	*	1	6%	*	3	19%	*	-	-	*	-	-	*	-	-	15	2%
Hispanic/Latina Females	*	-	-	*	1	6%	*	-	-	*	-	-	*	-	-	9	1%
White Males	3.0	7	41%	3.0	7	44%	3.9	9	47%	6.5	15	75%	4.0	9	69%	417	63%
White Females	2.6	6	35%	*	4	25%	*	4	21%	*	1	5%	*	3	23%	88	13%
Multi-Race Males	*	-	-	*	1	6%	*	-	-	*	-	-	*	-	-	15	2%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	5	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Total	3.3	17		3.1	16		3.7	19		3.9	20		2.6	13		661	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 10

Transmission Category ^a	Diagnoses of HIV Infection by Year											
	2016		2017		2018		2019		2020		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	5	56%	10	91%	6	46%	7	44%	8	80%	334	65%
Injection drug use (IDU)	1	11%	1	9%	1	8%	1	6%	-	-	28	5%
Male-to-male sexual contact and IDU	1	11%	-	-	1	8%	5	31%	-	-	34	7%
Heterosexual contact	2	22%	-	-	2	15%	1	6%	-	-	29	6%
Other/unknown	-	-	-	-	3	23%	2	13%	2	20%	88	17%
Subtotal	9	100%	11	100%	13	100%	16	100%	10	100%	513	100%
Female adult or adolescent												
Injection drug use	4	50%	2	40%	1	17%	-	-	1	33%	27	19%
Heterosexual contact	4	50%	2	40%	4	67%	4	100%	2	67%	105	74%
Other/unknown	-	-	1	20%	1	17%	-	-	-	-	10	7%
Subtotal	8	100%	5	100%	6	100%	4	100%	3	100%	142	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	17%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	17		16		19		20		13		661	

Trends in reported diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 10

Exposure Category ^b	Diagnoses of HIV Infection by Year											
	2016		2017		2018		2019		2020		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	5	29%	9	56%	6	32%	7	35%	8	62%	308	47%
Injection drug use (IDU) only	-	-	2	13%	-	-	1	5%	-	-	24	4%
Heterosexual contact only	6	35%	2	13%	6	32%	5	25%	2	15%	135	20%
Male-to-male sexual contact & IDU	1	6%	-	-	1	5%	3	15%	-	-	26	4%
IDU & Heterosexual contact	5	29%	1	6%	2	11%	-	-	1	8%	31	5%
Male-to-male sexual contact & Heterosexual contact	-	-	1	6%	-	-	-	-	-	-	26	4%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	2	10%	-	-	8	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	-	-	1	6%	4	21%	2	10%	2	15%	98	15%
Total	17		16		19		20		13		661	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020. Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 10

Characteristic	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex at birth						
Males	3	100%	1	100%	2	100%
Females	-	-	-	-	-	-
Age at AIDS diagnosis (yr)						
<13	-	-	-	-	-	-
13-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	1	33%	1	100%	-	-
35-39	-	-	-	-	-	-
40-44	-	-	-	-	-	-
45-49	-	-	-	-	-	-
50-54	-	-	-	-	-	-
55-64	1	33%	-	-	1	50%
65+	1	33%	-	-	1	50%
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African-American	-	-	-	-	-	-
Hispanic/Latinx	-	-	-	-	-	-
White	3	100%	1	100%	2	100%
Multi-Race	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African-American Males	-	-	-	-	-	-
Black/African-American Females	-	-	-	-	-	-
Hispanic/Latino Males	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-
White Males	3	100%	1	100%	2	100%
White Females	-	-	-	-	-	-
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
Total	3		1		2	

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 10

Transmission Category ^b	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2	67%	1	100%	1	50%
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Other/unknown	1	33%	-	-	1	50%
Subtotal	3	100%	1	100%	2	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	3		1		2	

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 10

Exposure Category ^c	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	33%	-	-	1	50%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	1	33%	1	100%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	1	33%	-	-	1	50%
Total	3		1		2	

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 10

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex at birth												
Males	11	92%	10	91%	5	63%	7	70%	3	100%	348	83%
Females	1	8%	1	9%	3	38%	3	30%	-	-	73	17%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	5	1%
13-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-	4	1%
20-24	-	-	-	-	1	13%	1	10%	-	-	15	4%
25-29	2	17%	-	-	1	13%	-	-	-	-	61	14%
30-34	1	8%	1	9%	2	25%	1	10%	1	33%	68	16%
35-39	-	-	1	9%	1	13%	-	-	-	-	60	14%
40-44	-	-	2	18%	-	-	-	-	-	-	57	14%
45-49	3	25%	2	18%	-	-	-	-	-	-	57	14%
50-54	2	17%	4	36%	1	13%	3	30%	-	-	44	10%
55-64	4	33%	1	9%	1	13%	5	50%	1	33%	41	10%
65+	-	-	-	-	1	13%	-	-	1	33%	9	2%
Race/Ethnicity^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
Black/African-American	-	-	1	9%	2	25%	2	20%	-	-	60	14%
Hispanic/Latinx	1	8%	1	9%	-	-	-	-	-	-	11	3%
White	11	92%	9	82%	6	75%	8	80%	3	100%	336	80%
Multi-Race	-	-	-	-	-	-	-	-	-	-	13	3%
Race/Ethnicity^a and Sex at birth												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	1	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
Black/African-American Males	-	-	1	9%	1	13%	-	-	-	-	36	9%
Black/African-American Females	-	-	-	-	1	13%	2	20%	-	-	24	6%
Hispanic/Latino Males	1	8%	1	9%	-	-	-	-	-	-	7	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	4	1%
White Males	10	83%	8	73%	4	50%	7	70%	3	100%	294	70%
White Females	1	8%	1	9%	2	25%	1	10%	-	-	42	10%
Multi-Race Males	-	-	-	-	-	-	-	-	-	-	10	2%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	3	1%
Total	12		11		8		10		3		421	

Notes:

AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 10

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	7	64%	8	80%	3	60%	5	71%	2	67%	234	67%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	15	4%
Male-to-male sexual contact and IDU	1	9%	-	-	-	-	-	-	-	-	17	5%
Heterosexual contact	1	9%	1	10%	1	20%	-	-	-	-	20	6%
Other/unknown	2	18%	1	10%	1	20%	2	29%	1	33%	62	18%
Subtotal	11	100%	10	100%	5	100%	7	100%	3	100%	348	100%
Female adult or adolescent												
Injection drug use	1	100%	1	100%	-	-	1	33%	-	-	10	14%
Heterosexual contact	-	-	-	-	2	67%	2	67%	-	-	55	80%
Other/unknown	-	-	-	-	1	33%	-	-	-	-	4	6%
Subtotal	1	100%	1	100%	3	100%	3	100%	-	-	69	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	3	75%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	25%
Subtotal	-	-	-	-	-	-	-	-	-	-	4	100%
Total	12		11		8		10		3		421	

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 10

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sexual contact only	7	58%	7	64%	3	38%	5	50%	1	33%	215	51%
Injection drug use (IDU) only	-	-	1	9%	-	-	-	-	-	-	12	3%
Heterosexual contact only	1	8%	1	9%	3	38%	2	20%	-	-	76	18%
Male-to-male sexual contact & IDU	1	8%	-	-	-	-	-	-	-	-	14	3%
IDU & Heterosexual contact	1	8%	-	-	-	-	1	10%	-	-	13	3%
Male-to-male sexual contact & Heterosexual contact	-	-	1	9%	-	-	-	-	1	33%	19	5%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	3	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	3	1%
Other/unknown	2	17%	1	9%	2	25%	2	20%	1	33%	66	16%
Total	12		11		8		10		3		421	

Notes:

AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode or transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 10

Counties: Allen, Auglaize, Champaign, Hancock, Hardin, Logan, Mercer, Paulding, Putnam, Shelby, Van Wert

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, HIV Prevention Planning Region 10

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	123.9	315	76%	150	72%	165	79%
Females	39.7	101	24%	57	28%	44	21%
Age at end of year							
<13	*	3	1%	2	1%	1	<1%
13-14	*	-	-	-	-	-	-
15-19	*	1	<1%	1	<1%	-	-
20-24	15.8	5	1%	4	2%	1	<1%
25-29	56.6	18	4%	16	8%	2	1%
30-34	146.6	43	10%	28	14%	15	7%
35-39	132.6	39	9%	27	13%	12	6%
40-44	148.4	43	10%	23	11%	20	10%
45-49	148.6	44	11%	24	12%	20	10%
50-54	205.7	64	15%	22	11%	42	20%
55-64	161.4	114	27%	45	22%	69	33%
65+	44.5	42	10%	15	7%	27	13%
Race/Ethnicity^b							
American Indian/Alaska Native	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander	*	3	1%	1	<1%	2	1%
Black/African-American	493.5	89	21%	44	21%	45	22%
Hispanic/Latinx	187.8	32	8%	19	9%	13	6%
White	59.8	274	66%	134	65%	140	67%
Multi-Race	179.4	17	4%	9	4%	8	4%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	2	<1%	1	<1%	1	<1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
Black/African-American Males	556.9	54	13%	27	13%	27	13%
Black/African-American Females	419.7	35	8%	17	8%	18	9%
Hispanic/Latino Males	278.1	25	6%	14	7%	11	5%
Hispanic/Latina Females	87.0	7	2%	5	2%	2	1%
White Males	96.6	220	53%	101	49%	119	57%
White Females	23.4	54	13%	33	16%	21	10%
Multi-Race Males	273.9	13	3%	7	3%	6	3%
Multi-Race Females	*	4	1%	2	1%	2	1%
Unknown	*	-	-	-	-	-	-
Total	81.7	416		207		209	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, HIV Prevention Planning Region 10

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	204	65%	95	64%	109	67%
Injection drug use (IDU)	13	4%	7	5%	6	4%
Male-to-male sexual contact and IDU	23	7%	12	8%	11	7%
Heterosexual contact	29	9%	12	8%	17	10%
Other/unknown	43	14%	23	15%	20	12%
Subtotal	312	100%	149	100%	163	100%
Female adult or adolescent						
Injection drug use	15	15%	10	18%	5	12%
Heterosexual contact	77	79%	40	73%	37	86%
Other/unknown	6	6%	5	9%	1	2%
Subtotal	98	100%	55	100%	43	100%
Child (<13 yrs at diagnosis)						
Perinatal	5	83%	3	100%	2	67%
Other/unknown	1	17%	-	-	1	33%
Subtotal	6	100%	3	100%	3	100%
Total	416		207		209	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, HIV Prevention Planning Region 10

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	184	44%	87	42%	97	46%
Injection drug use (IDU) only	9	2%	5	2%	4	2%
Heterosexual contact only	107	26%	52	25%	55	26%
Male-to-male sexual contact & IDU	18	4%	11	5%	7	3%
IDU & Heterosexual contact	19	5%	12	6%	7	3%
Male-to-male sexual contact & Heterosexual contact	20	5%	8	4%	12	6%
Male-to-male sexual contact & IDU & Heterosexual contact	5	1%	1	<1%	4	2%
Perinatal exposure	6	1%	4	2%	2	1%
Other/unknown	48	12%	27	13%	21	10%
Total	416		207		209	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	-	-	*	2	2%	*	1	3%	*	-	-	*	-	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	1	3%	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	*	2	2%	*	-	-	*	3	1%	*	-	-	-	-
25-29	*	-	-	*	-	-	478.8	7	8%	*	1	3%	32.3	9	3%	*	1	6%	-	-
30-34	*	-	-	*	-	-	988.5	12	13%	396.8	5	16%	96.2	25	9%	*	1	6%	-	-
35-39	*	-	-	*	-	-	656.8	8	9%	*	2	6%	102.9	27	10%	*	2	12%	-	-
40-44	*	-	-	*	-	-	896.1	10	11%	557.6	6	19%	95.9	25	9%	*	2	12%	-	-
45-49	*	-	-	*	1	33%	692.4	7	8%	552.5	5	16%	103.7	28	10%	*	3	18%	-	-
50-54	*	-	-	*	-	-	1,591.8	17	19%	848.7	6	19%	139.4	40	15%	*	1	6%	-	-
55-64	*	-	-	*	1	33%	1,028.5	22	25%	*	4	13%	124.2	82	30%	1,262.6	5	29%	-	-
65+	*	1	100%	*	1	33%	*	2	2%	*	1	3%	38.9	35	13%	*	2	12%	-	-
Total	*	1		*	3		493.5	89		187.8	32		59.8	274		179.4	17			

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	1	100%	-	-	-	-	33	63%	9	38%	151	69%	10	77%
Injection drug use (IDU)	-	-	-	-	-	-	4	8%	2	8%	7	3%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	4	8%	3	13%	16	7%	-	-
Heterosexual contact	-	-	-	50%	1	50%	6	12%	8	33%	13	6%	1	8%
Other/unknown	-	-	-	50%	1	50%	5	10%	2	8%	33	15%	2	15%
Subtotal	1	100%	2	100%	52	100%	24	100%	220	100%	13	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	-	-	4	11%	-	-	10	19%	1	25%
Heterosexual contact	-	-	-	100%	1	100%	29	83%	5	83%	39	75%	3	75%
Other/unknown	-	-	-	-	-	-	2	6%	1	17%	3	6%	-	-
Subtotal	-	-	1	100%	35	100%	6	100%	52	100%	4	100%	-	-
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	-	-	1	50%	2	100%	2	100%	-	-
Other/unknown	-	-	-	-	-	-	1	50%	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	2	100%	2	100%	2	100%	-	-
Total	1		3		89		32		274		17			

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	100%	-	-	-	-	30	34%	9	28%	136	50%	8	47%
Injection drug use (IDU) only	-	-	-	-	-	-	4	4%	-	-	5	2%	-	-
Heterosexual contact only	-	-	-	67%	2	67%	35	39%	13	41%	53	19%	4	24%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	4	4%	3	9%	11	4%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	4	4%	2	6%	12	4%	1	6%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	3	3%	-	-	15	5%	2	12%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	5	2%	-	-	
Perinatal exposure	-	-	-	-	-	-	1	1%	3	9%	2	1%	-	-
Other/unknown	-	-	1	33%	8	9%	2	6%	35	13%	2	12%	-	-
Total	1		3		89		32		274		17			

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 10, 2016-2020

Living with Diagnosed HIV Infection

Characteristic	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	108.6	277	74%	113.0	288	74%	115.3	294	74%	118.4	302	75%	123.9	315	76%
Females	38.2	98	26%	40.3	103	26%	39.9	102	26%	39.6	101	25%	39.7	101	24%
Age at end of year															
<13	*	1	<1%	*	1	<1%	*	2	1%	*	3	1%	*	3	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	1	<1%	*	-	-	*	-	-	*	-	-	*	1	<1%
20-24	23.9	8	2%	30.6	10	3%	21.8	7	2%	21.9	7	2%	15.8	5	1%
25-29	95.8	29	8%	82.7	26	7%	71.9	23	6%	53.1	17	4%	56.6	18	4%
30-34	102.6	30	8%	108.0	31	8%	114.6	33	8%	120.5	35	9%	146.6	43	10%
35-39	142.2	42	11%	120.2	36	9%	123.8	37	9%	114.6	34	8%	132.6	39	9%
40-44	121.0	35	9%	161.3	46	12%	161.6	46	12%	181.6	52	13%	148.4	43	10%
45-49	165.1	52	14%	166.2	52	13%	175.5	54	14%	165.2	50	12%	148.6	44	11%
50-54	185.5	66	18%	195.5	67	17%	164.2	54	14%	176.8	56	14%	205.7	64	15%
55-64	126.8	91	24%	136.0	98	25%	152.7	110	28%	160.6	115	29%	161.4	114	27%
65+	23.2	20	5%	27.3	24	6%	33.3	30	8%	36.8	34	8%	44.5	42	10%
Race/Ethnicity^b															
American Indian/Alaska Native	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander	*	3	1%	*	2	1%	*	4	1%	*	4	1%	*	3	1%
Black/African-American	426.1	76	20%	424.1	76	19%	433.6	78	20%	474.9	85	21%	493.5	89	21%
Hispanic/Latinx	133.4	20	5%	163.8	25	6%	157.7	25	6%	175.0	29	7%	187.8	32	8%
White	56.0	260	69%	58.5	271	69%	58.7	271	68%	58.4	269	67%	59.8	274	66%
Multi-Race	173.5	15	4%	180.8	16	4%	188.2	17	4%	163.2	15	4%	179.4	17	4%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	2	1%	*	1	<1%	*	3	1%	*	3	1%	*	2	<1%
Asian/Pacific Islander Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Black/African-American Males	472.5	45	12%	470.2	45	12%	436.5	42	11%	498.0	48	12%	556.9	54	13%
Black/African-American Females	372.8	31	8%	371.3	31	8%	430.3	36	9%	447.8	37	9%	419.7	35	8%
Hispanic/Latino Males	177.4	14	4%	198.8	16	4%	201.2	17	4%	238.7	21	5%	278.1	25	6%
Hispanic/Latina Females	84.5	6	2%	124.8	9	2%	108.0	8	2%	102.9	8	2%	87.0	7	2%
White Males	88.5	204	54%	92.6	213	54%	95.0	218	55%	94.3	216	54%	96.6	220	53%
White Females	23.9	56	15%	24.9	58	15%	22.8	53	13%	22.9	53	13%	23.4	54	13%
Multi-Race Males	254.7	11	3%	270.0	12	3%	287.3	13	3%	283.5	13	3%	273.9	13	3%
Multi-Race Females	*	4	1%	*	4	1%	*	4	1%	*	2	<1%	*	4	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	73.3	375		76.5	391		77.6	396		79.0	403		81.7	416	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 10, 2016-2020

Living with Diagnosed HIV Infection										
Transmission Category ^a	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	187	68%	198	69%	193	66%	191	64%	204	65%
Injection drug use (IDU)	15	5%	15	5%	16	5%	13	4%	13	4%
Male-to-male sexual contact and IDU	13	5%	11	4%	18	6%	21	7%	23	7%
Heterosexual contact	21	8%	24	8%	24	8%	31	10%	29	9%
Other/unknown	40	14%	39	14%	41	14%	43	14%	43	14%
Subtotal	276	100%	287	100%	292	100%	299	100%	312	100%
Female adult or adolescent										
Injection drug use	17	18%	19	19%	17	17%	13	13%	15	15%
Heterosexual contact	75	78%	76	75%	76	76%	79	80%	77	79%
Other/unknown	4	4%	6	6%	7	7%	7	7%	6	6%
Subtotal	96	100%	101	100%	100	100%	99	100%	98	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	100%	3	100%	4	100%	4	80%	5	83%
Other/unknown	-	-	-	-	-	-	1	20%	1	17%
Subtotal	3	100%	3	100%	4	100%	5	100%	6	100%
Total	375		391		396		403		416	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 10, 2016-2020

Living with Diagnosed HIV Infection										
Exposure Category ^b	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	168	45%	180	46%	173	44%	169	42%	184	44%
Injection drug use (IDU) only	7	2%	10	3%	10	3%	10	2%	9	2%
Heterosexual contact only	97	26%	101	26%	101	26%	111	28%	107	26%
Male-to-male sexual contact & IDU	9	2%	7	2%	13	3%	17	4%	18	4%
IDU & Heterosexual contact	25	7%	24	6%	23	6%	16	4%	19	5%
Male-to-male sexual contact & Heterosexual contact	19	5%	18	5%	20	5%	22	5%	20	5%
Male-to-male sexual contact & IDU & Heterosexual contact	4	1%	4	1%	5	1%	4	1%	5	1%
Perinatal exposure	3	1%	3	1%	4	1%	4	1%	6	1%
Other/unknown	43	11%	44	11%	47	12%	50	12%	48	12%
Total	375		391		396		403		416	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.