



Department of
Health

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 6

Counties: Athens, Belmont, Guernsey, Meigs, Monroe, Morgan,
Muskingum, Noble, Perry, Washington

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2025

Reported new diagnoses of HIV infection in 2024 by disease status and selected characteristics, HIV Prevention Planning Region 6

Characteristic	2024 diagnosis of HIV infection			Disease Status						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	2.4	5	50%	2	33%	2	67%	1	100%	
Females	2.5	5	50%	4	67%	1	33%	-	-	
Age at diagnosis (years)										
0-9	*	-	-	-	-	-	-	-	-	
10-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	*	1	10%	1	17%	-	-	-	-	
25-29	*	1	10%	1	17%	-	-	-	-	
30-34	*	-	-	-	-	-	-	-	-	
35-39	*	3	30%	2	33%	1	33%	-	-	
40-44	*	-	-	-	-	-	-	-	-	
45-49	*	2	20%	2	33%	-	-	-	-	
50-54	*	2	20%	-	-	2	67%	-	-	
55-64	*	-	-	-	-	-	-	-	-	
65+	*	1	10%	-	-	-	-	1	100%	
Race/Ethnicity ^b										
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
Black/African American	*	2	20%	1	17%	1	33%	-	-	
Hispanic/Latino	*	-	-	-	-	-	-	-	-	
White	2.1	8	80%	5	83%	2	67%	1	100%	
Multi-Race	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b and Sex										
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
Black/African American Males	*	1	10%	-	-	1	33%	-	-	
Black/African American Females	*	1	10%	1	17%	-	-	-	-	
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
White Males	*	4	40%	2	33%	1	33%	1	100%	
White Females	*	4	40%	3	50%	1	33%	-	-	
Multi-Race Males	*	-	-	-	-	-	-	-	-	
Multi-Race Females	*	-	-	-	-	-	-	-	-	
Total	2.4	10		6		3		1		

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by disease status and transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	3	60%	1	50%	1	50%	1	100%
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	2	40%	1	50%	1	50%	-	-
Subtotal	5	100%	2	100%	2	100%	1	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	5	100%	4	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	5	100%	4	100%	1	100%	-	-
Child (<13 years at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	10		6		3		1	

Reported new diagnoses of HIV infection in 2024 by disease status and exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	6	60%	5	83%	1	33%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	3	30%	1	17%	1	33%	1	100%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	1	10%	-	-	1	33%	-	-
Total	10		6		3		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 6

2024 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
0-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	-	-	*	-	-	*	1	13%	*	-	-
25-29	*	-	-	*	-	-	*	-	-	*	-	-	*	1	13%	*	-	-
30-34	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
35-39	*	-	-	*	-	-	*	2	100%	*	-	-	*	1	13%	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	2	25%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	2	25%	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	1	13%	*	-	-
Total	*	-	-	*	-	-	*	2	-	*	-	-	2.1	8	-	*	-	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 6

Transmission Category ^c	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	-	-	-	-	3	75%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	-	-	1	25%	-	-
Subtotal	-	-	-	-	1	100%	-	-	4	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	1	100%	-	-	4	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	4	100%	-	-
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	2	-	-	-	8	-	-	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 6

Exposure Category ^d	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	1	50%	-	-	5	63%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	3	38%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	50%	-	-	-	-	-	-
Total	-	-	-	-	2	-	-	-	8	-	-	-

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
Asterisk (*) indicates rate not calculated. for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 6

Diagnoses of HIV Infection by Year																	
Characteristic	2020			2021			2022			2023			2024			Cumulative Diagnoses	
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex																	
Males	5.3	11	85%	5.3	11	92%	6.8	14	82%	4.4	9	64%	2.4	5	50%	432	80%
Females	*	2	15%	*	1	8%	*	3	18%	2.5	5	36%	2.5	5	50%	107	20%
Age at diagnosis (years)																	
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	5	1%
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	1%
15-19	*	-	-	*	-	-	*	1	6%	*	-	-	*	-	-	8	1%
20-24	*	1	8%	*	1	8%	*	2	12%	*	1	7%	*	1	10%	58	11%
25-29	*	3	23%	*	-	-	25.2	6	35%	*	4	29%	*	1	10%	86	16%
30-34	*	2	15%	*	2	17%	*	1	6%	*	2	14%	*	-	-	86	16%
35-39	*	2	15%	21.3	5	42%	*	4	24%	*	-	-	*	3	30%	98	18%
40-44	*	2	15%	*	-	-	*	2	12%	*	2	14%	*	-	-	74	14%
45-49	*	1	8%	*	-	-	*	1	6%	*	-	-	*	2	20%	52	10%
50-54	*	1	8%	*	1	8%	*	-	-	*	1	7%	*	2	20%	28	5%
55-64	*	1	8%	*	2	17%	*	-	-	*	3	21%	*	-	-	30	6%
65+	*	-	-	*	1	8%	*	-	-	*	1	7%	*	1	10%	11	2%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	1	8%	*	-	-	*	-	-	*	-	-	1	<1%
Black/African American	*	2	15%	*	-	-	*	4	24%	*	1	7%	*	2	20%	44	8%
Hispanic/Latino	*	-	-	*	1	8%	*	-	-	*	-	-	*	-	-	15	3%
White	2.8	11	85%	2.6	10	83%	3.4	13	76%	3.4	13	93%	2.1	8	80%	465	86%
Multi-Race	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	10	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	1%
Race/Ethnicity ^b and Sex																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	1	8%	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African American Males	*	2	15%	*	-	-	*	4	24%	*	-	-	*	1	10%	27	5%
Black/African American Females	*	-	-	*	-	-	*	-	-	*	1	7%	*	1	10%	17	3%
Hispanic/Latino Males	*	-	-	*	1	8%	*	-	-	*	-	-	*	-	-	12	2%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	1%
White Males	4.7	9	69%	4.7	9	75%	5.3	10	59%	4.8	9	64%	*	4	40%	382	71%
White Females	*	2	15%	*	1	8%	*	3	18%	*	4	29%	*	4	40%	83	15%
Multi-Race Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	8	1%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	1%
Total	3.1	13		2.9	12		4.2	17		3.4	14		2.4	10		539	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	9	82%	6	55%	11	79%	4	44%	3	60%	266	62%
Injection drug use (IDU)	1	9%	1	9%	1	7%	1	11%	-	-	19	4%
Male-to-male sexual contact and IDU	-	-	1	9%	-	-	1	11%	-	-	23	5%
Heterosexual contact	1	9%	-	-	-	-	-	-	-	-	24	6%
Other/unknown	-	-	3	27%	2	14%	3	33%	2	40%	95	22%
Subtotal	11	100%	11	100%	14	100%	9	100%	5	100%	427	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	1	33%	1	20%	-	-	17	16%
Heterosexual contact	-	-	1	100%	2	67%	4	80%	5	100%	74	70%
Other/unknown	2	100%	-	-	-	-	-	-	-	-	15	14%
Subtotal	2	100%	1	100%	3	100%	5	100%	5	100%	106	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	67%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	33%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	13		12		17		14		10		539	

Trends in reported diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 6

Diagnoses of HIV Infection by Year												
Exposure Category ^b	2020		2021		2022		2023		2024		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	23%	3	25%	6	35%	4	29%	-	-	184	34%
Injection drug use (IDU) only	-	-	-	-	2	12%	-	-	-	-	8	1%
Heterosexual contact only	1	8%	2	17%	3	18%	4	29%	6	60%	129	24%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	13	2%
IDU & Heterosexual contact	1	8%	1	8%	-	-	2	14%	-	-	28	5%
Male-to-male sexual contact & Heterosexual contact	6	46%	3	25%	5	29%	-	-	3	30%	82	15%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	8%	-	-	1	7%	-	-	10	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	2	15%	2	17%	1	6%	3	21%	1	10%	80	15%
Total	13		12		17		14		10		539	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 6

Characteristic	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex						
Males	3	75%	-	-	3	75%
Females	1	25%	-	-	1	25%
Age at AIDS diagnosis (years)						
0-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	-	-	-	-	-	-
35-39	1	25%	-	-	1	25%
40-44	-	-	-	-	-	-
45-49	-	-	-	-	-	-
50-54	2	50%	-	-	2	50%
55-64	-	-	-	-	-	-
65+	1	25%	-	-	1	25%
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African American	1	25%	-	-	1	25%
Hispanic/Latino	-	-	-	-	-	-
White	3	75%	-	-	3	75%
Multi-Race	-	-	-	-	-	-
Race/Ethnicity^b and Sex						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African American Males	1	25%	-	-	1	25%
Black/African American Females	-	-	-	-	-	-
Hispanic/Latino Males	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-
White Males	2	50%	-	-	2	50%
White Females	1	25%	-	-	1	25%
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
Total	4		-		4	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 6

Transmission Category ^b	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2	67%	-	-	2	67%
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Other/unknown	1	33%	-	-	1	33%
Subtotal	3	100%	-	-	3	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	1	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-
Subtotal	1	100%	-	-	1	100%
Child (<13 years at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	4		-		4	

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 6

Exposure Category ^c	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	1	25%	-	-	1	25%
Male-to-male sexual contact & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	2	50%	-	-	2	50%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	1	25%	-	-	1	25%
Total	4		-		4	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 6

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	2	67%	5	71%	4	80%	1	50%	3	75%	286	86%
Females	1	33%	2	29%	1	20%	1	50%	1	25%	48	14%
Age at AIDS diagnosis (years)												
0-9	-	-	-	-	-	-	-	-	-	-	2	1%
10-14	-	-	-	-	-	-	-	-	-	-	2	1%
15-19	-	-	-	-	-	-	-	-	-	-	2	1%
20-24	-	-	-	-	-	-	-	-	-	-	5	1%
25-29	-	-	-	-	-	-	-	-	-	-	34	10%
30-34	1	33%	-	-	1	20%	-	-	-	-	50	15%
35-39	1	33%	4	57%	2	40%	-	-	1	25%	67	20%
40-44	-	-	-	-	1	20%	-	-	-	-	68	20%
45-49	1	33%	-	-	1	20%	1	50%	-	-	42	13%
50-54	-	-	1	14%	-	-	-	-	2	50%	28	8%
55-64	-	-	1	14%	-	-	1	50%	-	-	26	8%
65+	-	-	1	14%	-	-	-	-	1	25%	8	2%
Race/Ethnicity ^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander	-	-	1	14%	-	-	-	-	-	-	1	<1%
Black/African American	-	-	1	14%	-	-	-	-	1	25%	24	7%
Hispanic/Latino	-	-	1	14%	1	20%	-	-	-	-	9	3%
White	3	100%	4	57%	4	80%	2	100%	3	75%	292	87%
Multi-Race	-	-	-	-	-	-	-	-	-	-	8	2%
Race/Ethnicity ^a and Sex												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	1	14%	-	-	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
Black/African American Males	-	-	1	14%	-	-	-	-	1	25%	19	6%
Black/African American Females	-	-	-	-	-	-	-	-	-	-	5	1%
Hispanic/Latino Males	-	-	1	14%	1	20%	-	-	-	-	7	2%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	2	1%
White Males	2	67%	2	29%	3	60%	1	50%	2	50%	252	75%
White Females	1	33%	2	29%	1	20%	1	50%	1	25%	40	12%
Multi-Race Males	-	-	-	-	-	-	-	-	-	-	7	2%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	1	<1%
Total	3		7		5		2		4		334	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	1	50%	3	60%	1	25%	1	100%	2	67%	171	61%
Injection drug use (IDU)	-	-	1	20%	2	50%	-	-	-	-	16	6%
Male-to-male sexual contact and IDU	-	-	1	20%	-	-	-	-	-	-	19	7%
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	18	6%
Other/unknown	1	50%	-	-	1	25%	-	-	1	33%	58	21%
Subtotal	2	100%	5	100%	4	100%	1	100%	3	100%	282	100%
Female adult or adolescent												
Injection drug use	1	100%	-	-	-	-	-	-	-	-	9	19%
Heterosexual contact	-	-	2	100%	1	100%	1	100%	1	100%	35	73%
Other/unknown	-	-	-	-	-	-	-	-	-	-	4	8%
Subtotal	1	100%	2	100%	1	100%	1	100%	1	100%	48	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	2	50%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	50%
Subtotal	-	-	-	-	-	-	-	-	-	-	4	100%
Total	3		7		5		2		4		334	

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	33%	1	14%	1	20%	1	50%	-	-	121	36%
Injection drug use (IDU) only	-	-	-	-	2	40%	-	-	-	-	7	2%
Heterosexual contact only	-	-	2	29%	2	40%	1	50%	1	25%	77	23%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	12	4%
IDU & Heterosexual contact	1	33%	1	14%	-	-	-	-	-	-	18	5%
Male-to-male sexual contact & Heterosexual contact	-	-	2	29%	-	-	-	-	2	50%	50	15%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	14%	-	-	-	-	-	-	7	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	3	1%
Other/unknown	1	33%	-	-	-	-	-	-	1	25%	39	12%
Total	3		7		5		2		4		334	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.



Department of
Health

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 6

Counties: Athens, Belmont, Guernsey, Meigs, Monroe, Morgan,
Muskingum, Noble, Perry, Washington

As of June 30, 2025

Reported persons living with diagnosed HIV infection in 2024 by current disease status and selected characteristics, HIV Prevention Planning Region 6

Characteristic	Living with diagnosed HIV infection in 2024			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	140.2	289	78%	154	76%	135	81%
Females	39.4	80	22%	48	24%	32	19%
Age at end of year (years)							
0-9	*	2	1%	2	1%	-	-
10-14	*	1	<1%	1	<1%	-	-
15-19	*	2	1%	1	<1%	1	1%
20-24	*	4	1%	3	1%	1	1%
25-29	59.3	14	4%	12	6%	2	1%
30-34	104.1	26	7%	19	9%	7	4%
35-39	123.8	29	8%	18	9%	11	7%
40-44	177.1	43	12%	30	15%	13	8%
45-49	117.9	27	7%	17	8%	10	6%
50-54	203.4	51	14%	30	15%	21	13%
55-64	181.3	99	27%	42	21%	57	34%
65+	82.9	71	19%	27	13%	44	26%
Race/Ethnicity ^b							
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Asian/Pacific Islander	*	4	1%	3	1%	1	1%
Black/African American	272.6	30	8%	14	7%	16	10%
Hispanic/Latino	190.9	12	3%	7	3%	5	3%
White	81.1	306	83%	170	84%	136	81%
Multi-Race	125.4	12	3%	3	1%	9	5%
Unknown	*	4	1%	4	2%	-	-
Race/Ethnicity ^b and Sex							
American Indian/Alaska Native Males	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	4	1%	3	1%	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African American Males	314.0	22	6%	8	4%	14	8%
Black/African American Females	200.2	8	2%	6	3%	2	1%
Hispanic/Latino Males	296.2	10	3%	6	3%	4	2%
Hispanic/Latina Females	*	2	1%	1	<1%	1	1%
White Males	127.3	240	65%	132	65%	108	65%
White Females	35.0	66	18%	38	19%	28	17%
Multi-Race Males	205.1	10	3%	2	1%	8	5%
Multi-Race Females	*	2	1%	1	<1%	1	1%
Unknown	*	4	1%	4	2%	-	-
Total	90.2	369		202		167	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by current disease status and transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	196	69%	104	68%	92	70%
Injection drug use (IDU)	9	3%	6	4%	3	2%
Male-to-male sexual contact and IDU	23	8%	15	10%	8	6%
Heterosexual contact	12	4%	2	1%	10	8%
Other/unknown	44	15%	25	16%	19	14%
Subtotal	284	100%	152	100%	132	100%
Female adult or adolescent						
Injection drug use	10	13%	5	11%	5	16%
Heterosexual contact	62	78%	35	74%	27	84%
Other/unknown	7	9%	7	15%	-	-
Subtotal	79	100%	47	100%	32	100%
Child (<13 years at diagnosis)						
Perinatal	5	83%	3	100%	2	67%
Other/unknown	1	17%	-	-	1	33%
Subtotal	6	100%	3	100%	3	100%
Total	369		202		167	

Reported persons living with diagnosed HIV infection in 2024 by current disease status and exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	148	40%	83	41%	65	39%
Injection drug use (IDU) only	5	1%	3	1%	2	1%
Heterosexual contact only	90	24%	44	22%	46	28%
Male-to-male sexual contact & IDU	15	4%	10	5%	5	3%
IDU & Heterosexual contact	14	4%	8	4%	6	4%
Male-to-male sexual contact & Heterosexual contact	48	13%	21	10%	27	16%
Male-to-male sexual contact & IDU & Heterosexual contact	8	2%	5	2%	3	2%
Perinatal exposure	5	1%	3	1%	2	1%
Other/unknown	36	10%	25	12%	11	7%
Total	369		202		167	

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Dash (-) indicates no cases were reported for the given category.
^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and age, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2024																				
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
0-9	*	-	-	*	2	50%	*	-	-	*	-	-	*	-	-	*	-	-	-	-
10-14	*	-	-	*	-	-	*	1	3%	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	1	3%	*	1	8%	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	*	-	-	*	1	8%	*	3	1%	*	-	-	-	-
25-29	*	-	-	*	-	-	*	4	13%	*	1	8%	43.2	9	3%	*	-	-	-	-
30-34	*	-	-	*	-	-	*	1	3%	*	1	8%	103.8	23	8%	*	1	8%	-	-
35-39	*	-	-	*	-	-	*	1	3%	*	2	17%	115.6	25	8%	*	1	8%	-	-
40-44	*	-	-	*	2	50%	741.8	5	17%	*	2	17%	146.2	33	11%	*	1	8%	-	-
45-49	*	1	100%	*	-	-	*	-	-	*	-	-	121.1	26	8%	*	-	-	-	-
50-54	*	-	-	*	-	-	*	4	13%	*	2	17%	177.6	42	14%	*	2	17%	1	25%
55-64	*	-	-	*	-	-	673.4	8	27%	*	-	-	155.7	81	26%	1,394.4	7	58%	3	75%
65+	*	-	-	*	-	-	340.6	5	17%	*	2	17%	77.9	64	21%	*	-	-	-	-
Total	*	1		*	4		272.6	30		190.9	12		81.1	306		125.4	12		4	

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2024														
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/ Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	10	48%	6	60%	171	72%	9	90%	-	-
Injection drug use (IDU)	-	-	2	100%	-	-	1	10%	6	3%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	5%	1	10%	21	9%	-	-	-	-
Heterosexual contact	-	-	-	-	3	14%	1	10%	8	3%	-	-	-	-
Other/unknown	1	100%	-	-	7	33%	1	10%	32	13%	1	10%	2	100%
Subtotal	1	100%	2	100%	21	100%	10	100%	238	100%	10	100%	2	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	-	-	-	-	9	14%	1	50%	-	-
Heterosexual contact	-	-	-	-	7	88%	1	100%	53	80%	1	50%	-	-
Other/unknown	-	-	-	-	1	13%	-	-	4	6%	-	-	2	100%
Subtotal	-	-	-	-	8	100%	1	100%	66	100%	2	100%	2	100%
Child (<13 years at diagnosis)														
Perinatal	-	-	2	100%	1	100%	1	100%	1	50%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	50%	-	-	-	-
Subtotal	-	-	2	100%	1	100%	1	100%	2	100%	-	-	-	-
Total	1		4		30		12		306		12		4	

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2024														
Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	5	17%	4	33%	133	43%	6	50%	-	-
Injection drug use (IDU) only	-	-	1	25%	-	-	-	-	4	1%	-	-	-	-
Heterosexual contact only	1	100%	-	-	13	43%	2	17%	73	24%	1	8%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	1	3%	1	8%	13	4%	-	-	-	-
IDU & Heterosexual contact	-	-	1	25%	-	-	1	8%	11	4%	1	8%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	5	17%	2	17%	38	12%	3	25%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	8	3%	-	-	-	-
Perinatal exposure	-	-	2	50%	1	3%	1	8%	1	0%	-	-	-	-
Other/unknown	-	-	-	-	5	17%	1	8%	25	8%	1	8%	4	100%
Total	1		4		30		12		306		12		4	

Notes:
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 6, 2020-2024

Living with Diagnosed HIV Infection															
Characteristic	2020			2021			2022			2023			2024		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	127.6	266	80%	135.5	280	80%	132.1	271	79%	128.2	264	78%	140.2	289	78%
Females	32.6	68	20%	34.3	70	20%	34.8	70	21%	36.4	74	22%	39.4	80	22%
Age at end of year (years)															
0-9	*	2	1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
10-14	*	2	1%	*	1	<1%	*	1	<1%	*	-	-	*	1	<1%
15-19	*	-	-	*	1	<1%	*	1	<1%	*	2	1%	*	2	1%
20-24	15.8	5	1%	*	4	1%	16.2	5	1%	*	4	1%	*	4	1%
25-29	70.6	18	5%	45.6	11	3%	54.5	13	4%	59.4	14	4%	59.3	14	4%
30-34	112.8	27	8%	137.7	33	9%	90.7	22	6%	89.4	22	7%	104.1	26	7%
35-39	109.8	26	8%	123.5	29	8%	142.9	33	10%	112.6	26	8%	123.8	29	8%
40-44	133.6	31	9%	143.4	34	10%	124.9	30	9%	155.9	38	11%	177.1	43	12%
45-49	152.1	37	11%	159.2	37	11%	136.8	31	9%	102.3	23	7%	117.9	27	7%
50-54	219.0	57	17%	198.8	52	15%	207.7	54	16%	202.7	52	15%	203.4	51	14%
55-64	163.2	97	29%	176.6	103	29%	175.6	100	29%	179.4	100	30%	181.3	99	27%
65+	38.7	32	10%	53.1	43	12%	59.5	49	14%	65.4	55	16%	82.9	71	19%
Race/Ethnicity ^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	1	<1%
Asian/Pacific Islander	*	2	1%	*	3	1%	*	2	1%	*	3	1%	*	4	1%
Black/African American	244.0	25	7%	278.0	29	8%	266.3	28	8%	241.1	26	8%	272.6	30	8%
Hispanic/Latino	287.9	15	4%	284.3	15	4%	258.0	14	4%	274.2	16	5%	190.9	12	3%
White	70.1	272	81%	74.2	283	81%	74.0	279	82%	73.5	278	82%	81.1	306	83%
Multi-Race	184.4	16	5%	180.8	16	5%	156.2	14	4%	118.2	11	3%	125.4	12	3%
Unknown	*	4	1%	*	4	1%	*	4	1%	*	4	1%	*	4	1%
Race/Ethnicity ^b and Sex															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	1	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	2	1%	*	3	1%	*	2	1%	*	3	1%	*	4	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African American Males	280.2	18	5%	315.9	21	6%	294.4	20	6%	260.2	18	5%	314.0	22	6%
Black/African American Females	183.2	7	2%	211.4	8	2%	215.1	8	2%	207.0	8	2%	200.2	8	2%
Hispanic/Latino Males	447.8	12	4%	476.2	13	4%	385.2	11	3%	391.5	12	4%	296.2	10	3%
Hispanic/Latina Females	*	3	1%	*	2	1%	*	3	1%	*	4	1%	*	2	1%
White Males	113.6	219	66%	119.6	228	65%	118.6	224	66%	115.9	219	65%	127.3	240	65%
White Females	27.1	53	16%	28.8	55	16%	29.2	55	16%	31.1	59	17%	35.0	66	18%
Multi-Race Males	301.0	13	4%	295.2	13	4%	264.9	12	4%	212.6	10	3%	205.1	10	3%
Multi-Race Females	*	3	1%	*	3	1%	*	2	1%	*	1	<1%	*	2	1%
Unknown	*	4	1%	*	4	1%	*	4	1%	*	4	1%	*	4	1%
Total	80.1	334		85.2	350		83.9	341		82.6	338		90.2	369	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 6, 2020-2024

Living with Diagnosed HIV Infection										
Transmission Category ^a	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	183	70%	192	70%	190	71%	187	72%	196	69%
Injection drug use (IDU)	9	3%	11	4%	10	4%	6	2%	9	3%
Male-to-male sexual contact and IDU	19	7%	21	8%	19	7%	17	7%	23	8%
Heterosexual contact	13	5%	13	5%	12	5%	12	5%	12	4%
Other/unknown	37	14%	38	14%	35	13%	37	14%	44	15%
Subtotal	261	100%	275	100%	266	100%	259	100%	284	100%
Female adult or adolescent										
Injection drug use	10	15%	8	12%	8	12%	10	14%	10	13%
Heterosexual contact	49	73%	54	78%	54	78%	56	77%	62	78%
Other/unknown	8	12%	7	10%	7	10%	7	10%	7	9%
Subtotal	67	100%	69	100%	69	100%	73	100%	79	100%
Child (<13 years at diagnosis)										
Perinatal	5	83%	5	83%	5	83%	5	83%	5	83%
Other/unknown	1	17%	1	17%	1	17%	1	17%	1	17%
Subtotal	6	100%	6	100%	6	100%	6	100%	6	100%
Total	334		350		341		338		369	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 6, 2020-2024

Living with Diagnosed HIV Infection										
Exposure Category ^b	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	136	41%	143	41%	142	42%	141	42%	148	40%
Injection drug use (IDU) only	1	<1%	1	<1%	5	1%	5	1%	5	1%
Heterosexual contact only	76	23%	80	23%	78	23%	82	24%	90	24%
Male-to-male sexual contact & IDU	12	4%	12	3%	12	4%	10	3%	15	4%
IDU & Heterosexual contact	18	5%	18	5%	13	4%	11	3%	14	4%
Male-to-male sexual contact & Heterosexual contact	47	14%	49	14%	48	14%	46	14%	48	13%
Male-to-male sexual contact & IDU & Heterosexual contact	7	2%	9	3%	7	2%	7	2%	8	2%
Perinatal exposure	5	1%	5	1%	5	1%	5	1%	5	1%
Other/unknown	32	10%	33	9%	31	9%	31	9%	36	10%
Total	334		350		341		338		369	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.