

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 6

Counties: Athens, Belmont, Guernsey, Meigs, Monroe, Morgan, Muskingum, Noble, Perry, Washington

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2021

Reported new diagnoses of HIV infection in 2020 by disease status and selected characteristics, HIV Prevention Planning Region 6

Characteristic	2020 diagnosis of HIV infection			Disease Status						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex at birth										
Males	5.3	11	92%	9	90%	2	100%	-	-	
Females	*	1	8%	1	10%	-	-	-	-	
Age at diagnosis (yr)										
<13	*	-	-	-	-	-	-	-	-	
13-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	*	1	8%	1	10%	-	-	-	-	
25-29	*	3	25%	3	30%	-	-	-	-	
30-34	*	1	8%	1	10%	-	-	-	-	
35-39	*	2	17%	1	10%	1	50%	-	-	
40-44	*	2	17%	2	20%	-	-	-	-	
45-49	*	1	8%	-	-	1	50%	-	-	
50-54	*	1	8%	1	10%	-	-	-	-	
55-64	*	1	8%	1	10%	-	-	-	-	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity^b										
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
Black/African-American	*	2	17%	1	10%	1	50%	-	-	
Hispanic/Latinx	*	-	-	-	-	-	-	-	-	
White	2.6	10	83%	9	90%	1	50%	-	-	
Multi-Race	*	-	-	-	-	-	-	-	-	
Race/Ethnicity^b and Sex at birth										
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
Black/African-American Males	*	2	17%	1	10%	1	50%	-	-	
Black/African-American Females	*	-	-	-	-	-	-	-	-	
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
White Males	4.7	9	75%	8	80%	1	50%	-	-	
White Females	*	1	8%	1	10%	-	-	-	-	
Multi-Race Males	*	-	-	-	-	-	-	-	-	
Multi-Race Females	*	-	-	-	-	-	-	-	-	
Total	2.9	12		10		2		-		

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by disease status and transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	8	73%	6	67%	2	100%	-	-
Injection drug use (IDU)	1	9%	1	11%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-
Heterosexual contact	1	9%	1	11%	-	-	-	-
Other/unknown	1	9%	1	11%	-	-	-	-
Subtotal	11	100%	9	100%	2	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	12		10		2		-	

Reported new diagnoses of HIV infection in 2020 by disease status and exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	8	67%	6	60%	2	100%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	1	8%	1	10%	-	-	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	8%	1	10%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	2	17%	2	20%	-	-	-	-
Total	12		10		2		-	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 6

2020 diagnosis of HIV infection

Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	-	-	*	-	-	*	1	10%	*	-	-
25-29	*	-	-	*	-	-	*	1	50%	*	-	-	*	2	20%	*	-	-
30-34	*	-	-	*	-	-	*	-	-	*	-	-	*	1	10%	*	-	-
35-39	*	-	-	*	-	-	*	1	50%	*	-	-	*	1	10%	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	2	20%	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	1	10%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	1	10%	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	1	10%	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	*	2	-	*	-	-	2.6	10	-	*	-	-

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and transmission category, HIV Prevention Planning Region 6

2020 diagnosis of HIV infection

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	2	100%	-	-	6	67%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	11%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	1	11%	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	11%	-	-
Subtotal	-	-	-	-	2	100%	-	-	9	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	100%	-	-
Subtotal	-	-	-	-	-	-	-	-	1	100%	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	2	-	-	-	10	-	-	-

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and exposure category, HIV Prevention Planning Region 6

2020 diagnosis of HIV infection

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	2	100%	-	-	6	60%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	-	-	-	-	1	10%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	10%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	20%	-	-
Total	-	-	-	-	2	-	-	-	10	-	-	-

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 6

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2016			2017			2018			2019			2020			No.	%
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%		
Sex at birth																	
Males	2.8	6	67%	4.3	9	69%	2.4	5	71%	4.3	9	75%	5.3	11	92%	395	81%
Females	*	3	33%	*	4	31%	*	2	29%	*	3	25%	*	1	8%	92	19%
Age at diagnosis (yr)																	
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	6	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	7	1%
20-24	*	4	44%	*	4	31%	*	1	14%	*	1	8%	*	1	8%	53	11%
25-29	*	1	11%	*	4	31%	*	-	-	*	1	8%	*	3	25%	77	16%
30-34	*	1	11%	*	-	-	*	2	29%	*	-	-	*	1	8%	80	16%
35-39	*	-	-	*	3	23%	*	1	14%	*	3	25%	*	2	17%	86	18%
40-44	*	-	-	*	1	8%	*	-	-	*	4	33%	*	2	17%	70	14%
45-49	*	2	22%	*	1	8%	*	3	43%	*	2	17%	*	1	8%	48	10%
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	1	8%	25	5%
55-64	*	1	11%	*	-	-	*	-	-	*	1	8%	*	1	8%	25	5%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	8	2%
Race/Ethnicity^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African-American	*	-	-	*	1	8%	*	1	14%	*	2	17%	*	2	17%	39	8%
Hispanic/Latinx	*	1	11%	*	1	8%	*	-	-	*	-	-	*	-	-	14	3%
White	2.0	8	89%	2.5	10	77%	1.5	6	86%	2.3	9	75%	2.6	10	83%	421	86%
Multi-Race	*	-	-	*	1	8%	*	-	-	*	1	8%	*	-	-	9	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	1%
Race/Ethnicity^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African-American Males	*	-	-	*	-	-	*	-	-	*	1	8%	*	2	17%	23	5%
Black/African-American Females	*	-	-	*	1	8%	*	1	14%	*	1	8%	*	-	-	16	3%
Hispanic/Latino Males	*	1	11%	*	1	8%	*	-	-	*	-	-	*	-	-	11	2%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	1%
White Males	2.5	5	56%	3.6	7	54%	2.6	5	71%	4.1	8	67%	4.7	9	75%	351	72%
White Females	*	3	33%	*	3	23%	*	1	14%	*	1	8%	*	1	8%	70	14%
Multi-Race Males	*	-	-	*	1	8%	*	-	-	*	-	-	*	-	-	8	2%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	1	8%	*	-	-	1	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	1%
Total	2.1	9		3.1	13		1.7	7		2.9	12		2.9	12		487	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	4	67%	5	56%	3	60%	4	44%	8	73%	240	62%
Injection drug use (IDU)	-	-	1	11%	1	20%	2	22%	1	9%	16	4%
Male-to-male sexual contact and IDU	-	-	2	22%	1	20%	1	11%	-	-	21	5%
Heterosexual contact	-	-	1	11%	-	-	-	-	1	9%	26	7%
Other/unknown	2	33%	-	-	-	-	2	22%	1	9%	87	22%
Subtotal	6	100%	9	100%	5	100%	9	100%	11	100%	390	100%
Female adult or adolescent												
Injection drug use	1	33%	3	75%	1	50%	-	-	-	-	15	16%
Heterosexual contact	2	67%	1	25%	1	50%	3	100%	-	-	63	69%
Other/unknown	-	-	-	-	-	-	-	-	1	100%	13	14%
Subtotal	3	100%	4	100%	2	100%	3	100%	1	100%	91	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	67%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	33%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	9		13		7		12		12		487	

Trends in reported diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sexual contact only	4	44%	4	31%	3	43%	4	33%	8	67%	226	46%
Injection drug use (IDU) only	-	-	1	8%	-	-	-	-	-	-	12	2%
Heterosexual contact only	2	22%	2	15%	1	14%	3	25%	1	8%	89	18%
Male-to-male sexual contact & IDU	-	-	-	-	1	14%	1	8%	-	-	16	3%
IDU & Heterosexual contact	1	11%	3	23%	2	29%	2	17%	1	8%	19	4%
Male-to-male sexual contact & Heterosexual contact	-	-	1	8%	-	-	-	-	-	-	14	3%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	2	15%	-	-	-	-	-	-	5	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	2	22%	-	-	-	-	2	17%	2	17%	101	21%
Total	9		13		7		12		12		487	

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.
 Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 6

Characteristic	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex at birth						
Males	1	50%	1	100%	-	-
Females	1	50%	-	-	1	100%
Age at AIDS diagnosis (yr)						
<13	-	-	-	-	-	-
13-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	1	50%	-	-	1	100%
35-39	-	-	-	-	-	-
40-44	-	-	-	-	-	-
45-49	1	50%	1	100%	-	-
50-54	-	-	-	-	-	-
55-64	-	-	-	-	-	-
65+	-	-	-	-	-	-
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African-American	-	-	-	-	-	-
Hispanic/Latinx	-	-	-	-	-	-
White	2	100%	1	100%	1	100%
Multi-Race	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African-American Males	-	-	-	-	-	-
Black/African-American Females	-	-	-	-	-	-
Hispanic/Latino Males	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-
White Males	1	50%	1	100%	-	-
White Females	1	50%	-	-	1	100%
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
Total	2		1		1	

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 6

Transmission Category ^b	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	-	-	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-
Subtotal	1	100%	1	100%	-	-
Female adult or adolescent						
Injection drug use	1	100%	-	-	1	100%
Heterosexual contact	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	1	100%	-	-	1	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	2		1		1	

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 6

Exposure Category ^c	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-
Injection drug use (IDU) only	1	50%	-	-	1	100%
Heterosexual contact only	-	-	-	-	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	1	50%	1	100%	-	-
Total	2		1		1	

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 6

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex at birth												
Males	3	100%	6	86%	4	100%	3	75%	1	50%	257	87%
Females	-	-	1	14%	-	-	1	25%	1	50%	40	13%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	2	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	1%
15-19	-	-	-	-	-	-	-	-	-	-	2	1%
20-24	-	-	-	-	-	-	-	-	-	-	5	2%
25-29	-	-	2	29%	-	-	1	25%	-	-	33	11%
30-34	1	33%	-	-	-	-	-	-	1	50%	47	16%
35-39	-	-	2	29%	1	25%	1	25%	-	-	56	19%
40-44	-	-	-	-	-	-	1	25%	-	-	61	21%
45-49	-	-	1	14%	2	50%	1	25%	1	50%	37	12%
50-54	2	67%	-	-	1	25%	-	-	-	-	25	8%
55-64	-	-	1	14%	-	-	-	-	-	-	22	7%
65+	-	-	1	14%	-	-	-	-	-	-	5	2%
Race/Ethnicity^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
Black/African-American	-	-	1	14%	-	-	1	25%	-	-	20	7%
Hispanic/Latinx	-	-	1	14%	-	-	-	-	-	-	5	2%
White	3	100%	4	57%	4	100%	3	75%	2	100%	266	90%
Multi-Race	-	-	1	14%	-	-	-	-	-	-	6	2%
Race/Ethnicity^a and Sex at birth												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
Black/African-American Males	-	-	-	-	-	-	1	25%	-	-	15	5%
Black/African-American Females	-	-	1	14%	-	-	-	-	-	-	5	2%
Hispanic/Latino Males	-	-	1	14%	-	-	-	-	-	-	4	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	1	<1%
White Males	3	100%	4	57%	4	100%	2	50%	1	50%	232	78%
White Females	-	-	-	-	-	-	1	25%	1	50%	34	11%
Multi-Race Males	-	-	1	14%	-	-	-	-	-	-	6	2%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	-	-
Total	3		7		4		4		2		297	

Notes:

AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	2	67%	5	83%	2	50%	1	33%	-	-	153	60%
Injection drug use (IDU)	-	-	-	-	1	25%	-	-	-	-	12	5%
Male-to-male sexual contact and IDU	-	-	-	-	1	25%	-	-	-	-	17	7%
Heterosexual contact	-	-	1	17%	-	-	-	-	-	-	18	7%
Other/unknown	1	33%	-	-	-	-	2	67%	1	100%	53	21%
Subtotal	3	100%	6	100%	4	100%	3	100%	1	100%	253	100%
Female adult or adolescent												
Injection drug use	-	-	1	100%	-	-	-	-	1	100%	8	20%
Heterosexual contact	-	-	-	-	-	-	1	100%	-	-	28	70%
Other/unknown	-	-	-	-	-	-	-	-	-	-	4	10%
Subtotal	-	-	1	100%	-	-	1	100%	1	100%	40	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	2	50%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	50%
Subtotal	-	-	-	-	-	-	-	-	-	-	4	100%
Total	3		7		4		4		2		297	

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sexual contact only	2	67%	5	71%	2	50%	1	25%	-	-	146	49%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	1	50%	10	3%
Heterosexual contact only	-	-	1	14%	-	-	1	25%	-	-	46	15%
Male-to-male sexual contact & IDU	-	-	-	-	1	25%	-	-	-	-	14	5%
IDU & Heterosexual contact	-	-	1	14%	1	25%	-	-	-	-	10	3%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	7	2%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	3	1%
Perinatal exposure	1	33%	-	-	-	-	-	-	-	-	3	1%
Other/unknown	-	-	-	-	-	-	2	50%	1	50%	58	20%
Total	3		7		4		4		2		297	

Notes:

AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode or transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 6

Counties: Athens, Belmont, Guernsey, Meigs, Monroe, Morgan, Muskingum, Noble, Perry, Washington

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, HIV Prevention Planning Region 6

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	135.3	282	80%	142	78%	140	82%
Females	34.5	72	20%	41	22%	31	18%
Age at end of year							
<13	*	3	1%	2	1%	1	1%
13-14	*	1	<1%	1	1%	-	-
15-19	*	1	<1%	1	1%	-	-
20-24	15.8	5	1%	5	3%	-	-
25-29	74.6	19	5%	15	8%	4	2%
30-34	121.1	29	8%	21	11%	8	5%
35-39	118.2	28	8%	18	10%	10	6%
40-44	146.5	34	10%	21	11%	13	8%
45-49	156.2	38	11%	23	13%	15	9%
50-54	226.7	59	17%	24	13%	35	20%
55-64	174.9	104	29%	39	21%	65	38%
65+	40.0	33	9%	13	7%	20	12%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	4	1%	3	2%	1	1%
Black/African-American	292.8	30	8%	12	7%	18	11%
Hispanic/Latinx	268.7	14	4%	9	5%	5	3%
White	74.2	288	81%	151	83%	137	80%
Multi-Race	161.4	14	4%	4	2%	10	6%
Unknown	*	4	1%	4	2%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	4	1%	3	2%	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African-American Males	326.8	21	6%	7	4%	14	8%
Black/African-American Females	235.5	9	3%	5	3%	4	2%
Hispanic/Latino Males	447.8	12	3%	8	4%	4	2%
Hispanic/Latina Females	*	2	1%	1	1%	1	1%
White Males	120.4	232	66%	119	65%	113	66%
White Females	28.7	56	16%	32	17%	24	14%
Multi-Race Males	254.7	11	3%	3	2%	8	5%
Multi-Race Females	*	3	1%	1	1%	2	1%
Unknown	*	4	1%	4	2%	-	-
Total	84.9	354		183		171	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	192	69%	101	72%	91	66%
Injection drug use (IDU)	10	4%	6	4%	4	3%
Male-to-male sexual contact and IDU	21	8%	10	7%	11	8%
Heterosexual contact	13	5%	2	1%	11	8%
Other/unknown	41	15%	21	15%	20	15%
Subtotal	277	100%	140	100%	137	100%
Female adult or adolescent						
Injection drug use	10	14%	3	8%	7	23%
Heterosexual contact	53	75%	30	75%	23	74%
Other/unknown	8	11%	7	18%	1	3%
Subtotal	71	100%	40	100%	31	100%
Child (<13 yrs at diagnosis)						
Perinatal	5	83%	3	100%	2	67%
Other/unknown	1	17%	-	-	1	33%
Subtotal	6	100%	3	100%	3	100%
Total	354		183		171	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	182	51%	97	53%	85	50%
Injection drug use (IDU) only	10	3%	3	2%	7	4%
Heterosexual contact only	66	19%	32	17%	34	20%
Male-to-male sexual contact & IDU	16	5%	8	4%	8	5%
IDU & Heterosexual contact	10	3%	6	3%	4	2%
Male-to-male sexual contact & Heterosexual contact	10	3%	4	2%	6	4%
Male-to-male sexual contact & IDU & Heterosexual contact	5	1%	2	1%	3	2%
Perinatal exposure	5	1%	3	2%	2	1%
Other/unknown	50	14%	28	15%	22	13%
Total	354		183		171	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	2	50%	*	1	3%	*	-	-	*	-	-	*	-	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	1	7%	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	1	3%	*	-	-	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	*	1	3%	*	-	-	*	4	1%	*	-	-	-	-
25-29	*	-	-	*	-	-	*	2	7%	*	2	14%	61.0	14	5%	*	1	7%	-	-
30-34	*	-	-	*	-	-	*	-	-	*	2	14%	118.1	26	9%	*	1	7%	-	-
35-39	*	-	-	*	-	-	*	3	10%	*	2	14%	104.0	23	8%	*	-	-	-	-
40-44	*	-	-	*	-	-	*	1	3%	*	-	-	146.9	32	11%	*	1	7%	-	-
45-49	*	-	-	*	1	25%	*	4	13%	*	1	7%	121.7	28	10%	*	3	21%	1	25%
50-54	*	-	-	*	-	-	1,085.0	6	20%	*	3	21%	193.6	48	17%	*	2	14%	-	-
55-64	*	-	-	*	1	25%	914.4	11	37%	*	-	-	147.7	84	29%	889.7	5	36%	3	75%
65+	*	-	-	*	-	-	*	-	-	*	3	21%	36.4	29	10%	*	1	7%	-	-
Total	*	-	-	*	4	-	292.8	30	-	268.7	14	-	74.2	288	-	161.4	14	-	4	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	10	50%	6	50%	166	72%	10	91%	-	-
Injection drug use (IDU)	-	-	-	-	1	5%	1	8%	8	3%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	1	50%	-	-	1	8%	19	8%	-	-	-	-
Heterosexual contact	-	-	-	-	1	5%	1	8%	11	5%	-	-	-	-
Other/unknown	-	-	1	50%	8	40%	3	25%	26	11%	1	9%	2	100%
Subtotal	-	-	2	100%	20	100%	12	100%	230	100%	11	100%	2	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	1	11%	-	-	8	14%	1	33%	-	-
Heterosexual contact	-	-	-	-	7	78%	1	100%	43	77%	2	67%	-	-
Other/unknown	-	-	-	-	1	11%	-	-	5	9%	-	-	2	100%
Subtotal	-	-	-	-	9	100%	1	100%	56	100%	3	100%	2	100%
Child (<13 yrs at diagnosis)														
Perinatal	-	-	2	100%	1	100%	1	100%	1	50%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	50%	-	-	-	-
Subtotal	-	-	2	100%	1	100%	1	100%	2	100%	-	-	-	-
Total	-	-	4	-	30	-	14	-	288	-	14	-	4	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	9	30%	6	43%	157	55%	10	71%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	1	7%	8	3%	1	7%	-	-
Heterosexual contact only	-	-	-	-	8	27%	2	14%	54	19%	2	14%	-	-
Male-to-male sexual contact & IDU	-	-	1	25%	-	-	-	-	15	5%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	2	7%	-	-	8	3%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	1	3%	-	-	9	3%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	1	7%	4	1%	-	-	-	-
Perinatal exposure	-	-	2	50%	1	3%	1	7%	1	0%	-	-	-	-
Other/unknown	-	-	1	25%	9	30%	3	21%	32	11%	1	7%	4	100%
Total	-	-	4	-	30	-	14	-	288	-	14	-	4	-

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 6, 2016-2020

Living with Diagnosed HIV Infection

Characteristic	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	114.9	243	81%	114.7	242	80%	123.7	260	80%	131.3	275	80%	135.3	282	80%
Females	26.9	57	19%	28.8	61	20%	31.4	66	20%	32.9	69	20%	34.5	72	20%
Age at end of year															
<13	*	2	1%	*	2	1%	*	2	1%	*	1	<1%	*	3	1%
13-14	*	-	-	*	-	-	*	-	-	*	1	<1%	*	1	<1%
15-19	*	-	-	*	-	-	*	1	<1%	*	2	1%	*	1	<1%
20-24	17.8	6	2%	18.0	6	2%	18.5	6	2%	18.8	6	2%	15.8	5	1%
25-29	65.4	17	6%	84.1	22	7%	91.2	24	7%	84.9	22	6%	74.6	19	5%
30-34	92.4	22	7%	67.8	16	5%	72.3	17	5%	92.4	22	6%	121.1	29	8%
35-39	96.8	23	8%	112.1	27	9%	106.9	26	8%	120.8	29	8%	118.2	28	8%
40-44	106.6	25	8%	100.5	23	8%	97.3	22	7%	108.8	25	7%	146.5	34	10%
45-49	163.2	43	14%	172.1	45	15%	178.1	46	14%	175.6	44	13%	156.2	38	11%
50-54	216.9	62	21%	199.1	55	18%	230.6	62	19%	213.0	56	16%	226.7	59	17%
55-64	135.3	83	28%	139.8	86	28%	157.5	96	29%	171.0	103	30%	174.9	104	29%
65+	22.5	17	6%	27.1	21	7%	30.3	24	7%	40.7	33	10%	40.0	33	9%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	2	1%	*	2	1%	*	2	1%	*	2	1%	*	4	1%
Black/African-American	230.6	24	8%	209.6	22	7%	228.5	24	7%	286.2	30	9%	292.8	30	8%
Hispanic/Latinx	253.6	12	4%	267.5	13	4%	301.1	15	5%	297.0	15	4%	268.7	14	4%
White	62.8	248	83%	64.2	253	83%	68.9	270	83%	71.5	279	81%	74.2	288	81%
Multi-Race	124.3	10	3%	110.8	9	3%	132.8	11	3%	165.1	14	4%	161.4	14	4%
Unknown	*	4	1%	*	4	1%	*	4	1%	*	4	1%	*	4	1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	2	1%	*	2	1%	*	2	1%	*	2	1%	*	4	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African-American Males	276.3	18	6%	243.6	16	5%	257.7	17	5%	332.8	22	6%	326.8	21	6%
Black/African-American Females	154.2	6	2%	152.7	6	2%	179.1	7	2%	206.6	8	2%	235.5	9	3%
Hispanic/Latino Males	421.2	10	3%	452.3	11	4%	472.6	12	4%	462.8	12	3%	447.8	12	3%
Hispanic/Latina Females	*	2	1%	*	2	1%	*	3	1%	*	3	1%	*	2	1%
White Males	103.5	203	68%	104.3	204	67%	112.1	218	67%	116.7	226	66%	120.4	232	66%
White Females	22.6	45	15%	24.7	49	16%	26.4	52	16%	27.0	53	15%	28.7	56	16%
Multi-Race Males	200.6	8	3%	173.3	7	2%	217.2	9	3%	261.9	11	3%	254.7	11	3%
Multi-Race Females	*	2	1%	*	2	1%	*	2	1%	*	3	1%	*	3	1%
Unknown	*	4	1%	*	4	1%	*	4	1%	*	4	1%	*	4	1%
Total	70.9	300		71.7	303		77.5	326		82.1	344		84.9	354	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 6, 2016-2020

Living with Diagnosed HIV Infection										
Transmission Category ^a	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	171	71%	172	72%	179	70%	187	69%	192	69%
Injection drug use (IDU)	4	2%	6	3%	10	4%	9	3%	10	4%
Male-to-male sexual contact and IDU	15	6%	14	6%	18	7%	20	7%	21	8%
Heterosexual contact	12	5%	12	5%	13	5%	13	5%	13	5%
Other/unknown	38	16%	35	15%	37	14%	43	16%	41	15%
Subtotal	240	100%	239	100%	257	100%	272	100%	277	100%
Female adult or adolescent										
Injection drug use	8	14%	9	15%	13	20%	10	15%	10	14%
Heterosexual contact	43	77%	46	77%	47	72%	53	78%	53	75%
Other/unknown	5	9%	5	8%	5	8%	5	7%	8	11%
Subtotal	56	100%	60	100%	65	100%	68	100%	71	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	75%	3	75%	3	75%	3	75%	5	83%
Other/unknown	1	25%	1	25%	1	25%	1	25%	1	17%
Subtotal	4	100%	4	100%	4	100%	4	100%	6	100%
Total	300		303		326		344		354	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 6, 2016-2020

Living with Diagnosed HIV Infection										
Exposure Category ^b	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	162	54%	162	53%	170	52%	178	52%	182	51%
Injection drug use (IDU) only	7	2%	6	2%	11	3%	8	2%	10	3%
Heterosexual contact only	55	18%	58	19%	60	18%	66	19%	66	19%
Male-to-male sexual contact & IDU	12	4%	9	3%	14	4%	16	5%	16	5%
IDU & Heterosexual contact	5	2%	9	3%	12	4%	11	3%	10	3%
Male-to-male sexual contact & Heterosexual contact	9	3%	10	3%	9	3%	9	3%	10	3%
Male-to-male sexual contact & IDU & Heterosexual contact	3	1%	5	2%	4	1%	4	1%	5	1%
Perinatal exposure	4	1%	3	1%	3	1%	3	1%	5	1%
Other/unknown	43	14%	41	14%	43	13%	49	14%	50	14%
Total	300		303		326		344		354	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.