

Persons Living with Diagnosed HIV Infection Reported in Richland County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Richland County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	145.3	89	75%	43	77%	46	74%
Females	48.6	29	25%	13	23%	16	26%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	1	1%	1	2%	-	-
15-19	*	-	-	-	-	-	-
20-24	*	2	2%	1	2%	1	2%
25-29	98.4	8	7%	6	11%	2	3%
30-34	133.8	10	8%	6	11%	4	6%
35-39	171.9	12	10%	8	14%	4	6%
40-44	221.4	16	14%	6	11%	10	16%
45-49	85.6	6	5%	2	4%	4	6%
50-54	248.0	18	15%	7	13%	11	18%
55-64	198.0	32	27%	15	27%	17	27%
65+	52.7	13	11%	4	7%	9	15%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-
Black/African-American	377.0	42	36%	18	32%	24	39%
Hispanic/Latinx	*	4	3%	2	4%	2	3%
White	66.9	69	58%	35	63%	34	55%
Multi-Race	*	3	3%	1	2%	2	3%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African-American Males	417.2	29	25%	12	21%	17	27%
Black/African-American Females	310.3	13	11%	6	11%	7	11%
Hispanic/Latino Males	*	3	3%	2	4%	1	2%
Hispanic/Latina Females	*	1	1%	-	-	1	2%
White Males	108.1	55	47%	28	50%	27	44%
White Females	26.8	14	12%	7	13%	7	11%
Multi-Race Males	*	2	2%	1	2%	1	2%
Multi-Race Females	*	1	1%	-	-	1	2%
Unknown	*	-	-	-	-	-	-
Total	97.6	118		56		62	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Richland County

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	57	64%	28	65%	29	63%
Injection drug use (IDU)	1	1%	-	-	1	2%
Male-to-male sexual contact and IDU	3	3%	2	5%	1	2%
Heterosexual contact	9	10%	2	5%	7	15%
Other/unknown	19	21%	11	26%	8	17%
Subtotal	89	100%	43	100%	46	100%
Female adult or adolescent						
Injection drug use	3	12%	-	-	3	20%
Heterosexual contact	17	65%	6	55%	11	73%
Other/unknown	6	23%	5	45%	1	7%
Subtotal	26	100%	11	100%	15	100%
Child (<13 yrs at diagnosis)						
Perinatal	3	100%	2	100%	1	100%
Other/unknown	-	-	-	-	-	-
Subtotal	3	100%	2	100%	1	100%
Total	118		56		62	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Richland County

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	53	45%	26	46%	27	44%
Injection drug use (IDU) only	3	3%	-	-	3	5%
Heterosexual contact only	26	22%	8	14%	18	29%
Male-to-male sexual contact & IDU	3	3%	2	4%	1	2%
IDU & Heterosexual contact	1	1%	-	-	1	2%
Male-to-male sexual contact & Heterosexual contact	4	3%	2	4%	2	3%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	3	3%	2	4%	1	2%
Other/unknown	25	21%	16	29%	9	15%
Total	118		56		62	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Richland County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	-	-	*	<5	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	<5	-	*	1	1%	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	<5	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	-	-	*	<5	-	*	2	3%	*	-	-	*	-	-
25-29	*	-	-	*	-	-	*	3	7%	*	<5	-	78.2	5	7%	*	-	-	*	-	-
30-34	*	-	-	*	-	-	670.5	7	17%	*	<5	-	*	3	4%	*	-	-	*	-	-
35-39	*	-	-	*	-	-	731.7	6	14%	*	<5	-	86.3	5	7%	*	1	33%	*	-	-
40-44	*	-	-	*	-	-	*	3	7%	*	<5	-	179.5	11	16%	*	-	-	*	-	-
45-49	*	-	-	*	-	-	*	4	10%	*	<5	-	*	2	3%	*	-	-	*	-	-
50-54	*	-	-	*	-	-	1,157.0	7	17%	*	<5	-	155.6	10	14%	*	-	-	*	-	-
55-64	*	-	-	*	-	-	529.6	6	14%	*	<5	-	165.0	24	35%	*	2	67%	*	-	-
65+	*	-	-	*	-	-	457.3	6	14%	*	<5	-	26.3	6	9%	*	-	-	*	-	-
Total	*	-	-	*	-	-	377.0	42		*	4		66.9	69		*	3		*	-	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Richland County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	19	66%	1	33%	35	64%	2	100%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	2%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	3%	-	-	2	4%	-	-	-	-
Heterosexual contact	-	-	-	-	3	10%	1	33%	5	9%	-	-	-	-
Other/unknown	-	-	-	-	6	21%	1	33%	12	22%	-	-	-	-
Subtotal	-	-	-	-	29	100%	3	100%	55	100%	2	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	1	8%	-	-	2	18%	-	-	-	-
Heterosexual contact	-	-	-	-	10	77%	-	-	6	55%	1	100%	-	-
Other/unknown	-	-	-	-	2	15%	1	100%	3	27%	-	-	-	-
Subtotal	-	-	-	-	13	100%	1	100%	11	100%	1	100%	-	-
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	-	-	-	-	3	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	
Subtotal	-	-	-	-	-	-	-	-	3	100%	-	-	-	-
Total	-	-	-	-	42		4		69		3		-	-

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Richland County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	17	40%	1	25%	34	49%	1	33%	-	-
Injection drug use (IDU) only	-	-	-	-	1	2%	-	-	2	3%	-	-	-	-
Heterosexual contact only	-	-	-	-	13	31%	1	25%	11	16%	1	33%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	1	2%	-	-	2	3%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	1%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	2	5%	-	-	1	1%	1	33%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	3	4%	-	-	-	-
Other/unknown	-	-	-	-	8	19%	2	50%	15	22%	-	-	-	-
Total	-	-	-	-	42		4		69		3		-	-

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Richland County, 2016-2020

Characteristic	Living with Diagnosed HIV Infection														
	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	125.0	77	73%	129.4	79	71%	136.6	84	73%	134.9	83	72%	145.3	89	75%
Females	48.7	29	27%	55.4	33	29%	52.0	31	27%	53.7	32	28%	48.6	29	25%
Age at end of year															
<13	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	1	1%
15-19	*	-	-	*	-	-	*	3	3%	*	3	3%	*	-	-
20-24	*	2	2%	*	2	2%	*	1	1%	*	1	1%	*	2	2%
25-29	102.7	8	8%	88.8	7	6%	99.2	8	7%	84.9	7	6%	98.4	8	7%
30-34	197.1	14	13%	143.0	10	9%	82.9	6	5%	67.8	5	4%	133.8	10	8%
35-39	205.1	15	14%	209.9	15	13%	236.8	17	15%	212.6	15	13%	171.9	12	10%
40-44	85.7	6	6%	143.1	10	9%	197.3	14	12%	180.4	13	11%	221.4	16	14%
45-49	211.3	16	15%	217.6	16	14%	179.0	13	11%	139.1	10	9%	85.6	6	5%
50-54	172.9	14	13%	191.5	15	13%	196.6	15	13%	202.5	15	13%	248.0	18	15%
55-64	146.0	25	24%	170.7	29	26%	171.9	29	25%	205.1	34	30%	198.0	32	27%
65+	21.9	5	5%	30.1	7	6%	33.9	8	7%	45.9	11	10%	52.7	13	11%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African-American	324.9	37	35%	367.6	41	37%	396.8	45	39%	359.9	41	36%	377.0	42	36%
Hispanic/Latinx	318.5	7	7%	220.6	5	4%	208.6	5	4%	200.7	5	4%	*	4	3%
White	56.7	59	56%	59.9	62	55%	58.9	61	53%	61.9	64	56%	66.9	69	58%
Multi-Race	*	3	3%	*	4	4%	*	4	3%	190.8	5	4%	*	3	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African-American Males	291.5	21	20%	345.2	24	21%	406.0	29	25%	361.7	26	23%	417.2	29	25%
Black/African-American Females	382.2	16	15%	404.9	17	15%	381.2	16	14%	356.7	15	13%	310.3	13	11%
Hispanic/Latino Males	511.9	6	6%	*	4	4%	*	4	3%	*	4	3%	*	3	3%
Hispanic/Latina Females	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
White Males	93.2	48	45%	94.0	48	43%	93.8	48	42%	99.8	51	44%	108.1	55	47%
White Females	21.0	11	10%	26.7	14	13%	24.8	13	11%	24.8	13	11%	26.8	14	12%
Multi-Race Males	*	2	2%	*	3	3%	*	3	3%	*	2	2%	*	2	2%
Multi-Race Females	*	1	1%	*	1	1%	*	1	1%	*	3	3%	*	1	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	87.5	106		92.9	112		95.0	115		94.9	115		97.6	118	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Richland County, 2016-2020

Transmission Category ^a	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	54	70%	53	67%	60	71%	55	66%	57	64%
Injection drug use (IDU)	2	3%	3	4%	1	1%	1	1%	1	1%
Male-to-male sexual contact and IDU	1	1%	2	3%	1	1%	2	2%	3	3%
Heterosexual contact	5	6%	7	9%	8	10%	9	11%	9	10%
Other/unknown	15	19%	14	18%	14	17%	16	19%	19	21%
Subtotal	77	100%	79	100%	84	100%	83	100%	89	100%
Female adult or adolescent										
Injection drug use	3	11%	4	13%	3	10%	2	7%	3	12%
Heterosexual contact	19	68%	22	69%	20	69%	22	73%	17	65%
Other/unknown	6	21%	6	19%	6	21%	6	20%	6	23%
Subtotal	28	100%	32	100%	29	100%	30	100%	26	100%
Child (<13 yrs at diagnosis)										
Perinatal	1	100%	1	100%	2	100%	2	100%	3	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	2	100%	2	100%	3	100%
Total	106		112		115		115		118	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Richland County, 2016-2020

Exposure Category ^b	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	50	47%	49	44%	55	48%	50	43%	53	45%
Injection drug use (IDU) only	3	3%	5	4%	1	1%	2	2%	3	3%
Heterosexual contact only	24	23%	29	26%	28	24%	31	27%	26	22%
Male-to-male sexual contact & IDU	1	1%	1	1%	-	-	1	1%	3	3%
IDU & Heterosexual contact	2	2%	2	2%	3	3%	1	1%	1	1%
Male-to-male sexual contact & Heterosexual contact	4	4%	4	4%	5	4%	5	4%	4	3%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	1%	1	1%	1	1%	-	-
Perinatal exposure	1	1%	1	1%	2	2%	2	2%	3	3%
Other/unknown	21	20%	20	18%	20	17%	22	19%	25	21%
Total	106		112		115		115		118	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.