

Ohio Department of Health Seasonal Influenza Activity Summary

MMWR Week 47

November 17th – November 23rd, 2019

Current Influenza Activity:

Current Ohio Activity Level (Geographic Spread) – Local

Definition: Increased ILI in 1 region; ILI activity in other regions is not increased AND recent (within the past 3 weeks) lab evidence of influenza in region with increased ILI, OR 2 or more institutional outbreaks (ILI or lab confirmed) in 1 region; ILI activity in other regions is not increased AND recent (within the past 3 weeks) lab evidence of influenza in region with the outbreaks; virus activity is no greater than sporadic in other regions.

During MMWR Week 47, public health surveillance data sources indicate minimal intensity for influenza-like illness (ILI) in outpatient settings reported by Ohio’s sentinel providers. The percentage of emergency department visits with patients exhibiting constitutional symptoms are above baseline levels statewide; fever and ILI specified ED visits are also above baseline levels. Reported cases of influenza-associated hospitalizations are above the seasonal threshold*. There were 32 influenza-associated hospitalizations reported during MMWR Week 46.

Ohio Weekly Influenza-associated Hospitalizations by Ohio Public Health Region

| | |
|--------------|-----------|
| Central | 3 |
| East Central | 2 |
| Northeast | 4 |
| Northwest | 10 |
| Southeast | 2 |
| Southwest | 7 |
| West Central | 4 |
| Total | 32 |

Ohio Influenza Activity Summary Dashboard:

| Data Source | Current week value | Percent Change from last week ¹ | # of weeks ² | Trend Chart ³ |
|---|--------------------|--|-------------------------|--------------------------|
| Influenza-like Illness (ILI) Outpatient Data (ILINet Sentinel Provider Visits) | 0.81% | -13.83% | ↓ 2 | |
| Thermometer Sales (National Retail Data Monitor) | 1075 | 14.48% | ↑ 3 | |
| Fever and ILI Specified ED Visits (EpiCenter) | 2.12% | 3.92% | ↑ 10 | |
| Constitutional ED Visits (EpiCenter) | 9.42% | 2.61% | ↑ 6 | |
| Confirmed Influenza-associated Hospitalizations (Ohio Disease Reporting System) | 32 | 28.00% | ↑ 2 | |
| Outpatient Medical Claims Data ⁴ | 0.56% | 5.66% | ↑ 1 | |

¹Interpret percent changes with caution. Large variability may be exhibited in data sources with low weekly values.

²Number of weeks that the % change is increasing or decreasing.

³Black lines represent current week's data; red lines represent baseline averages

⁴Medical Claims Data provided by athenahealth®

*The seasonal threshold is 25 cases of influenza-associated hospitalizations; historical data demonstrate that once the weekly count exceeds 25 cases, the number of weekly cases thereafter will likely not decrease until after the peak of influenza activity for the season

State, Regional, and National Data:

Ohio Surveillance Data:

- **ODH lab** has reported **12 positive** influenza tests from specimens sent from sentinel ILINet providers and hospital clinical labs. 2019-2020 influenza season results: **(1) A/pdmH1N1; (7) A/H3N2; (4) Influenza B;** (through 11/23/2019).
- The **National Respiratory and Enteric Virus Surveillance System (NREVSS)** has reported **12,009** influenza RT-PCR tests performed at participating facilities. 2019-2020 influenza season positive results: **(6) A/pdmH1N1; (2) A/H3N2; (50) Flu A Not Subtyped; and (38) Flu B;** (through 11/23/2019).
- **0 pediatric influenza-associated mortalities** have been reported during the 2019-2020 season (through 11/23/2019).
- No **novel influenza A virus infections** have been reported during the 2019-2020 season (through 11/23/2019).
- Incidence of confirmed **influenza-associated hospitalizations** in 2019-2020 season = **139** (through 11/23/2019).

HHS Regional Surveillance Data*: During week 46 (November 10th – November 16th, 2019), the proportion of outpatient visits for ILI in Region 5 (Ohio is in Region 5) was 1.5%, which is **below** the regional baseline of 1.9%. Ohio (Local for week 47) and Pennsylvania reported Sporadic Activity; West Virginia, Michigan, Kentucky, and Indiana reported Local Activity.

National Surveillance Data*: During week 46 (November 10th – November 16th, 2019), the majority of U.S. states reported Minimal or Low influenza activity. The proportion of outpatient visits for ILI was 2.5%, which is **above** the national baseline of 2.4%. Four of 10 HHS regions reported ILI levels at or above their region-specific baseline level. Nationally, influenza B/Victoria viruses are the most commonly reported influenza viruses among children age 0-4 years (48% of reported viruses) and 5-24 years (56% of reported viruses), while A(H3N2) viruses are the most commonly reported influenza viruses among persons 65 years of age and older (70% of reported viruses). Among adults aged 25-64 years, approximately equal proportions of influenza A(H1N1)pdm09, A(H3N2) and B/Victoria viruses (33%, 24% and 31%, respectively) have been reported

*National-level and regional-level data are reported one week later than Ohio state-level data


FLUVIEW



A Weekly Influenza Surveillance Report Prepared by the Influenza Division

Weekly Influenza Activity Estimates Reported by State and Territorial Epidemiologists*

Week Ending Nov 16, 2019 - Week 46



*This map indicates geographic spread and does not measure the severity of influenza activity.

2019-2020 Influenza Vaccine Components:

| A/B | Virus | Trivalent (High-dose only) | Quadrivalent |
|-----|--|-------------------------------|--------------|
| A | A/Brisbane/02/2018 (H1N1)pdm09-like | X | X |
| A | A/Kansas/14/2017 (H3N2)-like virus | X | X |
| B | B/Colorado/06/2017-like virus (Victoria lineage) | X | X |
| B | Phuket/3073/2013-like (B/Yamagata lineage) | | X |

Antigenic Characterization:

CDC antigenically characterized 13 influenza viruses collected in the U.S. from September 29, 2019 to November 16, 2019.

Influenza A Viruses

- **A (H1N1)pdm09:** Two A(H1N1)pdm09 viruses were antigenically characterized by HI with ferret antisera, and all were antigenically similar (reacting at titers that were within 4-fold of the homologous virus titer) to cell-propagated A/Brisbane/02/2018-like reference viruses representing the A(H1N1)pdm09 component for the 2019-20 Northern Hemisphere influenza vaccines.
- **A (H3N2):** Antigenic characterization is pending.


Influenza B Viruses

- **B/Victoria:** 11 B/Victoria lineage viruses, including viruses from both co-circulating sub-clades, were antigenically characterized by HI with ferret antisera, and seven (63.6%) were antigenically similar to cell-propagated B/Colorado/06/2017-like reference viruses representing the B/Victoria component for the 2019-20 Northern Hemisphere influenza vaccines.
- **B/Yamagata:** Antigenic characterization is pending.

National activity levels and more information can be found at the following CDC pages:

- <http://www.cdc.gov/flu/weekly/usmap.htm>
- <http://www.cdc.gov/flu/>

Positive Influenza Tests (PCR), National Respiratory and Enteric Virus Surveillance System (NREVSS), 2019-2020 Influenza Season, Ohio



**Influenza-Associated Hospitalizations, Ohio
2019-2020 Season***


| County | Influenza-Associated Hospitalizations | Percent of All Influenza-Associated Hospitalizations | Rate per 100,000 Population† | County | Influenza-Associated Hospitalizations | Percent of All Influenza-Associated Hospitalizations | Rate per 100,000 Population† |
|------------|---------------------------------------|--|------------------------------|--------------|---------------------------------------|--|------------------------------|
| ADAMS | 0 | 0.0% | 0.00 | LOGAN | 0 | 0.0% | 0.00 |
| ALLEN | 11 | 7.9% | 10.35 | LORAIN | 6 | 4.3% | 1.99 |
| ASHLAND | 0 | 0.0% | 0.00 | LUCAS | 6 | 4.3% | 1.36 |
| ASHTABULA | 0 | 0.0% | 0.00 | MADISON | 0 | 0.0% | 0.00 |
| ATHENS | 0 | 0.0% | 0.00 | MAHONING | 1 | 0.7% | 0.42 |
| AUGLAIZE | 4 | 2.9% | 8.71 | MARION | 0 | 0.0% | 0.00 |
| BELMONT | 1 | 0.7% | 1.42 | MEDINA | 2 | 1.4% | 1.16 |
| BROWN | 1 | 0.7% | 2.23 | MEIGS | 0 | 0.0% | 0.00 |
| BUTLER | 5 | 3.6% | 1.36 | MERCER | 0 | 0.0% | 0.00 |
| CARROLL | 0 | 0.0% | 0.00 | MIAMI | 0 | 0.0% | 0.00 |
| CHAMPAIGN | 0 | 0.0% | 0.00 | MONROE | 0 | 0.0% | 0.00 |
| CLARK | 3 | 2.2% | 2.17 | MONTGOMERY | 10 | 7.2% | 1.87 |
| CLERMONT | 3 | 2.2% | 1.52 | MORGAN | 0 | 0.0% | 0.00 |
| CLINTON | 1 | 0.7% | 2.38 | MORROW | 0 | 0.0% | 0.00 |
| COLUMBIANA | 2 | 1.4% | 1.85 | MUSKINGUM | 2 | 1.4% | 2.32 |
| COSHOCTON | 1 | 0.7% | 2.71 | NOBLE | 0 | 0.0% | 0.00 |
| CRAWFORD | 0 | 0.0% | 0.00 | OTTAWA | 0 | 0.0% | 0.00 |
| CUYAHOGA | 9 | 6.5% | 0.70 | PAULDING | 0 | 0.0% | 0.00 |
| DARKE | 0 | 0.0% | 0.00 | PERRY | 1 | 0.7% | 2.77 |
| DEFIANCE | 1 | 0.7% | 2.56 | PICKAWAY | 0 | 0.0% | 0.00 |
| DELAWARE | 1 | 0.7% | 0.57 | PIKE | 0 | 0.0% | 0.00 |
| ERIE | 0 | 0.0% | 0.00 | PORTAGE | 1 | 0.7% | 0.62 |
| FAIRFIELD | 5 | 3.6% | 3.42 | PREBLE | 0 | 0.0% | 0.00 |
| FAYETTE | 0 | 0.0% | 0.00 | PUTNAM | 1 | 0.7% | 2.90 |
| FRANKLIN | 7 | 5.0% | 0.60 | RICHLAND | 0 | 0.0% | 0.00 |
| FULTON | 0 | 0.0% | 0.00 | ROSS | 0 | 0.0% | 0.00 |
| GALLIA | 0 | 0.0% | 0.00 | SANDUSKY | 1 | 0.7% | 1.64 |
| GEAUGA | 0 | 0.0% | 0.00 | SCIOTO | 2 | 1.4% | 2.52 |
| GREENE | 3 | 2.2% | 1.86 | SENECA | 0 | 0.0% | 0.00 |
| GUERNSEY | 1 | 0.7% | 2.49 | SHELBY | 2 | 1.4% | 4.05 |
| HAMILTON | 15 | 10.8% | 1.87 | STARK | 2 | 1.4% | 0.53 |
| HANCOCK | 0 | 0.0% | 0.00 | SUMMIT | 2 | 1.4% | 0.37 |
| HARDIN | 0 | 0.0% | 0.00 | TRUMBULL | 5 | 3.6% | 2.38 |
| HARRISON | 0 | 0.0% | 0.00 | TUSCARAWAS | 0 | 0.0% | 0.00 |
| HENRY | 0 | 0.0% | 0.00 | UNION | 0 | 0.0% | 0.00 |
| HIGHLAND | 0 | 0.0% | 0.00 | VAN WERT | 1 | 0.7% | 3.48 |
| HOCKING | 1 | 0.7% | 3.40 | VINTON | 0 | 0.0% | 0.00 |
| HOLMES | 3 | 2.2% | 7.08 | WARREN | 6 | 4.3% | 2.82 |
| HURON | 0 | 0.0% | 0.00 | WASHINGTON | 0 | 0.0% | 0.00 |
| JACKSON | 0 | 0.0% | 0.00 | WAYNE | 1 | 0.7% | 0.87 |
| JEFFERSON | 0 | 0.0% | 0.00 | WILLIAMS | 1 | 0.7% | 2.66 |
| KNOX | 2 | 1.4% | 3.28 | WOOD | 2 | 1.4% | 1.59 |
| LAKE | 0 | 0.0% | 0.00 | WYANDOT | 0 | 0.0% | 0.00 |
| LAWRENCE | 0 | 0.0% | 0.00 | UNKNOWN | 0 | 0.0% | * |
| LICKING | 4 | 2.9% | 2.40 | TOTAL | 139 | 100% | 1.20 |

*2019-2020 Season 09/29/2019 thru 09/26/2020; data through 11/24/2019


† Disease rates were calculated by number of cases per 100,000 residents using 2010 census data.

Source: Ohio Disease Reporting System


Influenza-associated Hospitalizations by Public Health Region and MMWR Week, Ohio, 2019-2020 Influenza Season




Ohio Constitutional ED Visits with 5 Year Baseline Average; 2019-2020




Ohio Fever & ILI Specified ED Visits with 5 Year Baseline Average; 2019-2020




Ohio Outpatient Influenza-like Illness Network (ILINet) with 5 Year Baseline Average; 2019-2020




AthenaHealth: Ohio Influenza Related Outpatient Medical Claims with 5 Year Baseline Average; 2019-2020



Ohio Thermometer Sales with 5 Year Baseline Average; 2019-2020



Ohio Confirmed Influenza-associated Hospitalizations by MMWR Week; 2019-2020 Season (n=139)



Sources of Influenza Surveillance Data

- **National Retail Data Monitor (NRDM)-OTC Drug Purchases:** The NRDM collects over-the-counter (OTC) drug sales information from approximately 1,420 Ohio chain drug stores and grocery stores. For influenza surveillance, thermometer and adult cold relief sales are monitored on a weekly basis.
- **Emergency Department Visits (EpiCenter):** EpiCenter collects emergency department chief complaint data from 180 hospitals and urgent care facilities across Ohio in real time and classifies them into symptom and syndrome categories. Chief complaints from the constitutional syndrome category and the fever + ILI symptoms classifier are analyzed for influenza surveillance.
- **Sentinel Providers (ILINet):** Sentinel providers, through the US Influenza-like Illness Surveillance Network (ILINet), collect outpatient influenza-like illness (ILI) data. ILI is defined as a fever (≥ 100 F), **and** cough and/or sore throat without another known cause. Providers report the total number of patients seen and the number of patients with ILI by age group on a weekly basis. Sentinel providers also submit specimens for influenza testing to the ODH laboratory throughout the influenza season. There are 70 sentinel providers enrolled in Ohio for the 2019-2020 season.
- **ODH Laboratory Surveillance:** The Ohio Department of Health Laboratory reports the number of specimens that test positive for influenza each week. Generally, specimens are submitted by sentinel provider participants. A subset of the positive specimens is sent to CDC for further testing during the season.
- **Influenza-associated Hospitalizations (ODRS):** Influenza-associated hospitalizations are reported to ODH from local health departments and hospitals by direct entry into the Ohio Disease Reporting System (ODRS). Hospitalizations can be used as an indicator of the severity of illness during a particular influenza season. This condition became reportable in 2009.
- **Influenza-associated Pediatric Mortality (ODRS):** Influenza-associated pediatric mortalities are reported into ODRS by local health department and hospital staff. Pediatric deaths can be an indicator of the severity of illness during the influenza season. This condition became reportable in 2005.
- **National Respiratory and Enteric Virus Surveillance System (NREVSS):** The National Respiratory and Enteric Virus Surveillance System (NREVSS) is a laboratory-based system that monitors temporal and geographic patterns associated with the detection of respiratory syncytial virus (RSV), human parainfluenza viruses (HPIV), respiratory and enteric adenoviruses and rotavirus. There are 19 facilities in Ohio that submit data to this system.
- **athenahealth®:** athenahealth is a technology and services company for medical billing and electronic health records. Diagnosis and procedure data from primary care visits are automatically queried to produce influenza related statistics.

Ohio Public Health Regions: These counties comprise the Ohio Public Health Regions described in the figures shown on pages 1 and 5.

| Central | | East Central | | Noth East | North West | | South East | | South West | West Central |
|-----------|----------|--------------|------------|-----------|------------|----------|------------|------------|------------|--------------|
| CRAWFORD | LOGAN | ASHLAND | RICHLAND | ASHTABULA | ALLEN | MERCER | ATHENS | MONROE | ADAMS | CHAMPAIGN |
| DELAWARE | MADISON | CARROLL | STARK | CUYAHOGA | AUGLAIZE | OTTAWA | BELMONT | MORGAN | BROWN | CLARK |
| FAIRFIELD | MARION | COLUMBIANA | SUMMIT | GEAUGA | DEFIANCE | PAULDING | COSHOCTON | MUSKINGUM | BUTLER | DARKE |
| FAYETTE | MORROW | HOLMES | TRUMBULL | LAKE | ERIE | PUTNAM | GALLIA | NOBLE | CLERMONT | GREENE |
| FRANKLIN | PICKAWAY | MAHONING | TUSCARAWAS | LORAIN | FULTON | SANDUSKY | GUERNSEY | PERRY | CLINTON | MIAMI |
| HARDIN | UNION | MEDINA | WAYNE | | HANCOCK | SENECA | HARRISON | PIKE | HAMILTON | MONTGOMERY |
| KNOX | WYANDOT | PORTAGE | | | HENRY | VAN WERT | HOCKING | ROSS | HIGHLAND | PREBLE |
| LICKING | | | | | HURON | WILLIAMS | JACKSON | SCIOTO | WARREN | SHELBY |
| | | | | | LUCAS | WOOD | JEFFERSON | VINTON | | |
| | | | | | | | LAWRENCE | WASHINGTON | | |
| | | | | | | | MEIGS | | | |

If you have any further questions or comments about surveillance for seasonal influenza for the State of Ohio, please contact the Infectious Disease Informatics and Vaccine Preventable Disease Epidemiology Unit at SMED@odh.ohio.gov or call (614) 995-5599.