

Persons Living with Diagnosed HIV Infection Reported in Greene County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Greene County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	221.4	186	85%	99	82%	87	89%
Females	38.3	33	15%	22	18%	11	11%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	1	<1%	1	1%	-	-
20-24	51.3	7	3%	6	5%	1	1%
25-29	292.0	34	16%	29	24%	5	5%
30-34	153.5	17	8%	11	9%	6	6%
35-39	197.4	21	10%	14	12%	7	7%
40-44	226.0	21	10%	12	10%	9	9%
45-49	185.5	17	8%	6	5%	11	11%
50-54	240.4	24	11%	9	7%	15	15%
55-64	218.6	50	23%	23	19%	27	28%
65+	88.0	27	12%	10	8%	17	17%
Race/Ethnicity^b							
American Indian/Alaska Native	*	1	<1%	1	1%	-	-
Asian/Pacific Islander	*	1	<1%	1	1%	-	-
Black/African-American	497.7	60	27%	34	28%	26	27%
Hispanic/Latinx	167.2	9	4%	5	4%	4	4%
White	98.1	139	63%	76	63%	63	64%
Multi-Race	174.7	9	4%	4	3%	5	5%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	<1%	1	1%	-	-
Asian/Pacific Islander Males	*	1	<1%	1	1%	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African-American Males	880.8	52	24%	29	24%	23	23%
Black/African-American Females	130.0	8	4%	5	4%	3	3%
Hispanic/Latino Males	333.3	9	4%	5	4%	4	4%
Hispanic/Latina Females	*	-	-	-	-	-	-
White Males	163.9	115	53%	60	50%	55	56%
White Females	33.6	24	11%	16	13%	8	8%
Multi-Race Males	349.7	9	4%	4	3%	5	5%
Multi-Race Females	*	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-
Total	128.7	219		121		98	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Greene County

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	134	73%	75	77%	59	69%
Injection drug use (IDU)	2	1%	1	1%	1	1%
Male-to-male sexual contact and IDU	8	4%	4	4%	4	5%
Heterosexual contact	8	4%	2	2%	6	7%
Other/unknown	32	17%	16	16%	16	19%
Subtotal	184	100%	98	100%	86	100%
Female adult or adolescent						
Injection drug use	7	22%	5	24%	2	18%
Heterosexual contact	23	72%	14	67%	9	82%
Other/unknown	2	6%	2	10%	-	-
Subtotal	32	100%	21	100%	11	100%
Child (<13 yrs at diagnosis)						
Perinatal	2	67%	2	100%	-	-
Other/unknown	1	33%	-	-	1	100%
Subtotal	3	100%	2	100%	1	100%
Total	219		121		98	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Greene County

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	130	59%	72	60%	58	59%
Injection drug use (IDU) only	4	2%	3	2%	1	1%
Heterosexual contact only	31	14%	16	13%	15	15%
Male-to-male sexual contact & IDU	7	3%	4	3%	3	3%
IDU & Heterosexual contact	5	2%	3	2%	2	2%
Male-to-male sexual contact & Heterosexual contact	4	2%	3	2%	1	1%
Male-to-male sexual contact & IDU & Heterosexual contact	1	<1%	-	-	1	1%
Perinatal exposure	2	1%	2	2%	-	-
Other/unknown	35	16%	18	15%	17	17%
Total	219		121		98	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Greene County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown			
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	
<13	*	<5	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-
13-14	*	<5	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-
15-19	*	<5	-	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-	*	-	-	-
20-24	*	<5	-	*	-	-	*	1	2%	*	-	-	59.8	6	4%	*	-	-	*	-	-	-
25-29	*	<5	-	*	1	100%	1,081.1	12	20%	*	-	-	216.8	20	14%	*	1	11%	*	-	-	-
30-34	*	<5	-	*	-	-	1,312.3	10	17%	*	2	22%	54.8	5	4%	*	-	-	*	-	-	-
35-39	*	<5	-	*	-	-	969.3	6	10%	*	2	22%	124.9	11	8%	*	2	22%	*	-	-	-
40-44	*	<5	-	*	-	-	1,092.9	6	10%	*	1	11%	153.9	12	9%	*	2	22%	*	-	-	-
45-49	*	<5	-	*	-	-	*	4	7%	*	-	-	165.6	13	9%	*	-	-	*	-	-	-
50-54	*	<5	-	*	-	-	1,306.0	7	12%	*	-	-	148.8	13	9%	*	3	33%	*	-	-	-
55-64	*	<5	-	*	-	-	671.1	8	13%	*	4	44%	182.2	37	27%	*	1	11%	*	-	-	-
65+	*	<5	-	*	-	-	384.3	5	8%	*	-	-	78.9	22	16%	*	-	-	*	-	-	-
Total	*	1	-	*	1	-	497.7	60	-	167.2	9	-	98.1	139	-	174.7	9	-	-	-	-	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Greene County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	1	100%	37	73%	8	89%	83	73%	5	56%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	1	11%	1	1%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	2%	-	-	5	4%	2	22%	-	-
Heterosexual contact	-	-	-	-	6	12%	-	-	2	2%	-	-	-	-
Other/unknown	-	-	-	-	7	14%	-	-	23	20%	2	22%	-	-
Subtotal	-	-	1	100%	51	100%	9	100%	114	100%	9	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	1	13%	-	-	6	26%	-	-	-	-
Heterosexual contact	1	100%	-	-	7	88%	-	-	15	65%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	9%	-	-	-	-
Subtotal	1	100%	-	-	8	100%	-	-	23	100%	-	-	-	-
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	1	100%	-	-	1	50%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	50%	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	2	100%	-	-	-	-
Total	1		1		60		9		139		9			

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Greene County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	1	100%	36	60%	7	78%	81	58%	5	56%	-	-
Injection drug use (IDU) only	-	-	-	-	1	2%	1	11%	2	1%	-	-	-	-
Heterosexual contact only	1	100%	-	-	13	22%	-	-	17	12%	-	-	-	-
Male-to-male sexual contact & IDU	-	-	-	-	1	2%	-	-	4	3%	2	22%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	5	4%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	1	2%	1	11%	2	1%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	1%	-	-	-	-
Perinatal exposure	-	-	-	-	1	2%	-	-	1	1%	-	-	-	-
Other/unknown	-	-	-	-	7	12%	-	-	26	19%	2	22%	-	-
Total	1		1		60		9		139		9			

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Greene County, 2016-2020

Characteristic	Living with Diagnosed HIV Infection														
	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	185.0	150	84%	189.0	155	84%	200.5	166	85%	210.4	175	85%	221.4	186	85%
Females	34.7	29	16%	34.2	29	16%	35.2	30	15%	35.0	30	15%	38.3	33	15%
Age at end of year															
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	2	1%	*	1	1%	*	1	1%	*	1	<1%	*	1	<1%
20-24	73.1	10	6%	72.2	10	5%	50.4	7	4%	36.9	5	2%	51.3	7	3%
25-29	204.2	24	13%	210.8	25	14%	281.0	33	17%	323.0	38	19%	292.0	34	16%
30-34	67.1	7	4%	86.4	9	5%	94.0	10	5%	110.6	12	6%	153.5	17	8%
35-39	224.2	21	12%	90.5	9	5%	154.9	16	8%	201.7	21	10%	197.4	21	10%
40-44	136.8	12	7%	253.7	22	12%	194.4	17	9%	198.5	18	9%	226.0	21	10%
45-49	233.9	23	13%	195.0	19	10%	239.6	23	12%	244.8	23	11%	185.5	17	8%
50-54	225.2	25	14%	258.8	28	15%	211.0	22	11%	205.5	21	10%	240.4	24	11%
55-64	189.3	43	24%	201.2	46	25%	209.2	48	24%	199.3	46	22%	218.6	50	23%
65+	44.3	12	7%	53.7	15	8%	65.9	19	10%	67.5	20	10%	88.0	27	12%
Race/Ethnicity^b															
American Indian/Alaska Native	*	1	1%	*	1	1%	*	1	1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander	*	2	1%	*	1	1%	*	-	-	*	1	<1%	*	1	<1%
Black/African-American	370.5	43	24%	376.1	44	24%	486.9	58	30%	476.6	57	28%	497.7	60	27%
Hispanic/Latinx	173.6	8	4%	171.6	8	4%	182.7	9	5%	179.9	9	4%	167.2	9	4%
White	83.0	115	64%	88.6	124	67%	84.7	119	61%	91.2	129	63%	98.1	139	63%
Multi-Race	230.3	10	6%	129.9	6	3%	185.8	9	5%	164.4	8	4%	174.7	9	4%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	1	1%	*	1	1%	*	1	1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander Males	*	2	1%	*	1	1%	*	-	-	*	1	<1%	*	1	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African-American Males	613.7	35	20%	664.5	38	21%	896.1	52	27%	841.2	49	24%	880.8	52	24%
Black/African-American Females	135.5	8	4%	100.3	6	3%	98.2	6	3%	130.4	8	4%	130.0	8	4%
Hispanic/Latino Males	297.1	7	4%	298.4	7	4%	326.3	8	4%	359.3	9	4%	333.3	9	4%
Hispanic/Latina Females	*	1	1%	*	1	1%	*	1	1%	*	-	-	*	-	-
White Males	140.5	96	54%	149.2	103	56%	139.8	97	49%	154.8	108	53%	163.9	115	53%
White Females	27.1	19	11%	29.6	21	11%	30.9	22	11%	29.3	21	10%	33.6	24	11%
Multi-Race Males	459.1	10	6%	256.4	6	3%	358.4	9	5%	327.2	8	4%	349.7	9	4%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	108.6	179		110.3	184		116.7	196		121.3	205		128.7	219	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Greene County, 2016-2020

Transmission Category ^a	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	113	76%	112	73%	121	74%	128	74%	134	73%
Injection drug use (IDU)	3	2%	3	2%	3	2%	3	2%	2	1%
Male-to-male sexual contact and IDU	5	3%	6	4%	7	4%	8	5%	8	4%
Heterosexual contact	9	6%	9	6%	12	7%	9	5%	8	4%
Other/unknown	18	12%	23	15%	21	13%	25	14%	32	17%
Subtotal	148	100%	153	100%	164	100%	173	100%	184	100%
Female adult or adolescent										
Injection drug use	3	11%	6	21%	5	17%	7	24%	7	22%
Heterosexual contact	22	81%	20	71%	21	72%	21	72%	23	72%
Other/unknown	2	7%	2	7%	3	10%	1	3%	2	6%
Subtotal	27	100%	28	100%	29	100%	29	100%	32	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	75%	2	67%	2	67%	2	67%	2	67%
Other/unknown	1	25%	1	33%	1	33%	1	33%	1	33%
Subtotal	4	100%	3	100%	3	100%	3	100%	3	100%
Total	179		184		196		205		219	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Greene County, 2016-2020

Exposure Category ^b	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	109	61%	109	59%	117	60%	124	60%	130	59%
Injection drug use (IDU) only	3	2%	4	2%	4	2%	4	2%	4	2%
Heterosexual contact only	31	17%	29	16%	33	17%	30	15%	31	14%
Male-to-male sexual contact & IDU	3	2%	4	2%	6	3%	7	3%	7	3%
IDU & Heterosexual contact	3	2%	5	3%	4	2%	6	3%	5	2%
Male-to-male sexual contact & Heterosexual contact	4	2%	3	2%	4	2%	4	2%	4	2%
Male-to-male sexual contact & IDU & Heterosexual contact	2	1%	2	1%	1	1%	1	<1%	1	<1%
Perinatal exposure	3	2%	2	1%	2	1%	2	1%	2	1%
Other/unknown	21	12%	26	14%	25	13%	27	13%	35	16%
Total	179		184		196		205		219	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.