



# Percentage of High School Students Who Felt So Sad or Hopeless Almost Every Day, Two Weeks or More In A Row, That They Stopped Doing Some Usual Activities, During the 12 Months Before the Survey, 2019




For this behavior, based on quadratic trend analyses using logistic regression models controlling for sex, race/ethnicity, and grade ( $p < 0.05$ ), the prevalence did not change from 1999 (28.7%) to 2011 (27.1%) and increased from 2011 (27.1%) to 2019 (33.0%).


# Percentage of High School Students Who Felt So Sad or Hopeless Almost Every Day, Two Weeks or More In A Row, That They Stopped Doing Some Usual Activities, During the 12 Months Before the Survey, 2019




# Percentage of High School Students Who Felt So Sad or Hopeless Almost Every Day, Two Weeks or More In A Row, That They Stopped Doing Some Usual Activities, During the 12 Months Before the Survey, 2019



# Percentage of High School Students Who Felt So Sad or Hopeless Almost Every Day, Two Weeks or More In A Row, That They Stopped Doing Some Usual Activities, During the 12 Months Before the Survey, 2019




# Percentage of High School Students Who Seriously Considered Attempting Suicide During the 12 Months Before the Survey, 2019




For this behavior, based on linear trend analyses using logistic regression models controlling for sex, race/ethnicity, and grade ( $p < 0.05$ ), the prevalence decreased from 1993 (28.1%) to 2019 (15.6%).


# Percentage of High School Students Who Seriously Considered Attempting Suicide During the 12 Months Before the Survey, 2019



# Percentage of High School Students Who Seriously Considered Attempting Suicide During the 12 Months Before the Survey, 2019




# Percentage of High School Students Who Seriously Considered Attempting Suicide During the 12 Months Before the Survey, 2019






# Percentage of High School Students Who Made a Plan About How They Would Attempt Suicide, During the 12 Months Before the Survey, 2019




For this behavior, based on linear trend analyses using logistic regression models controlling for sex, race/ethnicity, and grade ( $p < 0.05$ ), the prevalence decreased from 1993 (21.2%) to 2019 (10.5%).


# Percentage of High School Students Who Made a Plan About How They Would Attempt Suicide, During the 12 Months Before the Survey, 2019




# Percentage of High School Students Who Made a Plan About How They Would Attempt Suicide, During the 12 Months Before the Survey, 2019



# Percentage of High School Students Who Made a Plan About How They Would Attempt Suicide, During the 12 Months Before the Survey, 2019




# Percentage of High School Students Who Attempted Suicide One or More Times During the 12 Months Before the Survey, 2019



Missing bar indicates fewer than 100 students in the subgroup.


For this behavior, based on linear and quadratic trend analyses using logistic regression models controlling for sex, race/ethnicity, and grade ( $p < 0.05$ ), the prevalence decreased from 1993 (10.6%) to 2019 (6.8%).

# Percentage of High School Students Who Attempted Suicide One or More Times During the 12 Months Before the Survey, 2019




Missing bar indicates fewer than 100 students in the subgroup.

# Percentage of High School Students Who Attempted Suicide One or More Times During the 12 Months Before the Survey, 2019



Missing bar indicates fewer than 100 students in the subgroup.

# Percentage of High School Students Who Attempted Suicide One or More Times During the 12 Months Before the Survey, 2019




The prevalence for black students is significantly higher than for white students.

Missing bar indicates fewer than 100 students in the subgroup.




# Percentage of High School Students Who Had a Suicide Attempt That Resulted in an Injury, Poisoning, or Overdose That Had to Be Treated by a Doctor or Nurse, During the 12 Months Before the Survey, 2019




For this behavior, based on linear and quadratic trend analyses using logistic regression models controlling for sex, race/ethnicity, and grade ( $p < 0.05$ ), the prevalence did not change from 1993 (2.8%) to 2019 (2.7%).


# Percentage of High School Students Who Had a Suicide Attempt That Resulted in an Injury, Poisoning, or Overdose That Had to Be Treated by a Doctor or Nurse, During the 12 Months Before the Survey, 2019




# Percentage of High School Students Who Had a Suicide Attempt That Resulted in an Injury, Poisoning, or Overdose That Had to Be Treated by a Doctor or Nurse, During the 12 Months Before the Survey, 2019



# Percentage of High School Students Who Had a Suicide Attempt That Resulted in an Injury, Poisoning, or Overdose That Had to Be Treated by a Doctor or Nurse, During the 12 Months Before the Survey, 2019




# Percentage of High School Students Who Did Something to Purposely Hurt Themselves Without Wanting to Die, During the 12 Months Before the Survey, 2019




For this behavior, based on linear trend analyses using logistic regression models controlling for sex, race/ethnicity, and grade ( $p < 0.05$ ), the prevalence did not change from 2011(16.5%) to 2019(17.7%).


# Percentage of High School Students Who Did Something to Purposely Hurt Themselves Without Wanting to Die, During the 12 Months Before the Survey, 2019



# Percentage of High School Students Who Did Something to Purposely Hurt Themselves Without Wanting to Die, During the 12 Months Before the Survey, 2019




# Percentage of High School Students Who Did Something to Purposely Hurt Themselves Without Wanting to Die, During the 12 Months Before the Survey, 2019






# Percentage of High School Students Who Most of the Time or Always Get the Kind of Help They Need, Among Students Who Report Having Felt Sad, Empty, Hopeless, Angry, or Anxious, 2019



Missing bar indicates fewer than 100 students in the subgroup.


For this behavior, based on linear trend analyses using logistic regression models controlling for sex, race/ethnicity, and grade ( $p < 0.05$ ), the prevalence did not change from 2013(30.2%) to 2019(27.5%).

# Percentage of High School Students Who Most of the Time or Always Get the Kind of Help They Need, Among Students Who Report Having Felt Sad, Empty, Hopeless, Angry, or Anxious, 2019




Missing bar indicates fewer than 100 students in the subgroup.

# Percentage of High School Students Who Most of the Time or Always Get the Kind of Help They Need, Among Students Who Report Having Felt Sad, Empty, Hopeless, Angry, or Anxious, 2019



Missing bar indicates fewer than 100 students in the subgroup.

# Percentage of High School Students Who Most of the Time or Always Get the Kind of Help They Need, Among Students Who Report Having Felt Sad, Empty, Hopeless, Angry, or Anxious, 2019



Missing bar indicates fewer than 100 students in the subgroup.