

New Diagnoses of HIV Infection Reported in Franklin County

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2021

Reported new diagnoses of HIV infection in 2020 by disease status and selected characteristics, Franklin County

Characteristic	2020 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	23.8	154	75%	117	74%	30	81%	7	78%
Females	7.4	50	25%	41	26%	7	19%	2	22%
Age at diagnosis (yr)									
<13	*	1	<1%	1	1%	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	8.4	7	3%	7	4%	-	-	-	-
20-24	37.7	34	17%	31	20%	2	5%	1	11%
25-29	32.3	39	19%	34	22%	5	14%	-	-
30-34	31.7	38	19%	28	18%	8	22%	2	22%
35-39	23.4	23	11%	17	11%	5	14%	1	11%
40-44	16.7	14	7%	11	7%	3	8%	-	-
45-49	20.8	16	8%	11	7%	5	14%	-	-
50-54	17.3	13	6%	6	4%	3	8%	4	44%
55-64	11.4	17	8%	10	6%	6	16%	1	11%
65+	*	2	1%	2	1%	-	-	-	-
Race/Ethnicity^b									
American Indian/Alaska Native	*	1	<1%	1	1%	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
Black/African-American	39.7	123	60%	98	62%	18	49%	7	78%
Hispanic/Latinx	7.6	6	3%	6	4%	-	-	-	-
White	8.5	69	34%	50	32%	17	46%	2	22%
Multi-Race	12.0	5	2%	3	2%	2	5%	-	-
Race/Ethnicity^b and Sex at birth									
American Indian/Alaska Native Males	*	1	<1%	1	1%	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African-American Males	56.1	83	41%	66	42%	12	32%	5	56%
Black/African-American Females	24.7	40	20%	32	20%	6	16%	2	22%
Hispanic/Latino Males	14.7	6	3%	6	4%	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
White Males	15.3	61	30%	43	27%	16	43%	2	22%
White Females	1.9	8	4%	7	4%	1	3%	-	-
Multi-Race Males	*	3	1%	1	1%	2	5%	-	-
Multi-Race Females	*	2	1%	2	1%	-	-	-	-
Total	15.4	204		158		37		9	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by disease status and transmission category, Franklin County

Transmission Category ^a	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	94	61%	73	63%	19	63%	2	29%
Injection drug use (IDU)	9	6%	6	5%	2	7%	1	14%
Male-to-male sexual contact and IDU	14	9%	13	11%	-	-	1	14%
Heterosexual contact	9	6%	6	5%	2	7%	1	14%
Other/unknown	27	18%	18	16%	7	23%	2	29%
Subtotal	153	100%	116	100%	30	100%	7	100%
Female adult or adolescent								
Injection drug use	5	10%	4	10%	1	14%	-	-
Heterosexual contact	44	88%	36	88%	6	86%	2	100%
Other/unknown	1	2%	1	2%	-	-	-	-
Subtotal	50	100%	41	100%	7	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	204		158		37		9	

Reported new diagnoses of HIV infection in 2020 by disease status and exposure category, Franklin County

Exposure Category ^b	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	90	44%	70	44%	18	49%	2	22%
Injection drug use (IDU) only	8	4%	4	3%	3	8%	1	11%
Heterosexual contact only	53	26%	42	27%	8	22%	3	33%
Male-to-male sexual contact & IDU	11	5%	10	6%	-	-	1	11%
IDU & Heterosexual contact	6	3%	6	4%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	4	2%	3	2%	1	3%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	3	1%	3	2%	-	-	-	-
Perinatal exposure	1	<1%	1	1%	-	-	-	-
Other/unknown	28	14%	19	12%	7	19%	2	22%
Total	204		158		37		9	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and age at diagnosis, Franklin County

2020 diagnosis of HIV infection

Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	7	6%	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	121.3	26	21%	*	1	17%	11.5	6	9%	*	1	20%
25-29	*	-	-	*	-	-	80.7	22	18%	*	2	33%	17.6	13	19%	*	2	40%
30-34	*	-	-	*	-	-	105.5	27	22%	*	1	17%	12.0	9	13%	*	1	20%
35-39	*	-	-	*	-	-	50.4	11	9%	*	-	-	18.3	11	16%	*	1	20%
40-44	*	-	-	*	-	-	30.7	6	5%	*	1	17%	13.6	7	10%	*	-	-
45-49	*	1	100%	*	-	-	44.0	8	7%	*	1	17%	12.5	6	9%	*	-	-
50-54	*	-	-	*	-	-	53.2	9	7%	*	-	-	*	4	6%	*	-	-
55-64	*	-	-	*	-	-	15.7	5	4%	*	-	-	11.5	12	17%	*	-	-
65+	*	-	-	*	-	-	*	1	1%	*	-	-	*	1	1%	*	-	-
Total	*	1		*	-		39.7	123		7.6	6		8.5	69		12.0	5	

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and transmission category, Franklin County

2020 diagnosis of HIV infection

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	53	65%	5	83%	34	56%	2	67%
Injection drug use (IDU)	-	-	-	-	1	1%	-	-	7	11%	1	33%
Male-to-male sexual contact and IDU	-	-	-	-	2	2%	-	-	12	20%	-	-
Heterosexual contact	-	-	-	-	7	9%	-	-	2	3%	-	-
Other/unknown	1	100%	-	-	19	23%	1	17%	6	10%	-	-
Subtotal	1	100%	-	-	82	100%	6	100%	61	100%	3	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	2	5%	-	-	3	38%	-	-
Heterosexual contact	-	-	-	-	38	95%	-	-	4	50%	2	100%
Other/unknown	-	-	-	-	-	-	-	-	1	13%	-	-
Subtotal	-	-	-	-	40	100%	-	-	8	100%	2	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-	-	-
Total	1		-		123		6		69		5	

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and exposure category, Franklin County

2020 diagnosis of HIV infection

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	51	41%	5	83%	33	48%	1	20%
Injection drug use (IDU) only	-	-	-	-	2	2%	-	-	5	7%	1	20%
Heterosexual contact only	-	-	-	-	45	37%	-	-	6	9%	2	40%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	11	16%	-	-
IDU & Heterosexual contact	-	-	-	-	1	1%	-	-	5	7%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	2	2%	-	-	1	1%	1	20%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	2	2%	-	-	1	1%	-	-
Perinatal exposure	-	-	-	-	1	1%	-	-	-	-	-	-
Other/unknown	1	100%	-	-	19	15%	1	17%	7	10%	-	-
Total	1		-		123		6		69		5	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by selected characteristics, Franklin County

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2016			2017			2018			2019			2020			No.	%
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%		
Sex at birth																	
Males	26.2	162	82%	27.9	176	79%	23.4	150	75%	27.5	177	82%	23.8	154	75%	6,617	84%
Females	5.4	35	18%	7.1	47	21%	7.3	49	25%	5.6	38	18%	7.4	50	25%	1,256	16%
Age at diagnosis (yr)																	
<13	*	2	1%	*	-	-	*	3	2%	*	-	-	*	1	<1%	50	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	8	<1%
15-19	13.7	11	6%	13.2	11	5%	13.2	11	6%	12.0	10	5%	8.4	7	3%	291	4%
20-24	39.1	35	18%	36.3	33	15%	29.6	27	14%	45.5	41	19%	37.7	34	17%	1,041	13%
25-29	28.7	36	18%	42.6	54	24%	41.0	52	26%	35.7	44	20%	32.3	39	19%	1,428	18%
30-34	22.1	23	12%	29.6	32	14%	31.8	36	18%	27.4	32	15%	31.7	38	19%	1,511	19%
35-39	28.9	26	13%	24.8	23	10%	18.9	18	9%	17.6	17	8%	23.4	23	11%	1,256	16%
40-44	24.3	19	10%	20.3	16	7%	17.5	14	7%	23.3	19	9%	16.7	14	7%	920	12%
45-49	15.1	12	6%	18.7	15	7%	17.5	14	7%	12.7	10	5%	20.8	16	8%	626	8%
50-54	20.6	16	8%	20.9	16	7%	13.2	10	5%	24.1	18	8%	17.3	13	6%	373	5%
55-64	11.1	16	8%	12.9	19	9%	6.7	10	5%	12.0	18	8%	11.4	17	8%	302	4%
65+	*	1	1%	*	4	2%	*	4	2%	3.7	6	3%	*	2	1%	67	1%
Race/Ethnicity^b																	
American Indian/Alaska Native	*	-	-	*	1	<1%	*	-	-	*	-	-	*	1	<1%	7	<1%
Asian/Pacific Islander	*	4	2%	*	3	1%	*	2	1%	6.6	5	2%	*	-	-	52	1%
Black/African-American	37.6	106	54%	38.7	113	51%	37.7	113	57%	38.0	116	54%	39.7	123	60%	3,033	39%
Hispanic/Latinx	16.4	11	6%	19.7	14	6%	20.0	15	8%	22.3	17	8%	7.6	6	3%	321	4%
White	8.4	68	35%	10.4	85	38%	7.8	64	32%	8.4	69	32%	8.5	69	34%	4,192	53%
Multi-Race	22.1	8	4%	18.6	7	3%	12.9	5	3%	20.1	8	4%	12.0	5	2%	247	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	21	<1%
Race/Ethnicity^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	1	<1%	*	-	-	*	-	-	*	1	<1%	5	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Asian/Pacific Islander Males	*	4	2%	*	1	<1%	*	1	1%	13.5	5	2%	*	-	-	38	<1%
Asian/Pacific Islander Females	*	-	-	*	2	1%	*	1	1%	*	-	-	*	-	-	14	<1%
Black/African-American Males	61.3	82	42%	59.1	82	37%	58.8	84	42%	64.1	93	43%	56.1	83	41%	2,216	28%
Black/African-American Females	16.2	24	12%	20.2	31	14%	18.5	29	15%	14.4	23	11%	24.7	40	20%	817	10%
Hispanic/Latino Males	31.6	11	6%	35.4	13	6%	33.3	13	7%	40.6	16	7%	14.7	6	3%	278	4%
Hispanic/Latina Females	*	-	-	*	1	<1%	*	2	1%	*	1	<1%	*	-	-	43	1%
White Males	14.6	58	29%	18.2	73	33%	12.2	49	25%	14.2	57	27%	15.3	61	30%	3,855	49%
White Females	2.4	10	5%	2.9	12	5%	3.6	15	8%	2.9	12	6%	1.9	8	4%	337	4%
Multi-Race Males	39.5	7	4%	32.5	6	3%	*	3	2%	30.7	6	3%	*	3	1%	207	3%
Multi-Race Females	*	1	1%	*	1	<1%	*	2	1%	*	2	1%	*	2	1%	40	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	21	<1%
Total	15.6	197		17.3	223		15.2	199		16.3	215		15.4	204		7,873	

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by transmission category, Franklin County

Transmission Category ^a	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	110	68%	110	63%	102	68%	105	59%	94	61%	4,743	72%
Injection drug use (IDU)	5	3%	9	5%	5	3%	1	1%	9	6%	227	3%
Male-to-male sexual contact and IDU	6	4%	8	5%	8	5%	12	7%	14	9%	269	4%
Heterosexual contact	6	4%	3	2%	3	2%	12	7%	9	6%	276	4%
Other/unknown	34	21%	46	26%	31	21%	47	27%	27	18%	1,077	16%
Subtotal	161	100%	176	100%	149	100%	177	100%	153	100%	6,592	100%
Female adult or adolescent												
Injection drug use	5	15%	9	19%	11	23%	8	21%	5	10%	156	13%
Heterosexual contact	29	85%	37	79%	35	74%	29	76%	44	88%	950	77%
Other/unknown	-	-	1	2%	1	2%	1	3%	1	2%	123	10%
Subtotal	34	100%	47	100%	47	100%	38	100%	50	100%	1,229	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	-	-	3	100%	-	-	1	100%	42	81%
Other/unknown	-	-	-	-	-	-	-	-	-	-	10	19%
Subtotal	2	100%	-	-	3	100%	-	-	1	100%	52	100%
Total	197		223		199		215		204		7,873	

Trends in reported diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by exposure category, Franklin County

Exposure Category ^b	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sexual contact only	110	56%	108	48%	99	50%	101	47%	90	44%	4,568	58%
Injection drug use (IDU) only	5	3%	10	4%	7	4%	4	2%	8	4%	234	3%
Heterosexual contact only	35	18%	40	18%	38	19%	41	19%	53	26%	1,226	16%
Male-to-male sexual contact & IDU	5	3%	6	3%	8	4%	11	5%	11	5%	229	3%
IDU & Heterosexual contact	5	3%	8	4%	9	5%	5	2%	6	3%	149	2%
Male-to-male sexual contact & Heterosexual contact	-	-	2	1%	3	2%	4	2%	4	2%	175	2%
Male-to-male sexual contact & IDU & Heterosexual contact	1	1%	2	1%	-	-	1	<1%	3	1%	40	1%
Perinatal exposure	2	1%	-	-	3	2%	-	-	1	<1%	44	1%
Other/unknown	34	17%	47	21%	32	16%	48	22%	28	14%	1,208	15%
Total	197		223		199		215		204		7,873	

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.
 Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Persons Living with Diagnosed HIV Infection Reported in Franklin County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Franklin County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	677.8	4,385	79%	2,437	79%	1,948	80%
Females	170.3	1,154	21%	653	21%	501	20%
Age at end of year							
<13	4.4	10	<1%	9	<1%	1	<1%
13-14	15.7	5	<1%	1	<1%	4	<1%
15-19	29.9	25	<1%	20	1%	5	<1%
20-24	146.3	132	2%	113	4%	19	1%
25-29	354.0	427	8%	331	11%	96	4%
30-34	575.4	690	12%	481	16%	209	9%
35-39	592.3	581	10%	363	12%	218	9%
40-44	695.0	581	10%	317	10%	264	11%
45-49	795.0	612	11%	306	10%	306	12%
50-54	997.3	751	14%	368	12%	383	16%
55-64	873.3	1,300	23%	586	19%	714	29%
65+	253.2	425	8%	195	6%	230	9%
Race/Ethnicity^b							
American Indian/Alaska Native	289.4	7	<1%	6	<1%	1	<1%
Asian/Pacific Islander	61.3	48	1%	31	1%	17	1%
Black/African-American	806.3	2,501	45%	1,398	45%	1,103	45%
Hispanic/Latinx	459.0	363	7%	198	6%	165	7%
White	288.8	2,348	42%	1,319	43%	1,029	42%
Multi-Race	606.2	252	5%	118	4%	134	5%
Unknown	*	20	<1%	20	1%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	517.7	6	<1%	6	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Males	99.6	38	1%	23	1%	15	1%
Asian/Pacific Islander Females	24.9	10	<1%	8	<1%	2	<1%
Black/African-American Males	1,147.1	1,697	31%	938	30%	759	31%
Black/African-American Females	495.5	804	15%	460	15%	344	14%
Hispanic/Latino Males	740.7	302	5%	170	6%	132	5%
Hispanic/Latina Females	159.2	61	1%	28	1%	33	1%
White Males	531.7	2,118	38%	1,190	39%	928	38%
White Females	55.5	230	4%	129	4%	101	4%
Multi-Race Males	1,007.4	207	4%	93	3%	114	5%
Multi-Race Females	214.1	45	1%	25	1%	20	1%
Unknown	*	20	<1%	20	1%	-	-
Total	418.2	5,539		3,090		2,449	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Franklin County

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	3,120	71%	1,777	73%	1,343	69%
Injection drug use (IDU)	97	2%	38	2%	59	3%
Male-to-male sexual contact and IDU	195	4%	103	4%	92	5%
Heterosexual contact	235	5%	86	4%	149	8%
Other/unknown	717	16%	420	17%	297	15%
Subtotal	4,364	100%	2,424	100%	1,940	100%
Female adult or adolescent						
Injection drug use	108	10%	46	7%	62	13%
Heterosexual contact	921	82%	520	82%	401	82%
Other/unknown	95	8%	71	11%	24	5%
Subtotal	1,124	100%	637	100%	487	100%
Child (<13 yrs at diagnosis)						
Perinatal	45	88%	24	83%	21	95%
Other/unknown	6	12%	5	17%	1	5%
Subtotal	51	100%	29	100%	22	100%
Total	5,539		3,090		2,449	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Franklin County

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,992	54%	1,715	56%	1,277	52%
Injection drug use (IDU) only	95	2%	40	1%	55	2%
Heterosexual contact only	1,157	21%	607	20%	550	22%
Male-to-male sexual contact & IDU	162	3%	84	3%	78	3%
IDU & Heterosexual contact	110	2%	44	1%	66	3%
Male-to-male sexual contact & Heterosexual contact	128	2%	62	2%	66	3%
Male-to-male sexual contact & IDU & Heterosexual contact	33	1%	19	1%	14	1%
Perinatal exposure	47	1%	25	1%	22	1%
Other/unknown	815	15%	494	16%	321	13%
Total	5,539		3,090		2,449	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Franklin County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	2	4%	*	5	<1%	*	2	1%	*	1	<1%	*	-	-	-	-
13-14	*	-	-	*	-	-	*	4	<1%	*	-	-	*	-	-	*	1	<1%	-	-
15-19	*	-	-	*	1	2%	92.6	20	1%	*	2	1%	*	2	<1%	*	-	-	-	-
20-24	*	-	-	*	3	6%	424.6	91	4%	225.7	14	4%	40.3	21	1%	*	3	1%	-	-
25-29	*	1	14%	74.5	6	13%	902.6	246	10%	419.3	30	8%	162.5	120	5%	592.4	24	10%	-	-
30-34	*	1	14%	86.0	8	17%	1,534.9	393	16%	889.2	58	16%	253.8	191	8%	1,298.7	39	15%	-	-
35-39	*	-	-	*	2	4%	1,375.6	300	12%	700.1	43	12%	354.4	213	9%	1,119.2	23	9%	-	-
40-44	*	-	-	108.3	6	13%	1,333.7	261	10%	1,079.7	57	16%	441.3	227	10%	1,734.8	28	11%	2	10%
45-49	*	4	57%	160.1	8	17%	1,544.9	281	11%	821.7	36	10%	530.2	254	11%	2,135.5	29	12%	-	-
50-54	*	-	-	222.1	9	19%	1,597.0	270	11%	1,322.0	42	12%	796.5	398	17%	2,992.3	31	12%	1	5%
55-64	*	1	14%	*	1	2%	1,482.5	473	19%	1,412.8	59	16%	669.1	700	30%	3,302.2	53	21%	13	65%
65+	*	-	-	*	2	4%	531.6	157	6%	691.1	20	6%	172.3	221	9%	1,734.1	21	8%	4	20%
Total	289.4	7		61.3	48		806.3	2,501		459.0	363		288.8	2,348		606.2	252		20	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Franklin County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	5	83%	20	56%	1,049	62%	202	68%	1,699	80%	144	70%	1	6%
Injection drug use (IDU)	-	-	-	-	49	3%	6	2%	36	2%	6	3%	-	-
Male-to-male sexual contact and IDU	-	-	2	6%	48	3%	11	4%	114	5%	20	10%	-	-
Heterosexual contact	-	-	1	3%	150	9%	30	10%	40	2%	14	7%	-	-
Other/unknown	1	17%	13	36%	387	23%	50	17%	227	11%	23	11%	16	94%
Subtotal	6	100%	36	100%	1,683	100%	299	100%	2,116	100%	207	100%	17	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	49	6%	6	10%	48	21%	5	11%	-	-
Heterosexual contact	1	100%	4	50%	668	85%	52	88%	158	70%	38	86%	-	-
Other/unknown	-	-	4	50%	67	9%	1	2%	19	8%	1	2%	3	100%
Subtotal	1	100%	8	100%	784	100%	59	100%	225	100%	44	100%	3	100%
Child (<13 yrs at diagnosis)														
Perinatal	-	-	4	100%	30	88%	4	80%	6	86%	1	100%	-	-
Other/unknown	-	-	-	-	4	12%	1	20%	1	14%	-	-	-	-
Subtotal	-	-	4	100%	34	100%	5	100%	7	100%	1	100%	-	-
Total	7		48		2,501		363		2,348		252		20	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Franklin County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	5	71%	19	40%	972	39%	192	53%	1,665	71%	138	55%	1	5%
Injection drug use (IDU) only	-	-	-	-	51	2%	8	2%	32	1%	4	2%	-	-
Heterosexual contact only	1	14%	5	10%	818	33%	82	23%	199	8%	52	21%	-	-
Male-to-male sexual contact & IDU	-	-	2	4%	39	2%	11	3%	94	4%	16	6%	-	-
IDU & Heterosexual contact	-	-	-	-	47	2%	4	1%	52	2%	7	3%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	1	2%	77	3%	10	3%	34	1%	6	2%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	9	0%	-	-	20	1%	4	2%	-	-
Perinatal exposure	-	-	4	8%	31	1%	4	1%	6	0%	2	1%	-	-
Other/unknown	1	14%	17	35%	457	18%	52	14%	246	10%	23	9%	19	95%
Total	7		48		2,501		363		2,348		252		20	

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Franklin County, 2016-2020

Characteristic	Living with Diagnosed HIV Infection														
	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	646.7	3,991	81%	652.9	4,115	81%	647.6	4,145	80%	659.0	4,236	80%	677.8	4,385	79%
Females	146.1	946	19%	149.6	990	19%	155.5	1,042	20%	161.9	1,091	20%	170.3	1,154	21%
Age at end of year															
<13	5.4	12	<1%	4.9	11	<1%	5.7	13	<1%	4.4	10	<1%	4.4	10	<1%
13-14	*	4	<1%	*	4	<1%	*	4	<1%	15.7	5	<1%	15.7	5	<1%
15-19	29.8	24	<1%	31.2	26	1%	30.0	25	<1%	30.0	25	<1%	29.9	25	<1%
20-24	182.3	163	3%	162.8	148	3%	147.0	134	3%	153.2	138	3%	146.3	132	2%
25-29	378.7	475	10%	408.2	517	10%	390.1	495	10%	370.9	457	9%	354.0	427	8%
30-34	488.9	509	10%	496.5	537	11%	521.3	590	11%	543.9	636	12%	575.4	690	12%
35-39	600.4	540	11%	605.7	562	11%	572.8	546	11%	563.1	543	10%	592.3	581	10%
40-44	675.9	528	11%	667.6	526	10%	653.6	523	10%	670.7	546	10%	695.0	581	10%
45-49	900.7	714	14%	822.8	661	13%	802.3	640	12%	795.8	625	12%	795.0	612	11%
50-54	1,054.1	819	17%	1,066.3	816	16%	1,070.8	810	16%	1,028.0	769	14%	997.3	751	14%
55-64	637.0	921	19%	698.7	1,029	20%	738.7	1,100	21%	811.1	1,212	23%	873.3	1,300	23%
65+	159.1	228	5%	177.1	268	5%	195.1	307	6%	221.3	361	7%	253.2	425	8%
Race/Ethnicity^b															
American Indian/Alaska Native	*	4	<1%	202.0	5	<1%	202.0	5	<1%	245.7	6	<1%	289.4	7	<1%
Asian/Pacific Islander	57.7	37	1%	57.7	40	1%	55.2	41	1%	59.8	45	1%	61.3	48	1%
Black/African-American	738.7	2,083	42%	750.7	2,193	43%	755.4	2,267	44%	774.5	2,362	44%	806.3	2,501	45%
Hispanic/Latinx	409.1	275	6%	420.4	298	6%	417.0	312	6%	442.7	337	6%	459.0	363	7%
White	279.9	2,274	46%	280.7	2,301	45%	279.7	2,293	44%	281.5	2,303	43%	288.8	2,348	42%
Multi-Race	675.4	245	5%	662.4	249	5%	644.8	250	5%	639.5	254	5%	606.2	252	5%
Unknown	*	19	<1%	*	19	<1%	*	19	<1%	*	20	<1%	*	20	<1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	3	<1%	*	4	<1%	*	4	<1%	425.5	5	<1%	517.7	6	<1%
American Indian/Alaska Native Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander Males	88.0	28	1%	91.1	31	1%	84.9	31	1%	94.7	35	1%	99.6	38	1%
Asian/Pacific Islander Females	27.9	9	<1%	25.5	9	<1%	26.5	10	<1%	26.1	10	<1%	24.9	10	<1%
Black/African-American Males	1,075.4	1,439	29%	1,093.2	1,516	30%	1,088.3	1,556	30%	1,113.1	1,616	30%	1,147.1	1,697	31%
Black/African-American Females	434.6	644	13%	441.1	677	13%	452.4	711	14%	466.8	746	14%	495.5	804	15%
Hispanic/Latino Males	635.3	221	4%	658.5	242	5%	653.2	255	5%	705.3	278	5%	740.7	302	5%
Hispanic/Latina Females	166.5	54	1%	164.0	56	1%	159.3	57	1%	160.7	59	1%	159.2	61	1%
White Males	522.1	2,077	42%	522.7	2,097	41%	517.3	2,076	40%	518.3	2,076	39%	531.7	2,118	38%
White Females	47.5	197	4%	48.8	204	4%	51.8	217	4%	54.4	227	4%	55.5	230	4%
Multi-Race Males	1,168.8	207	4%	1,133.5	209	4%	1,086.2	207	4%	1,069.7	209	4%	1,007.4	207	4%
Multi-Race Females	204.7	38	1%	208.9	40	1%	218.1	43	1%	223.0	45	1%	214.1	45	1%
Unknown	*	19	<1%	*	19	<1%	*	19	<1%	*	20	<1%	*	20	<1%
Total	390.4	4,937		395.1	5,105		395.9	5,187		404.6	5,327		418.2	5,539	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Franklin County, 2016-2020

Transmission Category ^a	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	2,904	73%	2,978	73%	2,981	72%	3,008	71%	3,120	71%
Injection drug use (IDU)	83	2%	90	2%	99	2%	96	2%	97	2%
Male-to-male sexual contact and IDU	169	4%	176	4%	173	4%	180	4%	195	4%
Heterosexual contact	213	5%	215	5%	222	5%	234	6%	235	5%
Other/unknown	603	15%	637	16%	651	16%	697	17%	717	16%
Subtotal	3,972	100%	4,096	100%	4,126	100%	4,215	100%	4,364	100%
Female adult or adolescent										
Injection drug use	87	9%	93	10%	96	9%	107	10%	108	10%
Heterosexual contact	750	81%	787	81%	829	81%	860	81%	921	82%
Other/unknown	88	10%	89	9%	94	9%	97	9%	95	8%
Subtotal	925	100%	969	100%	1,019	100%	1,064	100%	1,124	100%
Child (<13 yrs at diagnosis)										
Perinatal	35	88%	36	90%	39	93%	42	88%	45	88%
Other/unknown	5	13%	4	10%	3	7%	6	13%	6	12%
Subtotal	40	100%	40	100%	42	100%	48	100%	51	100%
Total	4,937		5,105		5,187		5,327		5,539	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Franklin County, 2016-2020

Exposure Category ^b	Living with Diagnosed HIV Infection									
	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,760	56%	2,836	56%	2,848	55%	2,884	54%	2,992	54%
Injection drug use (IDU) only	79	2%	82	2%	95	2%	95	2%	95	2%
Heterosexual contact only	964	20%	1,003	20%	1,052	20%	1,095	21%	1,157	21%
Male-to-male sexual contact & IDU	134	3%	141	3%	144	3%	148	3%	162	3%
IDU & Heterosexual contact	91	2%	101	2%	100	2%	108	2%	110	2%
Male-to-male sexual contact & Heterosexual contact	144	3%	142	3%	133	3%	124	2%	128	2%
Male-to-male sexual contact & IDU & Heterosexual contact	35	1%	35	1%	29	1%	32	1%	33	1%
Perinatal exposure	37	1%	38	1%	41	1%	44	1%	47	1%
Other/unknown	693	14%	727	14%	745	14%	797	15%	815	15%
Total	4,937		5,105		5,187		5,327		5,539	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.