

# New Diagnoses of HIV Infection Reported in Franklin County

\*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2024

## Reported new diagnoses of HIV infection in 2023 by disease status and selected characteristics, Franklin County

Characteristic	2023 diagnosis of HIV infection			Disease Status					
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	24.3	158	76%	119	77%	32	73%	7	70%
Females	7.6	51	24%	36	23%	12	27%	3	30%
Age at diagnosis (years)									
0-9	*	-	-	-	-	-	-	-	-
10-14	*	-	-	-	-	-	-	-	-
15-19	11.4	10	5%	10	6%	-	-	-	-
20-24	38.2	34	16%	28	18%	6	14%	-	-
25-29	30.7	34	16%	26	17%	7	16%	1	10%
30-34	36.9	43	21%	32	21%	10	23%	1	10%
35-39	26.2	27	13%	21	14%	4	9%	2	20%
40-44	26.8	24	11%	19	12%	5	11%	-	-
45-49	15.8	12	6%	6	4%	4	9%	2	20%
50-54	14.4	11	5%	8	5%	3	7%	-	-
55-64	8.5	12	6%	3	2%	5	11%	4	40%
65+	*	2	1%	2	1%	-	-	-	-
Race/Ethnicity <sup>b</sup>									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	2	1%	1	1%	1	2%	-	-
Black/African American	36.8	119	57%	85	55%	28	64%	6	60%
Hispanic/Latino	28.0	28	13%	21	14%	7	16%	-	-
White	6.4	50	24%	42	27%	5	11%	3	30%
Multi-Race	22.6	10	5%	6	4%	3	7%	1	10%
Race/Ethnicity <sup>b</sup> and Sex at birth									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	2	1%	1	1%	1	2%	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African American Males	56.5	88	42%	64	41%	19	43%	5	50%
Black/African American Females	18.5	31	15%	21	14%	9	20%	1	10%
Hispanic/Latino Males	48.5	25	12%	18	12%	7	16%	-	-
Hispanic/Latina Females	*	3	1%	3	2%	-	-	-	-
White Males	9.4	36	17%	32	21%	3	7%	1	10%
White Females	3.5	14	7%	10	6%	2	5%	2	20%
Multi-Race Males	32.1	7	3%	4	3%	2	5%	1	10%
Multi-Race Females	*	3	1%	2	1%	1	2%	-	-
Total	15.8	209		155		44		10	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

## Reported new diagnoses of HIV infection in 2023 by disease status and transmission category, Franklin County

Transmission Category <sup>a</sup>	2023 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	93	59%	78	66%	14	44%	1	14%
Injection drug use (IDU)	3	2%	1	1%	2	6%	-	-
Male-to-male sexual contact and IDU	6	4%	6	5%	-	-	-	-
Heterosexual contact	2	1%	1	1%	1	3%	-	-
Other/unknown	54	34%	33	28%	15	47%	6	86%
Subtotal	158	100%	119	100%	32	100%	7	100%
<b>Female adult or adolescent</b>								
Injection drug use	3	6%	2	6%	1	8%	-	-
Heterosexual contact	38	75%	26	72%	10	83%	2	67%
Other/unknown	10	20%	8	22%	1	8%	1	33%
Subtotal	51	100%	36	100%	12	100%	3	100%
<b>Child (&lt;13 years at diagnosis)</b>								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
<b>Total</b>	<b>209</b>		<b>155</b>		<b>44</b>		<b>10</b>	

## Reported new diagnoses of HIV infection in 2023 by disease status and exposure category, Franklin County

Exposure Category <sup>b</sup>	2023 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	70	33%	58	37%	11	25%	1	10%
Injection drug use (IDU) only	1	<1%	1	1%	-	-	-	-
Heterosexual contact only	71	34%	47	30%	19	43%	5	50%
Male-to-male sexual contact & IDU	2	1%	2	1%	-	-	-	-
IDU & Heterosexual contact	5	2%	2	1%	3	7%	-	-
Male-to-male sexual contact & Heterosexual contact	23	11%	20	13%	3	7%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	4	2%	4	3%	-	-	-	-
Perinatal exposure	1	<1%	-	-	1	2%	-	-
Other/unknown	32	15%	21	14%	7	16%	4	40%
<b>Total</b>	<b>209</b>		<b>155</b>		<b>44</b>		<b>10</b>	

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and age at diagnosis, Franklin County

2023 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino <sup>a</sup>			White			Multi-Race		
	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	29.6	7	6%	*	-	-	*	3	6%	*	-	-
20-24	*	-	-	*	-	-	88.1	19	16%	69.9	6	21%	14.4	7	14%	*	2	20%
25-29	*	-	-	*	-	-	79.8	20	17%	70.2	6	21%	10.7	7	14%	*	1	10%
30-34	*	-	-	*	1	50%	84.5	24	20%	74.5	6	21%	16.0	11	22%	*	1	10%
35-39	*	-	-	*	-	-	62.5	15	13%	*	3	11%	11.4	7	14%	*	2	20%
40-44	*	-	-	*	-	-	56.4	12	10%	72.7	5	18%	11.2	6	12%	*	1	10%
45-49	*	-	-	*	-	-	33.2	6	5%	*	1	4%	*	4	8%	*	1	10%
50-54	*	-	-	*	1	50%	32.8	6	5%	*	1	4%	*	2	4%	*	1	10%
55-64	*	-	-	*	-	-	25.2	8	7%	*	-	-	*	3	6%	*	1	10%
65+	*	-	-	*	-	-	*	2	2%	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	2	-	36.8	119	-	28.0	28	-	6.4	50	-	22.6	10	-

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and transmission category, Franklin County

Transmission Category <sup>c</sup>	2023 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	2	100%	45	51%	16	64%	25	69%	5	71%
Injection drug use (IDU)	-	-	-	-	2	2%	1	4%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	4	5%	1	4%	1	3%	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	2	6%	-	-
Other/unknown	-	-	-	-	37	42%	7	28%	8	22%	2	29%
Subtotal	-	-	2	100%	88	100%	25	100%	36	100%	7	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	2	14%	1	33%
Heterosexual contact	-	-	-	-	25	81%	2	67%	10	71%	1	33%
Other/unknown	-	-	-	-	6	19%	1	33%	2	14%	1	33%
Subtotal	-	-	-	-	31	100%	3	100%	14	100%	3	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-		2		119		28		50		10	

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and exposure category, Franklin County

2023 diagnosis of HIV infection												
Exposure Category <sup>d</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	2	100%	33	28%	12	43%	18	36%	5	50%
Injection drug use (IDU) only	-	-	-	-	1	1%	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	48	40%	6	21%	14	28%	3	30%
Male-to-male sexual contact & IDU	-	-	-	-	2	2%	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	1	1%	1	4%	2	4%	1	10%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	12	10%	4	14%	7	14%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	2	2%	1	4%	1	2%	-	-
Perinatal exposure	-	-	-	-	1	1%	-	-	-	-	-	-
Other/unknown	-	-	-	-	19	16%	4	14%	8	16%	1	10%
Total	-	-	2	-	119	-	28	-	50	-	10	-

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

<sup>b</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

<sup>c</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>d</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

## Trends in reported new diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by selected characteristics, Franklin County

Diagnoses of HIV Infection by Year																	
Characteristic	2019			2020			2021			2022			2023			Cumulative Diagnoses	
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	No.	%
Sex at birth																	
Males	27.5	177	82%	24.0	155	75%	21.7	141	73%	22.4	146	75%	24.3	158	76%	7,026	83%
Females	5.6	38	18%	7.5	51	25%	7.6	51	27%	7.2	48	25%	7.6	51	24%	1,396	17%
Age at diagnosis (years)																	
0-9	*	-	-	*	1	<1%	*	-	-	*	1	1%	*	-	-	46	1%
10-14	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	13	<1%
15-19	12.0	10	5%	8.4	7	3%	14.0	12	6%	11.8	10	5%	11.4	10	5%	320	4%
20-24	45.5	41	19%	39.9	36	17%	26.9	24	13%	24.8	23	12%	38.2	34	16%	1,121	13%
25-29	35.7	44	20%	31.5	38	18%	36.7	42	22%	26.7	30	15%	30.7	34	16%	1,522	18%
30-34	27.4	32	15%	31.7	38	18%	25.2	30	16%	28.9	34	18%	36.9	43	21%	1,607	19%
35-39	17.6	17	8%	24.5	24	12%	14.0	14	7%	22.8	23	12%	26.2	27	13%	1,315	16%
40-44	23.3	19	9%	16.7	14	7%	27.7	24	13%	34.1	30	15%	26.8	24	11%	991	12%
45-49	12.7	10	5%	20.8	16	8%	17.2	13	7%	20.0	15	8%	15.8	12	6%	664	8%
50-54	24.1	18	8%	17.3	13	6%	18.1	14	7%	14.3	11	6%	14.4	11	5%	407	5%
55-64	12.0	18	8%	11.4	17	8%	9.5	14	7%	11.1	16	8%	8.5	12	6%	342	4%
65+	3.7	6	3%	*	2	1%	*	4	2%	*	1	1%	*	2	1%	74	1%
Race/Ethnicity <sup>b</sup>																	
American Indian/Alaska Native	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	6	<1%
Asian/Pacific Islander	6.6	5	2%	*	-	-	*	2	1%	*	3	2%	*	2	1%	53	1%
Black/African American	37.4	114	53%	40.3	125	61%	34.5	108	56%	33.2	106	55%	36.8	119	57%	3,318	39%
Hispanic/Latino	22.3	17	8%	7.6	6	3%	15.9	13	7%	24.8	21	11%	28.0	28	13%	392	5%
White	8.3	68	32%	8.4	68	33%	7.1	57	30%	6.6	52	27%	6.4	50	24%	4,296	51%
Multi-Race	27.7	11	5%	16.8	7	3%	25.8	11	6%	27.6	12	6%	22.6	10	5%	342	4%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	15	<1%
Race/Ethnicity <sup>b</sup> and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	4	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Asian/Pacific Islander Males	13.5	5	2%	*	-	-	*	2	1%	*	3	2%	*	2	1%	42	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	11	<1%
Black/African American Males	64.1	93	43%	56.8	84	41%	46.6	70	36%	50.1	77	40%	56.5	88	42%	2,418	29%
Black/African American Females	13.1	21	10%	25.3	41	20%	23.3	38	20%	17.5	29	15%	18.5	31	15%	900	11%
Hispanic/Latino Males	40.6	16	7%	14.7	6	3%	23.7	10	5%	43.4	19	10%	48.5	25	12%	338	4%
Hispanic/Latina Females	*	1	<1%	*	-	-	*	3	2%	*	2	1%	*	3	1%	54	1%
White Males	14.0	56	26%	15.1	60	29%	12.6	50	26%	9.7	38	20%	9.4	36	17%	3,934	47%
White Females	2.9	12	6%	1.9	8	4%	1.7	7	4%	3.5	14	7%	3.5	14	7%	362	4%
Multi-Race Males	35.8	7	3%	24.3	5	2%	37.9	8	4%	41.8	9	5%	32.1	7	3%	276	3%
Multi-Race Females	*	4	2%	*	2	1%	*	3	2%	*	3	2%	*	3	1%	66	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	15	<1%
Total	16.3	215		15.6	206		14.5	192		14.7	194		15.8	209		8,422	

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2019-2023) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2023.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

## Trends in reported new diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by transmission category, Franklin County

Transmission Category <sup>a</sup>	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2019		2020		2021		2022		2023			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	106	60%	96	62%	92	65%	71	49%	93	59%	4,982	71%
Injection drug use (IDU)	1	1%	9	6%	11	8%	7	5%	3	2%	247	4%
Male-to-male sexual contact and IDU	12	7%	14	9%	6	4%	6	4%	6	4%	291	4%
Heterosexual contact	12	7%	9	6%	4	3%	4	3%	2	1%	287	4%
Other/unknown	46	26%	26	17%	28	20%	57	39%	54	34%	1,194	17%
Subtotal	177	100%	154	100%	141	100%	145	100%	158	100%	7,001	100%
<b>Female adult or adolescent</b>												
Injection drug use	8	21%	5	10%	7	14%	6	13%	3	6%	173	13%
Heterosexual contact	29	76%	45	88%	40	78%	38	79%	38	75%	1,063	78%
Other/unknown	1	3%	1	2%	4	8%	4	8%	10	20%	134	10%
Subtotal	38	100%	51	100%	51	100%	48	100%	51	100%	1,370	100%
<b>Child (&lt;13 years at diagnosis)</b>												
Perinatal	-	-	1	100%	-	-	1	100%	-	-	43	84%
Other/unknown	-	-	-	-	-	-	-	-	-	-	8	16%
Subtotal	-	-	1	100%	-	-	1	100%	-	-	51	100%
<b>Total</b>	<b>215</b>		<b>206</b>		<b>192</b>		<b>194</b>		<b>209</b>		<b>8,422</b>	

## Trends in reported diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by exposure category, Franklin County

Exposure Category <sup>b</sup>	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2019		2020		2021		2022		2023			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	75	35%	73	35%	67	35%	53	27%	70	33%	3,984	47%
Injection drug use (IDU) only	2	1%	2	1%	6	3%	3	2%	1	<1%	123	1%
Heterosexual contact only	75	35%	74	36%	61	32%	73	38%	71	34%	1,830	22%
Male-to-male sexual contact & IDU	7	3%	10	5%	5	3%	4	2%	2	1%	189	2%
IDU & Heterosexual contact	7	3%	12	6%	12	6%	10	5%	5	2%	297	4%
Male-to-male sexual contact & Heterosexual contact	31	14%	23	11%	25	13%	18	9%	23	11%	996	12%
Male-to-male sexual contact & IDU & Heterosexual contact	5	2%	4	2%	1	1%	2	1%	4	2%	102	1%
Perinatal exposure	-	-	1	<1%	-	-	2	1%	1	<1%	46	1%
Other/unknown	13	6%	7	3%	15	8%	29	15%	32	15%	855	10%
Total	215		206		192		194		209		8,422	

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2019-2023) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2023.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

# Persons Living with Diagnosed HIV Infection Reported in Franklin County

As of June 30, 2024

## Reported persons living with diagnosed HIV infection in 2023 by current disease status and selected characteristics, Franklin County

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2023			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Sex at birth</b>							
Males	678.0	4,413	78%	2,552	77%	1,861	78%
Females	187.8	1,268	22%	749	23%	519	22%
<b>Age at end of year (years)</b>							
0-9	2.9	5	<1%	5	<1%	-	-
10-14	*	3	<1%	2	<1%	1	<1%
15-19	29.8	26	<1%	19	1%	7	<1%
20-24	178.5	159	3%	145	4%	14	1%
25-29	303.0	336	6%	280	8%	56	2%
30-34	604.3	705	12%	523	16%	182	8%
35-39	656.5	676	12%	434	13%	242	10%
40-44	739.0	662	12%	388	12%	274	12%
45-49	718.5	545	10%	291	9%	254	11%
50-54	836.3	638	11%	322	10%	316	13%
55-64	920.0	1,305	23%	605	18%	700	29%
65+	346.5	621	11%	287	9%	334	14%
<b>Race/Ethnicity<sup>b</sup></b>							
American Indian/Alaska Native	217.6	5	<1%	3	<1%	2	<1%
Asian/Pacific Islander	55.1	43	1%	29	1%	14	1%
Black/African American	821.4	2,659	47%	1,539	47%	1,120	47%
Hispanic/Latino	504.5	504	9%	311	9%	193	8%
White	273.9	2,131	38%	1,237	37%	894	38%
Multi-Race	738.2	326	6%	169	5%	157	7%
Unknown	*	13	<1%	13	<1%	-	-
<b>Race/Ethnicity<sup>b</sup> and Sex at birth</b>							
American Indian/Alaska Native Males	*	4	<1%	3	<1%	1	<1%
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Males	102.6	39	1%	26	1%	13	1%
Asian/Pacific Islander Females	*	4	<1%	3	<1%	1	<1%
Black/African American Males	1,133.0	1,764	31%	995	30%	769	32%
Black/African American Females	532.7	895	16%	544	16%	351	15%
Hispanic/Latino Males	831.0	428	8%	270	8%	158	7%
Hispanic/Latina Females	157.1	76	1%	41	1%	35	1%
White Males	497.7	1,905	34%	1,109	34%	796	33%
White Females	57.2	226	4%	128	4%	98	4%
Multi-Race Males	1,195.7	261	5%	137	4%	124	5%
Multi-Race Females	291.1	65	1%	32	1%	33	1%
Unknown	*	13	<1%	13	<1%	-	-
<b>Total</b>	<b>428.4</b>	<b>5,681</b>		<b>3,301</b>		<b>2,380</b>	

### Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.



## Reported persons living with diagnosed HIV infection in 2023 by current disease status and transmission category, Franklin County

Transmission Category <sup>a</sup>	Living with diagnosed HIV infection in		Current Disease Status			
	2023		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	3,039	69%	1,791	71%	1,248	67%
Injection drug use (IDU)	106	2%	51	2%	55	3%
Male-to-male sexual contact and IDU	176	4%	95	4%	81	4%
Heterosexual contact	217	5%	77	3%	140	8%
Other/unknown	849	19%	521	21%	328	18%
Subtotal	4,387	100%	2,535	100%	1,852	100%
<b>Female adult or adolescent</b>						
Injection drug use	97	8%	42	6%	55	11%
Heterosexual contact	997	81%	582	80%	415	82%
Other/unknown	142	11%	108	15%	34	7%
Subtotal	1,236	100%	732	100%	504	100%
<b>Child (&lt;13 years at diagnosis)</b>						
Perinatal	48	83%	24	71%	24	100%
Other/unknown	10	17%	10	29%	-	-
Subtotal	58	100%	34	100%	24	100%
<b>Total</b>	<b>5,681</b>		<b>3,301</b>		<b>2,380</b>	

## Reported persons living with diagnosed HIV infection in 2023 by current disease status and exposure category, Franklin County

Exposure Category <sup>b</sup>	Living with diagnosed HIV infection in		Current Disease Status			
	2023		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,459	43%	1,464	44%	995	42%
Injection drug use (IDU) only	35	1%	19	1%	16	1%
Heterosexual contact only	1,586	28%	848	26%	738	31%
Male-to-male sexual contact & IDU	110	2%	57	2%	53	2%
IDU & Heterosexual contact	168	3%	74	2%	94	4%
Male-to-male sexual contact & Heterosexual contact	577	10%	327	10%	250	11%
Male-to-male sexual contact & IDU & Heterosexual contact	66	1%	38	1%	28	1%
Perinatal exposure	52	1%	26	1%	26	1%
Other/unknown	628	11%	448	14%	180	8%
<b>Total</b>	<b>5,681</b>		<b>3,301</b>		<b>2,380</b>	

### Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and age, Franklin County

Living with diagnosed HIV infection in 2023																		
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino <sup>a</sup>			White			Multi-Race		
	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%
0-9	*	-	-	*	-	-	*	3	<1%	*	1	<1%	*	1	<1%	*	-	-
10-14	*	-	-	*	-	-	*	2	<1%	*	1	<1%	*	-	-	*	-	-
15-19	*	-	-	*	1	2%	88.9	21	1%	*	2	<1%	*	1	<1%	*	1	<1%
20-24	*	-	-	*	1	2%	486.7	105	4%	267.8	23	5%	49.5	24	1%	142.0	6	2%
25-29	*	1	20%	77.6	6	14%	814.0	204	8%	655.7	56	11%	81.4	53	2%	378.8	16	5%
30-34	*	1	20%	89.1	7	16%	1,341.2	381	14%	968.8	78	15%	275.0	189	9%	1,441.6	49	15%
35-39	*	-	-	*	3	7%	1,495.5	359	14%	1,162.6	86	17%	307.9	189	9%	1,627.7	39	12%
40-44	*	-	-	*	4	9%	1,470.0	313	12%	944.8	65	13%	442.1	236	11%	2,309.7	44	13%
45-49	*	1	20%	101.2	5	12%	1,503.1	272	10%	836.6	49	10%	414.1	188	9%	2,112.7	30	9%
50-54	*	1	20%	196.7	9	21%	1,532.8	280	11%	923.3	42	8%	570.3	271	13%	2,859.5	35	11%
55-64	*	1	20%	74.6	5	12%	1,487.0	473	18%	1,216.8	71	14%	715.7	684	32%	3,808.9	63	19%
65+	*	-	-	*	2	5%	751.1	246	9%	735.3	30	6%	220.5	295	14%	2,749.4	43	13%
Total	217.6	5		55.1	43		821.4	2,659		504.5	504		273.9	2,131		738.2	326	

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and transmission category, Franklin County

Living with diagnosed HIV infection in 2023												
Transmission Category <sup>c</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/ Latino <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	3	75%	15	41%	1,041	59%	277	66%	1,513	80%	189	73%
Injection drug use (IDU)	-	-	-	-	47	3%	6	1%	45	2%	8	3%
Male-to-male sexual contact and IDU	-	-	3	8%	50	3%	10	2%	91	5%	22	8%
Heterosexual contact	1	25%	1	3%	143	8%	24	6%	37	2%	11	4%
Other/unknown	-	-	18	49%	469	27%	105	25%	216	11%	30	12%
Subtotal	4	100%	37	100%	1,750	100%	422	100%	1,902	100%	260	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	42	5%	6	8%	39	18%	10	16%
Heterosexual contact	1	100%	2	50%	731	84%	52	70%	162	74%	49	78%
Other/unknown	-	-	2	50%	100	11%	16	22%	19	9%	4	6%
Subtotal	1	100%	4	100%	873	100%	74	100%	220	100%	63	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	2	100%	29	81%	6	75%	8	89%	3	100%
Other/unknown	-	-	-	-	7	19%	2	25%	1	11%	-	-
Subtotal	-	-	2	100%	36	100%	8	100%	9	100%	3	100%
Total	5		43		2,659		504		2,131		326	

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and exposure category, Franklin County

Living with diagnosed HIV infection in 2023												
Exposure Category <sup>d</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/ Latino <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2	40%	12	28%	748	28%	228	45%	1,321	62%	147	45%
Injection drug use (IDU) only	-	-	-	-	14	1%	3	1%	16	1%	2	1%
Heterosexual contact only	2	40%	9	21%	1,129	42%	115	23%	250	12%	81	25%
Male-to-male sexual contact & IDU	-	-	2	5%	27	1%	7	1%	63	3%	11	3%
IDU & Heterosexual contact	-	-	-	-	75	3%	9	2%	68	3%	16	5%
Male-to-male sexual contact & Heterosexual contact	1	20%	3	7%	292	11%	49	10%	190	9%	42	13%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	2%	23	1%	3	1%	28	1%	11	3%
Perinatal exposure	-	-	2	5%	32	1%	7	1%	8	0%	3	1%
Other/unknown	-	-	14	33%	319	12%	83	16%	187	9%	13	4%
Total	5		43		2,659		504		2,131		326	

Notes:  
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.  
Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.  
<sup>b</sup> The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.  
<sup>c</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.  
<sup>d</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

# Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Franklin County, 2019-2023

Characteristic	Living with Diagnosed HIV Infection														
	2019			2020			2021			2022			2023		
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%
<b>Sex at birth</b>															
Males	647.8	4,164	79%	663.6	4,293	79%	668.8	4,338	79%	666.0	4,333	78%	678.0	4,413	78%
Females	160.0	1,078	21%	166.9	1,131	21%	173.3	1,166	21%	179.7	1,206	22%	187.8	1,268	22%
<b>Age at end of year (years)</b>															
0-9	3.4	6	<1%	3.4	6	<1%	3.4	6	<1%	2.9	5	<1%	2.9	5	<1%
10-14	12.3	10	<1%	11.0	9	<1%	7.1	6	<1%	*	4	<1%	*	3	<1%
15-19	30.0	25	<1%	29.9	25	<1%	43.2	37	1%	37.6	32	1%	29.8	26	<1%
20-24	156.5	141	3%	147.5	133	2%	149.3	133	2%	140.0	130	2%	178.5	159	3%
25-29	368.5	454	9%	348.2	420	8%	311.3	356	6%	294.4	331	6%	303.0	336	6%
30-34	534.5	625	12%	564.6	677	12%	572.5	681	12%	573.2	675	12%	604.3	705	12%
35-39	561.0	541	10%	585.2	574	11%	601.1	600	11%	621.0	626	11%	656.5	676	12%
40-44	660.8	538	10%	683.0	571	11%	700.1	606	11%	714.3	628	11%	739.0	662	12%
45-49	785.6	617	12%	783.3	603	11%	742.1	560	10%	738.4	553	10%	718.5	545	10%
50-54	1,005.3	752	14%	984.1	741	14%	933.4	721	13%	868.1	669	12%	836.3	638	11%
55-64	792.4	1,184	23%	846.5	1,260	23%	891.1	1,316	24%	931.9	1,340	24%	920.0	1,305	23%
65+	213.9	349	7%	241.2	405	7%	287.5	482	9%	312.3	546	10%	346.5	621	11%
<b>Race/Ethnicity<sup>b</sup></b>															
American Indian/Alaska Native	*	4	<1%	*	4	<1%	*	4	<1%	*	4	<1%	217.6	5	<1%
Asian/Pacific Islander	50.5	38	1%	49.8	39	1%	49.8	39	1%	48.0	38	1%	55.1	43	1%
Black/African American	747.9	2,281	44%	773.4	2,399	44%	792.7	2,481	45%	794.0	2,535	46%	821.4	2,659	47%
Hispanic/Latino	434.8	331	6%	447.6	354	7%	458.8	375	7%	495.4	419	8%	504.5	504	9%
White	274.6	2,247	43%	280.2	2,278	42%	280.7	2,255	41%	276.8	2,195	40%	273.9	2,131	38%
Multi-Race	825.8	328	6%	810.7	337	6%	790.6	337	6%	771.3	335	6%	738.2	326	6%
Unknown	*	13	<1%	*	13	<1%	*	13	<1%	*	13	<1%	*	13	<1%
<b>Race/Ethnicity<sup>b</sup> and Sex at birth</b>															
American Indian/Alaska Native Males	*	3	<1%	*	3	<1%	*	3	<1%	*	3	<1%	*	4	<1%
American Indian/Alaska Native Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander Males	86.6	32	1%	86.5	33	1%	86.1	33	1%	87.9	34	1%	102.6	39	1%
Asian/Pacific Islander Females	15.7	6	<1%	15.0	6	<1%	15.0	6	<1%	*	4	<1%	*	4	<1%
Black/African American Males	1,074.6	1,560	30%	1,100.5	1,628	30%	1,111.8	1,669	30%	1,095.8	1,685	30%	1,133.0	1,764	31%
Black/African American Females	451.2	721	14%	475.2	771	14%	498.6	812	15%	513.6	850	15%	532.7	895	16%
Hispanic/Latino Males	679.9	268	5%	706.4	288	5%	730.0	308	6%	797.3	349	6%	831.0	428	8%
Hispanic/Latina Females	171.6	63	1%	172.2	66	1%	169.5	67	1%	171.5	70	1%	157.1	76	1%
White Males	505.5	2,025	39%	516.1	2,056	38%	515.8	2,041	37%	505.5	1,980	36%	497.7	1,905	34%
White Females	53.2	222	4%	53.5	222	4%	52.5	214	4%	53.6	215	4%	57.2	226	4%
Multi-Race Males	1,351.2	264	5%	1,328.6	273	5%	1,288.9	272	5%	1,254.0	270	5%	1,195.7	261	5%
Multi-Race Females	317.1	64	1%	304.4	64	1%	302.0	65	1%	296.8	65	1%	291.1	65	1%
Unknown	*	13	<1%	*	13	<1%	*	13	<1%	*	13	<1%	*	13	<1%
<b>Total</b>	<b>398.1</b>	<b>5,242</b>		<b>409.5</b>	<b>5,424</b>		<b>416.5</b>	<b>5,504</b>		<b>419.0</b>	<b>5,539</b>		<b>428.4</b>	<b>5,681</b>	

Notes:

Living with diagnosed HIV infection by year (2019-2023) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

## Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Franklin County, 2019-2023

Living with Diagnosed HIV Infection										
Transmission Category <sup>a</sup>	2019		2020		2021		2022		2023	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	2,979	72%	3,068	72%	3,087	72%	3,057	71%	3,039	69%
Injection drug use (IDU)	91	2%	94	2%	104	2%	104	2%	106	2%
Male-to-male sexual contact and IDU	178	4%	196	5%	195	5%	189	4%	176	4%
Heterosexual contact	233	6%	238	6%	229	5%	224	5%	217	5%
Other/unknown	662	16%	676	16%	701	16%	735	17%	849	19%
Subtotal	4,143	100%	4,272	100%	4,316	100%	4,309	100%	4,387	100%
<b>Female adult or adolescent</b>										
Injection drug use	109	10%	111	10%	112	10%	109	9%	97	8%
Heterosexual contact	849	81%	899	82%	926	82%	958	82%	997	81%
Other/unknown	92	9%	91	8%	97	9%	107	9%	142	11%
Subtotal	1,050	100%	1,101	100%	1,135	100%	1,174	100%	1,236	100%
<b>Child (&lt;13 years at diagnosis)</b>										
Perinatal	44	90%	46	90%	47	89%	50	89%	48	83%
Other/unknown	5	10%	5	10%	6	11%	6	11%	10	17%
Subtotal	49	100%	51	100%	53	100%	56	100%	58	100%
<b>Total</b>	<b>5,242</b>		<b>5,424</b>		<b>5,504</b>		<b>5,539</b>		<b>5,681</b>	

## Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Franklin County, 2019-2023

Living with Diagnosed HIV Infection										
Exposure Category <sup>b</sup>	2019		2020		2021		2022		2023	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,410	46%	2,478	46%	2,490	45%	2,469	45%	2,459	43%
Injection drug use (IDU) only	31	1%	31	1%	36	1%	36	1%	35	1%
Heterosexual contact only	1,367	26%	1,439	27%	1,483	27%	1,520	27%	1,586	28%
Male-to-male sexual contact & IDU	114	2%	127	2%	126	2%	124	2%	110	2%
IDU & Heterosexual contact	169	3%	174	3%	180	3%	177	3%	168	3%
Male-to-male sexual contact & Heterosexual contact	567	11%	588	11%	595	11%	586	11%	577	10%
Male-to-male sexual contact & IDU & Heterosexual contact	64	1%	69	1%	69	1%	65	1%	66	1%
Perinatal exposure	46	1%	48	1%	49	1%	53	1%	52	1%
Other/unknown	474	9%	470	9%	476	9%	509	9%	628	11%
Total	5,242		5,424		5,504		5,539		5,681	

Notes:

Living with diagnosed HIV infection by year (2019-2023) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.