



Department of  
Health

# New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 11/Ryan White Part A-Columbus

---

**Counties:** Delaware, Fairfield, Franklin, Licking, Madison, Morrow,  
Pickaway, Union

\*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2025

## Reported new diagnoses of HIV infection in 2024 by disease status and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	2024 diagnosis of HIV infection			Disease Status						
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	14.8	159	74%	129	76%	26	60%	4	100%	
Females	5.2	57	26%	40	24%	17	40%	-	-	
Age at diagnosis (years)										
0-9	*	-	-	-	-	-	-	-	-	
10-14	*	1	<1%	1	1%	-	-	-	-	
15-19	7.0	10	5%	10	6%	-	-	-	-	
20-24	25.1	35	16%	31	18%	3	7%	1	25%	
25-29	19.6	31	14%	27	16%	4	9%	-	-	
30-34	24.7	42	19%	33	20%	9	21%	-	-	
35-39	17.7	29	13%	19	11%	9	21%	1	25%	
40-44	13.3	20	9%	13	8%	6	14%	1	25%	
45-49	15.9	21	10%	12	7%	9	21%	-	-	
50-54	8.4	11	5%	10	6%	1	2%	-	-	
55-64	4.9	12	6%	10	6%	1	2%	1	25%	
65+	*	4	2%	3	2%	1	2%	-	-	
Race/Ethnicity <sup>b</sup>										
American Indian/Alaska Native	*	1	<1%	1	1%	-	-	-	-	
Asian/Pacific Islander	*	4	2%	3	2%	1	2%	-	-	
Black/African American	29.4	112	52%	87	51%	23	53%	2	50%	
Hispanic/Latino	22.5	31	14%	25	15%	5	12%	1	25%	
White	4.2	60	28%	46	27%	13	30%	1	25%	
Multi-Race	12.6	8	4%	7	4%	1	2%	-	-	
Race/Ethnicity <sup>b</sup> and Sex										
American Indian/Alaska Native Males	*	1	<1%	1	1%	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	3	1%	3	2%	-	-	-	-	
Asian/Pacific Islander Females	*	1	<1%	-	-	1	2%	-	-	
Black/African American Males	40.7	76	35%	62	37%	12	28%	2	50%	
Black/African American Females	18.5	36	17%	25	15%	11	26%	-	-	
Hispanic/Latino Males	33.8	24	11%	20	12%	3	7%	1	25%	
Hispanic/Latina Females	10.5	7	3%	5	3%	2	5%	-	-	
White Males	6.7	48	22%	36	21%	11	26%	1	25%	
White Females	1.6	12	6%	10	6%	2	5%	-	-	
Multi-Race Males	22.3	7	3%	7	4%	-	-	-	-	
Multi-Race Females	*	1	<1%	-	-	1	2%	-	-	
Total	10.0	216		169		43		4		

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Reported new diagnoses of HIV infection in 2024 by disease status and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category <sup>a</sup>	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>								
Male-to-male sexual contact	93	59%	80	63%	11	42%	2	50%
Injection drug use (IDU)	4	3%	2	2%	2	8%	-	-
Male-to-male sexual contact and IDU	8	5%	6	5%	2	8%	-	-
Heterosexual contact	6	4%	5	4%	1	4%	-	-
Other/unknown	47	30%	35	27%	10	38%	2	50%
Subtotal	158	100%	128	100%	26	100%	4	100%
<b>Female adult or adolescent</b>								
Injection drug use	4	7%	2	5%	2	12%	-	-
Heterosexual contact	45	79%	32	80%	13	76%	-	-
Other/unknown	8	14%	6	15%	2	12%	-	-
Subtotal	57	100%	40	100%	17	100%	-	-
<b>Child (&lt;13 years at diagnosis)</b>								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
<b>Total</b>	<b>216</b>		<b>169</b>		<b>43</b>		<b>4</b>	

## Reported new diagnoses of HIV infection in 2024 by disease status and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category <sup>b</sup>	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	67	31%	61	36%	5	12%	1	25%
Injection drug use (IDU) only	2	1%	1	1%	1	2%	-	-
Heterosexual contact only	77	36%	54	32%	22	51%	1	25%
Male-to-male sexual contact & IDU	6	3%	5	3%	1	2%	-	-
IDU & Heterosexual contact	6	3%	3	2%	3	7%	-	-
Male-to-male sexual contact & Heterosexual contact	26	12%	19	11%	6	14%	1	25%
Male-to-male sexual contact & IDU & Heterosexual contact	2	1%	1	1%	1	2%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	30	14%	25	15%	4	9%	1	25%
<b>Total</b>	<b>216</b>		<b>169</b>		<b>43</b>		<b>4</b>	

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

2024 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino <sup>a</sup>			White			Multi-Race		
	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
0-14	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	17.6	5	4%	*	3	10%	*	1	2%	*	1	13%
20-24	*	1	100%	*	-	-	77.2	20	18%	73.3	9	29%	*	3	5%	*	2	25%
25-29	*	-	-	*	-	-	62.5	18	16%	*	1	3%	9.0	9	15%	*	3	38%
30-34	*	-	-	*	2	50%	53.0	18	16%	61.6	7	23%	14.0	15	25%	*	-	-
35-39	*	-	-	*	1	25%	64.7	19	17%	49.5	5	16%	*	4	7%	*	-	-
40-44	*	-	-	*	1	25%	31.4	8	7%	*	4	13%	6.0	6	10%	*	1	13%
45-49	*	-	-	*	-	-	40.3	9	8%	*	-	-	12.2	11	18%	*	1	13%
50-54	*	-	-	*	-	-	*	4	4%	*	1	3%	6.5	6	10%	*	-	-
55-64	*	-	-	*	-	-	21.7	8	7%	*	-	-	*	4	7%	*	-	-
65+	*	-	-	*	-	-	*	2	2%	*	1	3%	*	1	2%	*	-	-
Total	*	1		*	4		29.4	112		22.5	31		4.2	60		12.6	8	

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category <sup>c</sup>	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	1	100%	2	67%	42	56%	14	58%	28	58%	6	86%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	4	8%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	1	4%	7	15%	-	-
Heterosexual contact	-	-	-	-	4	5%	2	8%	-	-	-	-
Other/unknown	-	-	1	33%	29	39%	7	29%	9	19%	1	14%
Subtotal	1	100%	3	100%	75	100%	24	100%	48	100%	7	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	1	3%	-	-	3	25%	-	-
Heterosexual contact	-	-	1	100%	29	81%	6	86%	8	67%	1	100%
Other/unknown	-	-	-	-	6	17%	1	14%	1	8%	-	-
Subtotal	-	-	1	100%	36	100%	7	100%	12	100%	1	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	100%	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-	-	-
Total	1		4		112		31		60		8	

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

2024 diagnosis of HIV infection												
Exposure Category <sup>d</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	100%	1	25%	31	28%	11	35%	20	33%	3	38%
Injection drug use (IDU) only	-	-	-	-	1	1%	-	-	1	2%	-	-
Heterosexual contact only	-	-	2	50%	51	46%	12	39%	10	17%	2	25%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	1	3%	5	8%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	6	10%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	1	25%	11	10%	3	10%	8	13%	3	38%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	2	3%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	18	16%	4	13%	8	13%	-	-
Total	1		4		112		31		60		8	

Notes:  
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.  
Asterisk (\*) indicates rate not calculated, for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.  
<sup>b</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.  
<sup>c</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.  
<sup>d</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Diagnoses of HIV Infection by Year																	
Characteristic	2020			2021			2022			2023			2024			Cumulative Diagnoses	
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	No.	%
Sex																	
Males	16.7	171	75%	15.0	155	73%	16.3	170	76%	17.2	180	77%	14.8	159	74%	7,942	83%
Females	5.4	57	25%	5.4	57	27%	5.1	54	24%	5.1	54	23%	5.2	57	26%	1,600	17%
Age at diagnosis (years)																	
0-9	*	1	<1%	*	-	-	*	1	<1%	*	-	-	*	-	-	53	1%
10-14	*	-	-	*	1	<1%	*	-	-	*	-	-	*	1	<1%	16	<1%
15-19	6.7	9	4%	9.4	13	6%	8.0	11	5%	7.1	10	4%	7.0	10	5%	357	4%
20-24	27.1	36	16%	19.6	26	12%	17.9	25	11%	28.1	38	16%	25.1	35	16%	1,256	13%
25-29	26.5	43	19%	30.2	47	22%	23.3	36	16%	26.7	41	18%	19.6	31	14%	1,696	18%
30-34	25.3	42	18%	19.9	33	16%	23.5	39	17%	27.6	46	20%	24.7	42	19%	1,799	19%
35-39	18.8	28	12%	11.8	18	8%	21.4	33	15%	20.9	33	14%	17.7	29	13%	1,519	16%
40-44	11.9	16	7%	18.5	26	12%	21.7	31	14%	17.1	25	11%	13.3	20	9%	1,132	12%
45-49	13.2	17	7%	11.0	14	7%	13.5	17	8%	10.1	13	6%	15.9	21	10%	759	8%
50-54	11.0	14	6%	11.4	15	7%	8.3	11	5%	9.2	12	5%	8.4	11	5%	471	5%
55-64	8.1	20	9%	5.6	14	7%	7.4	18	8%	5.8	14	6%	4.9	12	6%	394	4%
65+	*	2	1%	1.7	5	2%	*	2	1%	*	2	1%	*	4	2%	90	1%
Race/Ethnicity <sup>b</sup>																	
American Indian/Alaska Native	*	-	-	*	1	<1%	*	-	-	*	-	-	*	1	<1%	8	<1%
Asian/Pacific Islander	*	-	-	*	2	1%	*	3	1%	*	4	2%	*	4	2%	62	1%
Black/African American	37.8	131	57%	32.7	115	54%	30.5	110	49%	33.2	122	52%	29.4	112	52%	3,588	38%
Hispanic/Latino	7.2	7	3%	13.8	14	7%	20.8	22	10%	25.1	31	13%	22.5	31	14%	453	5%
White	5.5	80	35%	4.8	70	33%	5.0	73	33%	4.8	69	29%	4.2	60	28%	5,054	53%
Multi-Race	17.7	10	4%	17.1	10	5%	26.7	16	7%	13.0	8	3%	12.6	8	4%	362	4%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	15	<1%
Race/Ethnicity <sup>b</sup> and Sex																	
American Indian/Alaska Native Males	*	-	-	*	1	<1%	*	-	-	*	-	-	*	1	<1%	6	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Asian/Pacific Islander Males	*	-	-	*	2	1%	*	3	1%	*	4	2%	*	3	1%	49	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	1	<1%	13	<1%
Black/African American Males	52.3	88	39%	43.0	74	35%	44.7	79	35%	50.5	91	39%	40.7	76	35%	2,614	27%
Black/African American Females	24.1	43	19%	22.8	41	19%	16.9	31	14%	16.5	31	13%	18.5	36	17%	974	10%
Hispanic/Latino Males	14.0	7	3%	21.0	11	5%	36.6	20	9%	44.0	28	12%	33.8	24	11%	386	4%
Hispanic/Latina Females	*	-	-	*	3	1%	*	2	1%	*	3	1%	10.5	7	3%	67	1%
White Males	9.6	69	30%	8.2	59	28%	7.6	55	25%	7.3	52	22%	6.7	48	22%	4,583	48%
White Females	1.5	11	5%	1.5	11	5%	2.5	18	8%	2.3	17	7%	1.6	12	6%	471	5%
Multi-Race Males	25.1	7	3%	27.6	8	4%	43.7	13	6%	16.4	5	2%	22.3	7	3%	290	3%
Multi-Race Females	*	3	1%	*	2	1%	*	3	1%	*	3	1%	*	1	<1%	72	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	15	<1%
Total	11.0	228		10.2	212		10.7	224		11.1	234		10.0	216		9,542	

Notes:  
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category <sup>a</sup>	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	105	62%	101	65%	81	48%	108	60%	93	59%	5,551	70%
Injection drug use (IDU)	9	5%	12	8%	10	6%	4	2%	4	3%	300	4%
Male-to-male sexual contact and IDU	16	9%	9	6%	10	6%	8	4%	8	5%	352	4%
Heterosexual contact	9	5%	4	3%	4	2%	3	2%	6	4%	329	4%
Other/unknown	31	18%	29	19%	64	38%	57	32%	47	30%	1,381	17%
Subtotal	170	100%	155	100%	169	100%	180	100%	158	100%	7,913	100%
<b>Female adult or adolescent</b>												
Injection drug use	7	12%	7	12%	7	13%	4	7%	4	7%	199	13%
Heterosexual contact	49	86%	46	81%	42	78%	41	76%	45	79%	1,210	77%
Other/unknown	1	2%	4	7%	5	9%	9	17%	8	14%	160	10%
Subtotal	57	100%	57	100%	54	100%	54	100%	57	100%	1,569	100%
<b>Child (&lt;13 years at diagnosis)</b>												
Perinatal	1	100%	-	-	1	100%	-	-	-	-	50	83%
Other/unknown	-	-	-	-	-	-	-	-	1	100%	10	17%
Subtotal	1	100%	-	-	1	100%	-	-	1	100%	60	100%
<b>Total</b>	<b>228</b>		<b>212</b>		<b>224</b>		<b>234</b>		<b>216</b>		<b>9,542</b>	

## Trends in reported diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Diagnoses of HIV Infection by Year												
Exposure Category <sup>b</sup>	2020		2021		2022		2023		2024		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	80	35%	74	35%	61	27%	80	34%	67	31%	4,407	46%
Injection drug use (IDU) only	2	1%	6	3%	4	2%	1	<1%	2	1%	137	1%
Heterosexual contact only	80	35%	68	32%	82	37%	76	32%	77	36%	2,095	22%
Male-to-male sexual contact & IDU	12	5%	7	3%	5	2%	3	1%	6	3%	219	2%
IDU & Heterosexual contact	14	6%	13	6%	13	6%	7	3%	6	3%	362	4%
Male-to-male sexual contact & Heterosexual contact	25	11%	27	13%	20	9%	28	12%	26	12%	1,142	12%
Male-to-male sexual contact & IDU & Heterosexual contact	4	2%	2	1%	5	2%	5	2%	2	1%	133	1%
Perinatal exposure	1	<1%	-	-	2	1%	1	<1%	-	-	54	1%
Other/unknown	10	4%	15	7%	32	14%	33	14%	30	14%	993	10%
Total	228		212		224		234		216		9,542	

### Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
<b>Sex</b>						
Males	64	65%	28	65%	36	65%
Females	34	35%	15	35%	19	35%
<b>Age at AIDS diagnosis (years)</b>						
0-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	1	1%	1	2%	-	-
20-24	4	4%	-	-	4	7%
25-29	9	9%	5	12%	4	7%
30-34	19	19%	9	21%	10	18%
35-39	15	15%	5	12%	10	18%
40-44	16	16%	6	14%	10	18%
45-49	9	9%	-	-	9	16%
50-54	8	8%	6	14%	2	4%
55-64	13	13%	8	19%	5	9%
65+	4	4%	3	7%	1	2%
<b>Race/Ethnicity<sup>b</sup></b>						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	1	1%	-	-	1	2%
Black/African American	48	49%	20	47%	28	51%
Hispanic/Latino	16	16%	7	16%	9	16%
White	27	28%	11	26%	16	29%
Multi-Race	6	6%	5	12%	1	2%
<b>Race/Ethnicity<sup>b</sup> and Sex</b>						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	1	1%	-	-	1	2%
Black/African American Males	27	28%	11	26%	16	29%
Black/African American Females	21	21%	9	21%	12	22%
Hispanic/Latino Males	12	12%	5	12%	7	13%
Hispanic/Latina Females	4	4%	2	5%	2	4%
White Males	21	21%	8	19%	13	24%
White Females	6	6%	3	7%	3	5%
Multi-Race Males	4	4%	4	9%	-	-
Multi-Race Females	2	2%	1	2%	1	2%
<b>Total</b>	<b>98</b>		<b>43</b>		<b>55</b>	

**Notes:**

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

<sup>a</sup> Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category <sup>b</sup>	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	32	51%	16	59%	16	44%
Injection drug use (IDU)	3	5%	1	4%	2	6%
Male-to-male sexual contact and IDU	4	6%	2	7%	2	6%
Heterosexual contact	1	2%	-	-	1	3%
Other/unknown	23	37%	8	30%	15	42%
Subtotal	63	100%	27	100%	36	100%
<b>Female adult or adolescent</b>						
Injection drug use	4	12%	2	13%	2	11%
Heterosexual contact	25	74%	10	67%	15	79%
Other/unknown	5	15%	3	20%	2	11%
Subtotal	34	100%	15	100%	19	100%
<b>Child (&lt;13 years at diagnosis)</b>						
Perinatal	1	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-
<b>Total</b>	<b>98</b>		<b>43</b>		<b>55</b>	

## Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category <sup>c</sup>	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis <sup>a</sup>	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	21	21%	13	30%	8	15%
Injection drug use (IDU) only	2	2%	1	2%	1	2%
Heterosexual contact only	38	39%	12	28%	26	47%
Male-to-male sexual contact & IDU	2	2%	1	2%	1	2%
IDU & Heterosexual contact	5	5%	2	5%	3	5%
Male-to-male sexual contact & Heterosexual contact	11	11%	3	7%	8	15%
Male-to-male sexual contact & IDU & Heterosexual contact	2	2%	1	2%	1	2%
Perinatal exposure	1	1%	1	2%	-	-
Other/unknown	16	16%	9	21%	7	13%
<b>Total</b>	<b>98</b>		<b>43</b>		<b>55</b>	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

<sup>b</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>c</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.



# Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	81	80%	71	68%	62	72%	84	73%	64	65%	4,542	84%
Females	20	20%	34	32%	24	28%	31	27%	34	35%	856	16%
Age at AIDS diagnosis (years)												
0-9	-	-	-	-	-	-	-	-	-	-	24	<1%
10-14	1	1%	-	-	-	-	-	-	-	-	6	<1%
15-19	-	-	1	1%	1	1%	-	-	1	1%	49	1%
20-24	6	6%	6	6%	2	2%	9	8%	4	4%	325	6%
25-29	14	14%	8	8%	8	9%	11	10%	9	9%	687	13%
30-34	14	14%	20	19%	15	17%	25	22%	19	19%	1,085	20%
35-39	14	14%	16	15%	12	14%	20	17%	15	15%	986	18%
40-44	7	7%	17	16%	15	17%	9	8%	16	16%	857	16%
45-49	9	9%	10	10%	9	10%	12	10%	9	9%	581	11%
50-54	8	8%	9	9%	6	7%	11	10%	8	8%	392	7%
55-64	19	19%	14	13%	15	17%	16	14%	13	13%	339	6%
65+	9	9%	4	4%	3	3%	2	2%	4	4%	67	1%
Race/Ethnicity <sup>a</sup>												
American Indian/Alaska Native	-	-	1	1%	-	-	-	-	-	-	4	<1%
Asian/Pacific Islander	1	1%	-	-	1	1%	1	1%	1	1%	23	<1%
Black/African American	52	51%	54	51%	37	43%	65	57%	48	49%	1,884	35%
Hispanic/Latino	4	4%	3	3%	13	15%	14	12%	16	16%	234	4%
White	39	39%	44	42%	29	34%	31	27%	27	28%	3,062	57%
Multi-Race	5	5%	3	3%	6	7%	4	3%	6	6%	191	4%
Race/Ethnicity <sup>a</sup> and Sex												
American Indian/Alaska Native Males	-	-	1	1%	-	-	-	-	-	-	2	<1%
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	1	1%	-	-	1	1%	1	1%	-	-	19	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	1	1%	4	<1%
Black/African American Males	35	35%	30	29%	27	31%	46	40%	27	28%	1,371	25%
Black/African American Females	17	17%	24	23%	10	12%	19	17%	21	21%	513	10%
Hispanic/Latino Males	4	4%	2	2%	11	13%	12	10%	12	12%	194	4%
Hispanic/Latina Females	-	-	1	1%	2	2%	2	2%	4	4%	40	1%
White Males	38	38%	37	35%	19	22%	22	19%	21	21%	2,806	52%
White Females	1	1%	7	7%	10	12%	9	8%	6	6%	256	5%
Multi-Race Males	3	3%	1	1%	4	5%	3	3%	4	4%	150	3%
Multi-Race Females	2	2%	2	2%	2	2%	1	1%	2	2%	41	1%
Total	101		105		86		115		98		5,398	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

## Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category <sup>a</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	49	61%	43	61%	25	40%	40	48%	32	51%	3,185	70%
Injection drug use (IDU)	2	3%	5	7%	3	5%	2	2%	3	5%	202	4%
Male-to-male sexual contact and IDU	1	1%	2	3%	2	3%	4	5%	4	6%	183	4%
Heterosexual contact	7	9%	2	3%	2	3%	2	2%	1	2%	243	5%
Other/unknown	21	26%	19	27%	30	48%	35	42%	23	37%	711	16%
Subtotal	80	100%	71	100%	62	100%	83	100%	63	100%	4,524	100%
<b>Female adult or adolescent</b>												
Injection drug use	3	15%	4	12%	3	13%	3	10%	4	12%	129	15%
Heterosexual contact	15	75%	27	79%	15	65%	22	71%	25	74%	639	76%
Other/unknown	2	10%	3	9%	5	22%	6	19%	5	15%	70	8%
Subtotal	20	100%	34	100%	23	100%	31	100%	34	100%	838	100%
<b>Child (&lt;13 years at diagnosis)</b>												
Perinatal	1	100%	-	-	1	100%	1	100%	1	100%	30	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	6	17%
Subtotal	1	100%	-	-	1	100%	1	100%	1	100%	36	100%
<b>Total</b>	<b>101</b>		<b>105</b>		<b>86</b>		<b>115</b>		<b>98</b>		<b>5,398</b>	

## Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category <sup>b</sup>	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	42	42%	34	32%	20	23%	34	30%	21	21%	2,500	46%
Injection drug use (IDU) only	1	1%	2	2%	1	1%	-	-	2	2%	90	2%
Heterosexual contact only	31	31%	39	37%	34	40%	44	38%	38	39%	1,215	23%
Male-to-male sexual contact & IDU	1	1%	2	2%	2	2%	1	1%	2	2%	109	2%
IDU & Heterosexual contact	4	4%	7	7%	5	6%	5	4%	5	5%	241	4%
Male-to-male sexual contact & Heterosexual contact	7	7%	9	9%	5	6%	6	5%	11	11%	683	13%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	3	3%	2	2%	74	1%
Perinatal exposure	1	1%	-	-	1	1%	2	2%	1	1%	32	1%
Other/unknown	14	14%	12	11%	18	21%	20	17%	16	16%	454	8%
Total	101		105		86		115		98		5,398	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

# New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 11/Ryan White Part A-Columbus

---

**Counties:** Delaware, Fairfield, Franklin, Licking, Madison, Morrow,  
Pickaway, Union

## Reported persons living with diagnosed HIV infection in 2024 by current disease status and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	Living with diagnosed HIV infection in 2024			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Sex</b>							
Males	475.9	5,098	77%	2,976	77%	2,122	77%
Females	141.3	1,542	23%	910	23%	632	23%
<b>Age at end of year (years)</b>							
0-9	2.2	6	<1%	6	<1%	-	-
10-14	5.0	7	<1%	6	<1%	1	<1%
15-19	26.0	37	1%	30	1%	7	<1%
20-24	118.4	165	2%	145	4%	20	1%
25-29	235.9	374	6%	313	8%	61	2%
30-34	441.6	750	11%	558	14%	192	7%
35-39	508.0	832	13%	549	14%	283	10%
40-44	498.8	749	11%	460	12%	289	10%
45-49	497.1	658	10%	359	9%	299	11%
50-54	551.4	719	11%	358	9%	361	13%
55-64	635.7	1,545	23%	726	19%	819	30%
65+	247.8	798	12%	376	10%	422	15%
<b>Race/Ethnicity<sup>b</sup></b>							
American Indian/Alaska Native	181.0	7	<1%	5	<1%	2	<1%
Asian/Pacific Islander	45.5	60	1%	43	1%	17	1%
Black/African American	774.6	2,955	45%	1,730	45%	1,225	44%
Hispanic/Latino	440.1	605	9%	385	10%	220	8%
White	182.9	2,642	40%	1,516	39%	1,126	41%
Multi-Race	564.4	358	5%	194	5%	164	6%
Unknown	*	13	<1%	13	<1%	-	-
<b>Race/Ethnicity<sup>b</sup> and Sex</b>							
American Indian/Alaska Native Males	314.0	6	<1%	5	<1%	1	<1%
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Males	75.7	49	1%	35	1%	14	1%
Asian/Pacific Islander Females	16.4	11	<1%	8	<1%	3	<1%
Black/African American Males	1,024.1	1,914	29%	1,098	28%	816	30%
Black/African American Females	535.0	1,041	16%	632	16%	409	15%
Hispanic/Latino Males	714.6	508	8%	331	9%	177	6%
Hispanic/Latina Females	146.1	97	1%	54	1%	43	2%
White Males	325.5	2,328	35%	1,337	34%	991	36%
White Females	43.1	314	5%	179	5%	135	5%
Multi-Race Males	893.4	281	4%	158	4%	123	4%
Multi-Race Females	240.8	77	1%	36	1%	41	1%
Unknown	*	13	<1%	13	<1%	-	-
<b>Total</b>	<b>307.1</b>	<b>6,640</b>		<b>3,886</b>		<b>2,754</b>	

### Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

<sup>a</sup>The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

**Reported persons living with diagnosed HIV infection in 2024 by current disease status and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus**

Transmission Category <sup>a</sup>	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	3,472	69%	2,062	70%	1,410	67%
Injection drug use (IDU)	122	2%	59	2%	63	3%
Male-to-male sexual contact and IDU	208	4%	114	4%	94	4%
Heterosexual contact	258	5%	95	3%	163	8%
Other/unknown	1,001	20%	622	21%	379	18%
Subtotal	5,061	100%	2,952	100%	2,109	100%
<b>Female adult or adolescent</b>						
Injection drug use	119	8%	55	6%	64	10%
Heterosexual contact	1,180	79%	676	76%	504	82%
Other/unknown	200	13%	154	17%	46	7%
Subtotal	1,499	100%	885	100%	614	100%
<b>Child (&lt;13 years at diagnosis)</b>						
Perinatal	66	83%	36	73%	30	97%
Other/unknown	14	18%	13	27%	1	3%
Subtotal	80	100%	49	100%	31	100%
<b>Total</b>	<b>6,640</b>		<b>3,886</b>		<b>2,754</b>	

**Reported persons living with diagnosed HIV infection in 2024 by current disease status and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus**

Exposure Category <sup>b</sup>	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,763	42%	1,662	43%	1,101	40%
Injection drug use (IDU) only	44	1%	22	1%	22	1%
Heterosexual contact only	1,881	28%	1,002	26%	879	32%
Male-to-male sexual contact & IDU	123	2%	66	2%	57	2%
IDU & Heterosexual contact	197	3%	92	2%	105	4%
Male-to-male sexual contact & Heterosexual contact	707	11%	400	10%	307	11%
Male-to-male sexual contact & IDU & Heterosexual contact	85	1%	48	1%	37	1%
Perinatal exposure	73	1%	41	1%	32	1%
Other/unknown	767	12%	553	14%	214	8%
<b>Total</b>	<b>6,640</b>		<b>3,886</b>		<b>2,754</b>	

Notes:  
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.  
 Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.  
  
<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and age, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2024																				
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino <sup>a</sup>			White			Multi-Race			Unknown	
	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	Rate <sup>b</sup>	No.	%	No.	%
0-9	*	-	-	*	-	-	*	3	<1%	*	2	<1%	*	1	<1%	*	-	-	-	-
10-14	*	-	-	*	1	2%	16.6	5	<1%	*	1	<1%	*	-	-	*	-	-	-	-
15-19	*	-	-	*	2	3%	91.7	26	1%	38.3	5	1%	*	2	<1%	*	2	1%	-	-
20-24	*	1	14%	*	2	3%	420.5	109	4%	211.7	26	4%	22.2	19	1%	134.5	8	2%	-	-
25-29	*	1	14%	*	3	5%	760.6	219	7%	499.6	59	10%	69.0	69	3%	403.9	23	6%	-	-
30-34	*	1	14%	111.4	14	23%	1,124.9	382	13%	809.2	92	15%	201.1	215	8%	976.2	46	13%	-	-
35-39	*	-	-	*	3	5%	1,481.0	435	15%	1,010.6	102	17%	226.3	242	9%	1,457.3	50	14%	-	-
40-44	*	1	14%	42.4	5	8%	1,295.7	330	11%	856.7	84	14%	281.5	282	11%	1,731.1	47	13%	-	-
45-49	*	-	-	123.1	11	18%	1,396.6	312	11%	718.4	61	10%	270.3	244	9%	1,426.5	30	8%	-	-
50-54	*	1	14%	128.6	10	17%	1,407.8	307	10%	723.3	47	8%	339.3	313	12%	2,252.7	41	11%	-	-
55-64	*	2	29%	67.5	7	12%	1,409.3	520	18%	1,013.1	86	14%	462.4	852	32%	2,770.1	70	20%	8	62%
65+	*	-	-	*	2	3%	792.9	307	10%	671.9	40	7%	152.7	403	15%	1,495.3	41	11%	5	38%
Total	181.0	7		45.5	60		774.6	2,955		440.1	605		182.9	2,642		564.4	358		13	

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2024														
Transmission Category <sup>c</sup>	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	4	67%	19	40%	1,114	59%	340	68%	1,794	77%	200	72%	1	8%
Injection drug use (IDU)	-	-	-	-	45	2%	6	1%	61	3%	10	4%	-	-
Male-to-male sexual contact and IDU	-	-	3	6%	54	3%	11	2%	121	5%	19	7%	-	-
Heterosexual contact	1	17%	3	6%	158	8%	29	6%	56	2%	11	4%	-	-
Other/unknown	1	17%	22	47%	522	28%	115	23%	291	13%	39	14%	11	92%
Subtotal	6	100%	47	100%	1,893	100%	501	100%	2,323	100%	279	100%	12	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	44	4%	9	10%	56	18%	10	13%	-	-
Heterosexual contact	1	100%	5	56%	841	83%	64	70%	215	70%	54	72%	-	-
Other/unknown	-	-	4	44%	129	13%	19	21%	36	12%	11	15%	1	100%
Subtotal	1	100%	9	100%	1,014	100%	92	100%	307	100%	75	100%	1	100%
Child (<13 years at diagnosis)														
Perinatal	-	-	4	100%	39	81%	9	75%	10	83%	4	100%	-	-
Other/unknown	-	-	-	-	9	19%	3	25%	2	17%	-	-	-	-
Subtotal	-	-	4	100%	48	100%	12	100%	12	100%	4	100%	-	-
Total	7		60		2,955		605		2,642		358		13	

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2024														
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino <sup>a</sup>		White		Multi-Race		Unknown	
Exposure Category <sup>d</sup>	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	43%	15	25%	789	27%	279	46%	1,522	58%	154	43%	1	8%
Injection drug use (IDU) only	-	-	-	-	15	1%	4	1%	23	1%	2	1%	-	-
Heterosexual contact only	2	29%	14	23%	1,287	44%	136	22%	349	13%	93	26%	-	-
Male-to-male sexual contact & IDU	-	-	2	3%	29	1%	7	1%	76	3%	9	3%	-	-
IDU & Heterosexual contact	-	-	-	-	74	3%	11	2%	94	4%	18	5%	-	-
Male-to-male sexual contact & Heterosexual contact	1	14%	4	7%	325	11%	61	10%	270	10%	46	13%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	2%	25	1%	4	1%	45	2%	10	3%	-	-
Perinatal exposure	-	-	5	8%	44	1%	10	2%	10	0%	4	1%	-	-
Other/unknown	1	14%	19	32%	367	12%	93	15%	253	10%	22	6%	12	92%
Total	7		60		2,955		605		2,642		358		13	

Notes:  
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.  
Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.  
<sup>b</sup> The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.  
<sup>c</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.  
<sup>d</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus, 2020-2024

Living with Diagnosed HIV Infection															
Characteristic	2020			2021			2022			2023			2024		
	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%	Rate <sup>a</sup>	No.	%
Sex															
Males	467.8	4,778	79%	468.2	4,840	79%	467.6	4,868	78%	472.5	4,954	78%	475.9	5,098	77%
Females	120.4	1,268	21%	124.5	1,312	21%	127.7	1,350	22%	132.6	1,417	22%	141.3	1,542	23%
Age at end of year (years)															
0-9	3.0	8	<1%	2.6	7	<1%	3.0	8	<1%	3.0	8	<1%	2.2	6	<1%
10-14	7.5	10	<1%	5.1	7	<1%	4.4	6	<1%	4.4	6	<1%	5.0	7	<1%
15-19	23.2	31	1%	29.8	41	1%	26.3	36	1%	20.0	28	<1%	26.0	37	1%
20-24	112.1	149	2%	113.0	150	2%	103.2	144	2%	127.2	172	3%	118.4	165	2%
25-29	285.3	463	8%	255.4	398	6%	240.8	372	6%	237.7	365	6%	235.9	374	6%
30-34	436.2	723	12%	447.2	743	12%	449.2	745	12%	464.0	772	12%	441.6	750	11%
35-39	415.2	620	10%	420.8	643	10%	435.9	672	11%	459.2	724	11%	508.0	832	13%
40-44	467.2	629	10%	476.7	669	11%	489.6	700	11%	491.7	719	11%	498.8	749	11%
45-49	522.8	675	11%	496.4	631	10%	490.2	619	10%	478.9	615	10%	497.1	658	10%
50-54	669.5	851	14%	629.9	828	13%	582.7	768	12%	562.8	738	12%	551.4	719	11%
55-64	575.6	1,424	24%	599.0	1,487	24%	623.4	1,522	24%	627.1	1,522	24%	635.7	1,545	23%
65+	160.9	463	8%	190.1	548	9%	207.9	626	10%	226.5	702	11%	247.8	798	12%
Race/Ethnicity <sup>b</sup>															
American Indian/Alaska Native	126.9	5	<1%	125.4	5	<1%	128.5	5	<1%	154.5	6	<1%	181.0	7	<1%
Asian/Pacific Islander	40.5	44	1%	42.1	47	1%	40.4	47	1%	41.8	50	1%	45.5	60	1%
Black/African American	726.3	2,520	42%	741.2	2,610	42%	742.2	2,675	43%	757.7	2,788	44%	774.6	2,955	45%
Hispanic/Latino	390.7	380	6%	402.3	407	7%	436.9	461	7%	427.1	527	8%	440.1	605	9%
White	186.9	2,731	45%	185.8	2,713	44%	183.0	2,657	43%	182.6	2,631	41%	182.9	2,642	40%
Multi-Race	624.2	353	6%	610.8	357	6%	600.1	360	6%	578.2	356	6%	564.4	358	5%
Unknown	*	13	<1%	*	13	<1%	*	13	<1%	*	13	<1%	*	13	<1%
Race/Ethnicity <sup>b</sup> and Sex															
American Indian/Alaska Native Males	*	4	<1%	*	4	<1%	*	4	<1%	262.5	5	<1%	314.0	6	<1%
American Indian/Alaska Native Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander Males	67.7	36	1%	69.2	38	1%	66.3	38	1%	71.6	42	1%	75.7	49	1%
Asian/Pacific Islander Females	14.4	8	<1%	15.9	9	<1%	15.2	9	<1%	13.1	8	<1%	16.4	11	<1%
Black/African American Males	1,012.6	1,705	28%	1,015.5	1,747	28%	1,006.4	1,777	29%	1,020.9	1,840	29%	1,024.1	1,914	29%
Black/African American Females	456.4	815	13%	479.1	863	14%	488.4	898	14%	505.0	948	15%	535.0	1,041	16%
Hispanic/Latino Males	606.0	304	5%	632.3	331	5%	697.2	381	6%	701.6	447	7%	714.6	508	8%
Hispanic/Latina Females	161.4	76	1%	155.7	76	1%	157.3	80	1%	134.0	80	1%	146.1	97	1%
White Males	338.3	2,435	40%	334.9	2,423	39%	328.8	2,370	38%	326.6	2,330	37%	325.5	2,328	35%
White Females	39.9	296	5%	39.4	290	5%	39.3	287	5%	41.4	301	5%	43.1	314	5%
Multi-Race Males	1,010.4	282	5%	984.3	285	5%	960.7	286	5%	913.1	278	4%	893.4	281	4%
Multi-Race Females	247.9	71	1%	244.1	72	1%	244.9	74	1%	250.6	78	1%	240.8	77	1%
Unknown	*	13	<1%	*	13	<1%	*	13	<1%	*	13	<1%	*	13	<1%
Total	291.4	6,046		294.7	6,152		296.4	6,218		300.9	6,371		307.1	6,640	

Notes:  
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (\*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.  
<sup>a</sup> The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.  
<sup>b</sup> Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

**Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus, 2020-2024**

Living with Diagnosed HIV Infection										
Transmission Category <sup>a</sup>	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>										
Male-to-male sexual contact	3,403	72%	3,429	71%	3,412	71%	3,421	70%	3,472	69%
Injection drug use (IDU)	106	2%	117	2%	118	2%	120	2%	122	2%
Male-to-male sexual contact and IDU	223	5%	222	5%	220	5%	204	4%	208	4%
Heterosexual contact	267	6%	263	5%	260	5%	251	5%	258	5%
Other/unknown	752	16%	781	16%	828	17%	922	19%	1,001	20%
Subtotal	4,751	100%	4,812	100%	4,838	100%	4,918	100%	5,061	100%
<b>Female adult or adolescent</b>										
Injection drug use	123	10%	125	10%	121	9%	113	8%	119	8%
Heterosexual contact	997	81%	1,029	81%	1,058	81%	1,105	80%	1,180	79%
Other/unknown	110	9%	118	9%	130	10%	159	12%	200	13%
Subtotal	1,230	100%	1,272	100%	1,309	100%	1,377	100%	1,499	100%
<b>Child (&lt;13 years at diagnosis)</b>										
Perinatal	59	91%	61	90%	64	90%	65	86%	66	83%
Other/unknown	6	9%	7	10%	7	10%	11	14%	14	18%
Subtotal	65	100%	68	100%	71	100%	76	100%	80	100%
<b>Total</b>	<b>6,046</b>		<b>6,152</b>		<b>6,218</b>		<b>6,371</b>		<b>6,640</b>	

**Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus, 2020-2024**

Living with Diagnosed HIV Infection										
Exposure Category <sup>b</sup>	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,724	45%	2,747	45%	2,734	44%	2,731	43%	2,763	42%
Injection drug use (IDU) only	34	1%	40	1%	41	1%	37	1%	44	1%
Heterosexual contact only	1,603	27%	1,657	27%	1,702	27%	1,770	28%	1,881	28%
Male-to-male sexual contact & IDU	138	2%	139	2%	137	2%	125	2%	123	2%
IDU & Heterosexual contact	195	3%	202	3%	198	3%	196	3%	197	3%
Male-to-male sexual contact & Heterosexual contact	677	11%	680	11%	676	11%	688	11%	707	11%
Male-to-male sexual contact & IDU & Heterosexual contact	85	1%	83	1%	83	1%	79	1%	85	1%
Perinatal exposure	62	1%	64	1%	69	1%	72	1%	73	1%
Other/unknown	528	9%	540	9%	578	9%	673	11%	767	12%
Total	6,046		6,152		6,218		6,371		6,640	

Notes:  
 Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.  
 Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.  
<sup>b</sup> Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.  
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.