

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 11/Ryan White Part A- Columbus

Counties: Delaware, Fairfield, Franklin, Licking, Madison, Morrow, Pickaway, Union

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2021

Reported new diagnoses of HIV infection in 2020 by disease status and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	2020 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	16.8	172	75%	133	74%	31	76%	8	80%
Females	5.5	58	25%	46	26%	10	24%	2	20%
Age at diagnosis (yr)									
<13	*	1	<1%	1	1%	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	6.7	9	4%	9	5%	-	-	-	-
20-24	27.1	36	16%	33	18%	2	5%	1	10%
25-29	27.1	44	19%	38	21%	6	15%	-	-
30-34	26.5	44	19%	34	19%	8	20%	2	20%
35-39	18.1	27	12%	20	11%	6	15%	1	10%
40-44	11.9	16	7%	12	7%	3	7%	1	10%
45-49	13.2	17	7%	12	7%	5	12%	-	-
50-54	11.0	14	6%	7	4%	3	7%	4	40%
55-64	8.1	20	9%	11	6%	8	20%	1	10%
65+	*	2	1%	2	1%	-	-	-	-
Race/Ethnicity^b									
American Indian/Alaska Native	*	1	<1%	1	1%	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
Black/African-American	38.0	132	57%	106	59%	19	46%	7	70%
Hispanic/Latinx	7.2	7	3%	7	4%	-	-	-	-
White	5.7	84	37%	62	35%	19	46%	3	30%
Multi-Race	10.6	6	3%	3	2%	3	7%	-	-
Race/Ethnicity^b and Sex at birth									
American Indian/Alaska Native Males	*	1	<1%	1	1%	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African-American Males	52.9	89	39%	72	40%	12	29%	5	50%
Black/African-American Females	24.1	43	19%	34	19%	7	17%	2	20%
Hispanic/Latino Males	14.0	7	3%	7	4%	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
White Males	10.0	72	31%	52	29%	17	41%	3	30%
White Females	1.6	12	5%	10	6%	2	5%	-	-
Multi-Race Males	*	3	1%	1	1%	2	5%	-	-
Multi-Race Females	*	3	1%	2	1%	1	2%	-	-
Total	11.1	230		179		41		10	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^bHispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by disease status and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^a	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	104	61%	83	63%	19	61%	2	25%
Injection drug use (IDU)	9	5%	6	5%	2	6%	1	13%
Male-to-male sexual contact and IDU	16	9%	15	11%	-	-	1	13%
Heterosexual contact	9	5%	6	5%	2	6%	1	13%
Other/unknown	33	19%	22	17%	8	26%	3	38%
Subtotal	171	100%	132	100%	31	100%	8	100%
Female adult or adolescent								
Injection drug use	7	12%	5	11%	2	20%	-	-
Heterosexual contact	49	84%	39	85%	8	80%	2	100%
Other/unknown	2	3%	2	4%	-	-	-	-
Subtotal	58	100%	46	100%	10	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	1	100%	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Total	230		179		41		10	

Reported new diagnoses of HIV infection in 2020 by disease status and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^b	2020 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	100	43%	80	45%	18	44%	2	20%
Injection drug use (IDU) only	9	4%	5	3%	3	7%	1	10%
Heterosexual contact only	58	25%	45	25%	10	24%	3	30%
Male-to-male sexual contact & IDU	13	6%	12	7%	-	-	1	10%
IDU & Heterosexual contact	7	3%	6	3%	1	2%	-	-
Male-to-male sexual contact & Heterosexual contact	4	2%	3	2%	1	2%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	3	1%	3	2%	-	-	-	-
Perinatal exposure	1	<1%	1	1%	-	-	-	-
Other/unknown	35	15%	24	13%	8	20%	3	30%
Total	230		179		41		10	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

2020 diagnosis of HIV infection

Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	1	1%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	28.4	7	5%	*	1	14%	*	1	1%	*	1	17%
20-24	*	-	-	*	-	-	115.5	28	21%	*	1	14%	6.8	6	7%	*	1	17%
25-29	*	-	-	*	-	-	76.4	23	17%	*	2	29%	14.7	16	19%	*	3	50%
30-34	*	-	-	*	-	-	102.1	29	22%	*	1	14%	11.5	13	15%	*	1	17%
35-39	*	-	-	*	-	-	56.7	14	11%	*	-	-	11.7	12	14%	*	1	17%
40-44	*	-	-	*	-	-	26.6	6	5%	*	1	14%	9.5	9	11%	*	-	-
45-49	*	1	100%	*	-	-	37.8	8	6%	*	1	14%	7.6	7	8%	*	-	-
50-54	*	-	-	*	-	-	46.4	9	7%	*	-	-	5.2	5	6%	*	-	-
55-64	*	-	-	*	-	-	16.8	6	5%	*	-	-	7.2	14	17%	*	-	-
65+	*	-	-	*	-	-	*	1	1%	*	-	-	*	1	1%	*	-	-
Total	*	1		*	-		38.0	132		7.2	7		5.7	84		10.6	6	

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

2020 diagnosis of HIV infection

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	56	64%	6	86%	40	56%	2	67%
Injection drug use (IDU)	-	-	-	-	1	1%	-	-	7	10%	1	33%
Male-to-male sexual contact and IDU	-	-	-	-	2	2%	-	-	14	19%	-	-
Heterosexual contact	-	-	-	-	7	8%	-	-	2	3%	-	-
Other/unknown	1	100%	-	-	22	25%	1	14%	9	13%	-	-
Subtotal	1	100%	-	-	88	100%	7	100%	72	100%	3	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	2	5%	-	-	4	33%	1	33%
Heterosexual contact	-	-	-	-	41	95%	-	-	6	50%	2	67%
Other/unknown	-	-	-	-	-	-	-	-	2	17%	-	-
Subtotal	-	-	-	-	43	100%	-	-	12	100%	3	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	1	100%	-	-	-	-	-	-
Total	1		-		132		7		84		6	

Reported new diagnoses of HIV infection in 2020 by race/ethnicity and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

2020 diagnosis of HIV infection

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	54	41%	6	86%	39	46%	1	17%
Injection drug use (IDU) only	-	-	-	-	2	2%	-	-	6	7%	1	17%
Heterosexual contact only	-	-	-	-	48	36%	-	-	8	10%	2	33%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	13	15%	-	-
IDU & Heterosexual contact	-	-	-	-	1	1%	-	-	5	6%	1	17%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	2	2%	-	-	1	1%	1	17%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	2	2%	-	-	1	1%	-	-
Perinatal exposure	-	-	-	-	1	1%	-	-	-	-	-	-
Other/unknown	1	100%	-	-	22	17%	1	14%	11	13%	-	-
Total	1		-		132		7		84		6	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2016			2017			2018			2019			2020			No.	%
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%		
Sex at birth																	
Males	18.2	177	80%	19.6	194	80%	16.8	169	76%	18.6	188	81%	16.8	172	75%	7,330	84%
Females	4.3	43	20%	4.9	50	20%	5.2	54	24%	4.3	45	19%	5.5	58	25%	1,391	16%
Age at diagnosis (yr)																	
<13	*	2	1%	*	-	-	*	3	1%	*	-	-	*	1	<1%	60	1%
13-14	*	1	<1%	*	-	-	*	-	-	*	-	-	*	-	-	10	<1%
15-19	10.8	14	6%	9.0	12	5%	9.0	12	5%	7.5	10	4%	6.7	9	4%	316	4%
20-24	29.0	38	17%	27.1	36	15%	23.2	31	14%	33.1	44	19%	27.1	36	16%	1,137	13%
25-29	24.4	40	18%	34.3	57	23%	32.2	54	24%	29.2	48	21%	27.1	44	19%	1,551	18%
30-34	16.9	25	11%	23.1	35	14%	23.5	37	17%	22.2	36	15%	26.5	44	19%	1,656	19%
35-39	19.6	27	12%	19.0	27	11%	14.5	21	9%	12.2	18	8%	18.1	27	12%	1,413	16%
40-44	16.5	21	10%	14.9	19	8%	15.5	20	9%	16.0	21	9%	11.9	16	7%	1,040	12%
45-49	11.3	15	7%	11.9	16	7%	13.5	18	8%	9.9	13	6%	13.2	17	7%	697	8%
50-54	15.4	20	9%	12.5	16	7%	8.7	11	5%	14.3	18	8%	11.0	14	6%	426	5%
55-64	6.7	16	7%	9.1	22	9%	4.9	12	5%	7.7	19	8%	8.1	20	9%	338	4%
65+	*	1	<1%	*	4	2%	*	4	2%	2.2	6	3%	*	2	1%	77	1%
Race/Ethnicity^b																	
American Indian/Alaska Native	*	-	-	*	1	<1%	*	-	-	*	-	-	*	1	<1%	8	<1%
Asian/Pacific Islander	8.4	7	3%	*	3	1%	*	2	1%	4.9	5	2%	*	-	-	57	1%
Black/African-American	35.1	110	50%	36.3	118	48%	36.2	121	54%	35.3	120	52%	38.0	132	57%	3,178	36%
Hispanic/Latinx	15.9	13	6%	16.3	14	6%	16.5	15	7%	19.4	18	8%	7.2	7	3%	343	4%
White	5.5	80	36%	6.8	99	41%	5.4	79	35%	5.5	81	35%	5.7	84	37%	4,839	55%
Multi-Race	20.4	10	5%	17.7	9	4%	11.4	6	3%	16.7	9	4%	10.6	6	3%	273	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	23	<1%
Race/Ethnicity^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	1	<1%	*	-	-	*	-	-	*	1	<1%	6	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Asian/Pacific Islander Males	14.6	6	3%	*	1	<1%	*	1	<1%	9.9	5	2%	*	-	-	42	<1%
Asian/Pacific Islander Females	*	1	<1%	*	2	1%	*	1	<1%	*	-	-	*	-	-	15	<1%
Black/African-American Males	56.2	85	39%	55.3	87	36%	56.2	91	41%	58.8	97	42%	52.9	89	39%	2,327	27%
Black/African-American Females	15.4	25	11%	18.5	31	13%	17.5	30	13%	13.1	23	10%	24.1	43	19%	851	10%
Hispanic/Latino Males	28.4	12	5%	29.2	13	5%	27.5	13	6%	33.3	16	7%	14.0	7	3%	293	3%
Hispanic/Latina Females	*	1	<1%	*	1	<1%	*	2	1%	*	2	1%	*	-	-	50	1%
White Males	9.3	66	30%	11.8	85	35%	8.5	61	27%	8.9	64	27%	10.0	72	31%	4,418	51%
White Females	1.9	14	6%	1.9	14	6%	2.4	18	8%	2.3	17	7%	1.6	12	5%	421	5%
Multi-Race Males	33.3	8	4%	28.0	7	3%	*	3	1%	22.6	6	3%	*	3	1%	225	3%
Multi-Race Females	*	2	1%	*	2	1%	*	3	1%	*	3	1%	*	3	1%	48	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	23	<1%
Total	11.1	220		12.1	244		10.9	223		11.3	233		11.1	230		8,721	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^a	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	117	66%	120	62%	110	65%	112	60%	104	61%	5,191	71%
Injection drug use (IDU)	7	4%	9	5%	6	4%	3	2%	9	5%	273	4%
Male-to-male sexual contact and IDU	8	5%	9	5%	10	6%	12	6%	16	9%	309	4%
Heterosexual contact	6	3%	5	3%	6	4%	12	6%	9	5%	313	4%
Other/unknown	38	22%	51	26%	36	21%	49	26%	33	19%	1,215	17%
Subtotal	176	100%	194	100%	168	100%	188	100%	171	100%	7,301	100%
Female adult or adolescent												
Injection drug use	6	14%	9	18%	12	23%	9	20%	7	12%	175	13%
Heterosexual contact	34	81%	40	80%	37	71%	34	76%	49	84%	1,040	77%
Other/unknown	2	5%	1	2%	3	6%	2	4%	2	3%	142	10%
Subtotal	42	100%	50	100%	52	100%	45	100%	58	100%	1,357	100%
Child (<13 yrs at diagnosis)												
Perinatal	2	100%	-	-	3	100%	-	-	1	100%	52	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	11	17%
Subtotal	2	100%	-	-	3	100%	-	-	1	100%	63	100%
Total	220		244		223		233		230		8,721	

Trends in reported diagnoses of HIV infection by year of diagnosis (2016-2020) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^b	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sexual contact only	117	53%	118	48%	107	48%	108	46%	100	43%	4,995	57%
Injection drug use (IDU) only	6	3%	10	4%	7	3%	4	2%	9	4%	271	3%
Heterosexual contact only	40	18%	45	18%	43	19%	46	20%	58	25%	1,353	16%
Male-to-male sexual contact & IDU	7	3%	7	3%	10	4%	11	5%	13	6%	265	3%
IDU & Heterosexual contact	7	3%	8	3%	11	5%	8	3%	7	3%	177	2%
Male-to-male sexual contact & Heterosexual contact	-	-	2	1%	3	1%	4	2%	4	2%	196	2%
Male-to-male sexual contact & IDU & Heterosexual contact	1	<1%	2	1%	-	-	1	<1%	3	1%	44	1%
Perinatal exposure	3	1%	-	-	3	1%	-	-	1	<1%	55	1%
Other/unknown	39	18%	52	21%	39	17%	51	22%	35	15%	1,365	16%
Total	220		244		223		233		230		8,721	

Notes:
 Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2016-2020) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2020.
 Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex at birth						
Males	72	81%	31	82%	41	80%
Females	17	19%	7	18%	10	20%
Age at AIDS diagnosis (yr)						
<13	-	-	-	-	-	-
13-14	1	1%	1	3%	-	-
15-19	-	-	-	-	-	-
20-24	6	7%	2	5%	4	8%
25-29	11	12%	4	11%	7	14%
30-34	11	12%	3	8%	8	16%
35-39	11	12%	4	11%	7	14%
40-44	6	7%	2	5%	4	8%
45-49	9	10%	4	11%	5	10%
50-54	8	9%	1	3%	7	14%
55-64	19	21%	10	26%	9	18%
65+	7	8%	7	18%	-	-
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	1	1%	1	3%	-	-
Black/African-American	44	49%	19	50%	25	49%
Hispanic/Latinx	2	2%	-	-	2	4%
White	37	42%	16	42%	21	41%
Multi-Race	5	6%	2	5%	3	6%
Race/Ethnicity^b and Sex at birth						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	1	1%	1	3%	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African-American Males	30	34%	13	34%	17	33%
Black/African-American Females	14	16%	6	16%	8	16%
Hispanic/Latino Males	2	2%	-	-	2	4%
Hispanic/Latina Females	-	-	-	-	-	-
White Males	36	40%	16	42%	20	39%
White Females	1	1%	-	-	1	2%
Multi-Race Males	3	3%	1	3%	2	4%
Multi-Race Females	2	2%	1	3%	1	2%
Total	89		38		51	

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^b	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	41	58%	19	63%	22	54%
Injection drug use (IDU)	2	3%	-	-	2	5%
Male-to-male sexual contact and IDU	1	1%	-	-	1	2%
Heterosexual contact	7	10%	3	10%	4	10%
Other/unknown	20	28%	8	27%	12	29%
Subtotal	71	100%	30	100%	41	100%
Female adult or adolescent						
Injection drug use	3	18%	2	29%	1	10%
Heterosexual contact	12	71%	3	43%	9	90%
Other/unknown	2	12%	2	29%	-	-
Subtotal	17	100%	7	100%	10	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	6%	1	14%	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	1	6%	1	14%	-	-
Total	89		38		51	

Reported diagnoses of AIDS in 2020 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^c	AIDS Diagnoses					
	2020 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	40	45%	19	50%	21	41%
Injection drug use (IDU) only	3	3%	1	3%	2	4%
Heterosexual contact only	19	21%	6	16%	13	25%
Male-to-male sexual contact & IDU	1	1%	-	-	1	2%
IDU & Heterosexual contact	2	2%	1	3%	1	2%
Male-to-male sexual contact & Heterosexual contact	1	1%	-	-	1	2%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	1	1%	1	3%	-	-
Other/unknown	22	25%	10	26%	12	24%
Total	89		38		51	

Notes:

Data reflects all persons with an AIDS diagnosis in 2020 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex at birth												
Males	91	82%	82	77%	73	75%	77	73%	72	81%	4,109	86%
Females	20	18%	24	23%	24	25%	29	27%	17	19%	686	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	1	1%	-	-	-	-	24	1%
13-14	-	-	-	-	-	-	-	-	1	1%	5	<1%
15-19	1	1%	1	1%	3	3%	1	1%	-	-	44	1%
20-24	9	8%	12	11%	8	8%	9	8%	6	7%	286	6%
25-29	16	14%	7	7%	8	8%	18	17%	11	12%	626	13%
30-34	12	11%	12	11%	19	20%	23	22%	11	12%	979	20%
35-39	19	17%	15	14%	14	14%	11	10%	11	12%	878	18%
40-44	14	13%	19	18%	12	12%	12	11%	6	7%	775	16%
45-49	9	8%	10	9%	10	10%	5	5%	9	10%	518	11%
50-54	18	16%	12	11%	10	10%	12	11%	8	9%	342	7%
55-64	10	9%	16	15%	11	11%	12	11%	19	21%	270	6%
65+	3	3%	2	2%	1	1%	3	3%	7	8%	48	1%
Race/Ethnicity^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander	1	1%	-	-	-	-	2	2%	1	1%	20	<1%
Black/African-American	50	45%	55	52%	49	51%	50	47%	44	49%	1,585	33%
Hispanic/Latinx	9	8%	7	7%	6	6%	7	7%	2	2%	174	4%
White	49	44%	39	37%	40	41%	45	42%	37	42%	2,868	60%
Multi-Race	2	2%	5	5%	2	2%	2	2%	5	6%	146	3%
Race/Ethnicity^a and Sex at birth												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	1	1%	-	-	-	-	2	2%	1	1%	18	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Black/African-American Males	37	33%	40	38%	38	39%	35	33%	30	34%	1,172	24%
Black/African-American Females	13	12%	15	14%	11	11%	15	14%	14	16%	413	9%
Hispanic/Latino Males	8	7%	5	5%	6	6%	6	6%	2	2%	149	3%
Hispanic/Latina Females	1	1%	2	2%	-	-	1	1%	-	-	25	1%
White Males	43	39%	33	31%	28	29%	34	32%	36	40%	2,648	55%
White Females	6	5%	6	6%	12	12%	11	10%	1	1%	220	5%
Multi-Race Males	2	2%	4	4%	1	1%	-	-	3	3%	122	3%
Multi-Race Females	-	-	1	1%	1	1%	2	2%	2	2%	24	1%
Total	111		106		97		106		89		4,795	

Notes:

AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent												
Male-to-male sexual contact	59	65%	50	61%	37	51%	38	49%	41	58%	2,933	72%
Injection drug use (IDU)	4	4%	4	5%	6	8%	4	5%	2	3%	180	4%
Male-to-male sexual contact and IDU	2	2%	7	9%	3	4%	4	5%	1	1%	161	4%
Heterosexual contact	5	5%	1	1%	5	7%	3	4%	7	10%	225	5%
Other/unknown	21	23%	20	24%	21	29%	28	36%	20	28%	594	15%
Subtotal	91	100%	82	100%	72	100%	77	100%	71	100%	4,093	100%
Female adult or adolescent												
Injection drug use	3	15%	5	21%	12	50%	7	24%	3	18%	108	16%
Heterosexual contact	17	85%	16	67%	10	42%	20	69%	12	71%	513	77%
Other/unknown	-	-	3	13%	2	8%	2	7%	2	12%	48	7%
Subtotal	20	100%	24	100%	24	100%	29	100%	17	100%	669	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	1	100%	-	-	1	100%	26	79%
Other/unknown	-	-	-	-	-	-	-	-	-	-	7	21%
Subtotal	-	-	-	-	1	100%	-	-	1	100%	33	100%
Total	111		106		97		106		89		4,795	

Trends in reported diagnoses of AIDS by year of diagnosis (2016-2020) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2016		2017		2018		2019		2020		No.	%
	No.	%	No.	%	No.	%	No.	%	No.	%		
Male-to-male sexual contact only	57	51%	50	47%	35	36%	38	36%	40	45%	2,824	59%
Injection drug use (IDU) only	4	4%	4	4%	5	5%	6	6%	3	3%	179	4%
Heterosexual contact only	22	20%	17	16%	15	15%	23	22%	19	21%	738	15%
Male-to-male sexual contact & IDU	2	2%	5	5%	3	3%	3	3%	1	1%	137	3%
IDU & Heterosexual contact	3	3%	5	5%	13	13%	5	5%	2	2%	109	2%
Male-to-male sexual contact & Heterosexual contact	2	2%	-	-	2	2%	-	-	1	1%	109	2%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	2	2%	-	-	1	1%	-	-	24	1%
Perinatal exposure	-	-	-	-	1	1%	-	-	1	1%	27	1%
Other/unknown	21	19%	23	22%	23	24%	30	28%	22	25%	648	14%
Total	111		106		97		106		89		4,795	

Notes:

AIDS diagnoses by year (2016-2020) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2020.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode or transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Counties: Delaware, Fairfield, Franklin, Licking, Madison, Morrow, Pickaway, Union

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	Living with diagnosed HIV infection			Current Disease Status			
	in 2020			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex at birth							
Males	478.7	4,889	79%	2,708	79%	2,181	79%
Females	123.0	1,296	21%	724	21%	572	21%
Age at end of year							
<13	3.7	13	<1%	12	<1%	1	<1%
13-14	11.3	6	<1%	2	<1%	4	<1%
15-19	23.2	31	1%	25	1%	6	<1%
20-24	109.8	146	2%	124	4%	22	1%
25-29	291.5	473	8%	366	11%	107	4%
30-34	445.3	738	12%	515	15%	223	8%
35-39	417.2	623	10%	394	11%	229	8%
40-44	479.1	645	10%	344	10%	301	11%
45-49	533.7	689	11%	340	10%	349	13%
50-54	679.0	863	14%	427	12%	436	16%
55-64	595.0	1,472	24%	664	19%	808	29%
65+	168.9	486	8%	219	6%	267	10%
Race/Ethnicity^b							
American Indian/Alaska Native	203.0	8	<1%	7	<1%	1	<1%
Asian/Pacific Islander	52.5	57	1%	38	1%	19	1%
Black/African-American	756.0	2,623	42%	1,465	43%	1,158	42%
Hispanic/Latinx	395.9	385	6%	211	6%	174	6%
White	191.8	2,803	45%	1,554	45%	1,249	45%
Multi-Race	509.3	288	5%	136	4%	152	6%
Unknown	*	21	<1%	21	1%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	357.0	7	<1%	7	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Males	82.7	44	1%	28	1%	16	1%
Asian/Pacific Islander Females	23.4	13	<1%	10	<1%	3	<1%
Black/African-American Males	1,053.0	1,773	29%	986	29%	787	29%
Black/African-American Females	476.0	850	14%	479	14%	371	13%
Hispanic/Latino Males	627.9	315	5%	177	5%	138	5%
Hispanic/Latina Females	148.7	70	1%	34	1%	36	1%
White Males	347.4	2,500	40%	1,386	40%	1,114	40%
White Females	40.9	303	5%	168	5%	135	5%
Multi-Race Males	834.8	233	4%	107	3%	126	5%
Multi-Race Females	192.0	55	1%	29	1%	26	1%
Unknown	*	21	<1%	21	1%	-	-
Total	298.1	6,185		3,432		2,753	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	3,458	71%	1,960	73%	1,498	69%
Injection drug use (IDU)	109	2%	41	2%	68	3%
Male-to-male sexual contact and IDU	221	5%	114	4%	107	5%
Heterosexual contact	267	5%	98	4%	169	8%
Other/unknown	807	17%	479	18%	328	15%
Subtotal	4,862	100%	2,692	100%	2,170	100%
Female adult or adolescent						
Injection drug use	121	10%	51	7%	70	13%
Heterosexual contact	1,019	81%	562	80%	457	82%
Other/unknown	117	9%	88	13%	29	5%
Subtotal	1,257	100%	701	100%	556	100%
Child (<13 yrs at diagnosis)						
Perinatal	59	89%	34	87%	25	93%
Other/unknown	7	11%	5	13%	2	7%
Subtotal	66	100%	39	100%	27	100%
Total	6,185		3,432		2,753	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3,314	54%	1,892	55%	1,422	52%
Injection drug use (IDU) only	106	2%	43	1%	63	2%
Heterosexual contact only	1,287	21%	661	19%	626	23%
Male-to-male sexual contact & IDU	183	3%	92	3%	91	3%
IDU & Heterosexual contact	124	2%	49	1%	75	3%
Male-to-male sexual contact & Heterosexual contact	144	2%	68	2%	76	3%
Male-to-male sexual contact & IDU & Heterosexual contact	38	1%	22	1%	16	1%
Perinatal exposure	62	1%	36	1%	26	1%
Other/unknown	927	15%	569	17%	358	13%
Total	6,185		3,432		2,753	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	2	4%	*	7	<1%	*	3	1%	*	1	<1%	*	-	-	-	-
13-14	*	-	-	*	-	-	*	5	<1%	*	-	-	*	-	-	*	1	<1%	-	-
15-19	*	-	-	*	2	4%	89.2	22	1%	*	3	1%	*	4	<1%	*	-	-	-	-
20-24	*	-	-	62.7	5	9%	387.9	94	4%	197.8	15	4%	30.8	27	1%	94.5	5	2%	-	-
25-29	*	1	13%	60.7	6	11%	863.4	260	10%	380.4	32	8%	134.5	146	5%	564.4	28	10%	-	-
30-34	*	1	13%	66.3	8	14%	1,429.6	406	15%	751.0	59	15%	193.2	219	8%	1,205.5	45	16%	-	-
35-39	*	-	-	*	3	5%	1,263.7	312	12%	562.2	43	11%	232.5	239	9%	981.5	26	9%	-	-
40-44	*	-	-	69.6	6	11%	1,211.2	273	10%	893.0	59	15%	289.1	273	10%	1,455.9	32	11%	2	10%
45-49	*	4	50%	145.1	11	19%	1,397.5	296	11%	679.3	38	10%	334.7	310	11%	1,609.4	30	10%	-	-
50-54	*	1	13%	172.9	10	18%	1,469.4	285	11%	1,077.6	44	11%	504.7	485	17%	2,517.0	37	13%	1	5%
55-64	*	1	13%	*	1	2%	1,398.9	499	19%	1,279.6	68	18%	424.4	829	30%	2,591.3	61	21%	13	62%
65+	*	-	-	*	3	5%	505.6	164	6%	545.6	21	5%	111.9	270	10%	1,197.3	23	8%	5	24%
Total	203.0	8		52.5	57		756.0	2,623		395.9	385		191.8	2,803		509.3	288		21	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	6	86%	22	52%	1,085	62%	211	68%	1,973	79%	160	69%	1	6%
Injection drug use (IDU)	-	-	-	-	51	3%	7	2%	45	2%	6	3%	-	-
Male-to-male sexual contact and IDU	-	-	3	7%	51	3%	11	4%	134	5%	22	10%	-	-
Heterosexual contact	-	-	2	5%	161	9%	31	10%	58	2%	15	6%	-	-
Other/unknown	1	14%	15	36%	409	23%	52	17%	286	11%	28	12%	16	94%
Subtotal	7	100%	42	100%	1,757	100%	312	100%	2,496	100%	231	100%	17	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	51	6%	7	11%	57	19%	6	11%	-	-
Heterosexual contact	1	100%	6	55%	707	86%	57	88%	205	69%	43	80%	-	-
Other/unknown	-	-	5	45%	67	8%	1	2%	35	12%	5	9%	4	100%
Subtotal	1	100%	11	100%	825	100%	65	100%	297	100%	54	100%	4	100%
Child (<13 yrs at diagnosis)														
Perinatal	-	-	4	100%	37	90%	7	88%	8	80%	3	100%	-	-
Other/unknown	-	-	-	-	4	10%	1	13%	2	20%	-	-	-	-
Subtotal	-	-	4	100%	41	100%	8	100%	10	100%	3	100%	-	-
Total	8		57		2,623		385		2,803		288		21	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	6	75%	21	37%	1,007	38%	200	52%	1,926	69%	153	53%	1	5%
Injection drug use (IDU) only	-	-	-	-	53	2%	10	3%	39	1%	4	1%	-	-
Heterosexual contact only	1	13%	8	14%	868	33%	88	23%	264	9%	58	20%	-	-
Male-to-male sexual contact & IDU	-	-	3	5%	41	2%	11	3%	111	4%	17	6%	-	-
IDU & Heterosexual contact	-	-	-	-	49	2%	4	1%	63	2%	8	3%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	1	2%	78	3%	11	3%	47	2%	7	2%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	10	0%	-	-	23	1%	5	2%	-	-
Perinatal exposure	-	-	5	9%	38	1%	7	2%	8	0%	4	1%	-	-
Other/unknown	1	13%	19	33%	479	18%	54	14%	322	11%	32	11%	20	95%
Total	8		57		2,623		385		2,803		288		21	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus, 2016-2020

Living with Diagnosed HIV Infection										
Transmission Category ^a	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	3,217	72%	3,299	72%	3,306	72%	3,352	71%	3,458	71%
Injection drug use (IDU)	97	2%	104	2%	109	2%	108	2%	109	2%
Male-to-male sexual contact and IDU	202	5%	210	5%	202	4%	205	4%	221	5%
Heterosexual contact	242	5%	242	5%	254	6%	264	6%	267	5%
Other/unknown	684	15%	721	16%	736	16%	783	17%	807	17%
Subtotal	4,442	100%	4,576	100%	4,607	100%	4,712	100%	4,862	100%
Female adult or adolescent										
Injection drug use	96	9%	100	9%	105	9%	118	10%	121	10%
Heterosexual contact	825	80%	870	81%	917	81%	960	80%	1,019	81%
Other/unknown	104	10%	104	10%	111	10%	117	10%	117	9%
Subtotal	1,025	100%	1,074	100%	1,133	100%	1,195	100%	1,257	100%
Child (<13 yrs at diagnosis)										
Perinatal	48	89%	50	91%	51	91%	56	89%	59	89%
Other/unknown	6	11%	5	9%	5	9%	7	11%	7	11%
Subtotal	54	100%	55	100%	56	100%	63	100%	66	100%
Total	5,521		5,705		5,796		5,970		6,185	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus, 2016-2020

Living with Diagnosed HIV Infection										
Exposure Category ^b	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3,058	55%	3,141	55%	3,159	55%	3,211	54%	3,314	54%
Injection drug use (IDU) only	88	2%	91	2%	101	2%	102	2%	106	2%
Heterosexual contact only	1,068	19%	1,113	20%	1,172	20%	1,225	21%	1,287	21%
Male-to-male sexual contact & IDU	163	3%	171	3%	167	3%	167	3%	183	3%
IDU & Heterosexual contact	105	2%	113	2%	113	2%	124	2%	124	2%
Male-to-male sexual contact & Heterosexual contact	159	3%	158	3%	147	3%	141	2%	144	2%
Male-to-male sexual contact & IDU & Heterosexual contact	39	1%	39	1%	35	1%	38	1%	38	1%
Perinatal exposure	51	1%	53	1%	54	1%	59	1%	62	1%
Other/unknown	790	14%	826	14%	848	15%	903	15%	927	15%
Total	5,521		5,705		5,796		5,970		6,185	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.