



Department of
Health

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 5

Counties: Carroll, Coshocton, Harrison, Holmes, Jefferson, Stark,
Tuscarawas, Wayne

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2025

Reported new diagnoses of HIV infection in 2024 by disease status and selected characteristics, HIV Prevention Planning Region 5

Characteristic	2024 diagnosis of HIV infection			Disease Status						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex										
Males	10.0	38	86%	32	86%	5	83%	1	100%	
Females	1.5	6	14%	5	14%	1	17%	-	-	
Age at diagnosis (years)										
0-9	*	-	-	-	-	-	-	-	-	
10-14	*	-	-	-	-	-	-	-	-	
15-19	*	1	2%	1	3%	-	-	-	-	
20-24	12.8	6	14%	5	14%	1	17%	-	-	
25-29	17.7	8	18%	7	19%	1	17%	-	-	
30-34	14.7	7	16%	7	19%	-	-	-	-	
35-39	17.7	8	18%	8	22%	-	-	-	-	
40-44	13.2	6	14%	5	14%	1	17%	-	-	
45-49	*	3	7%	1	3%	2	33%	-	-	
50-54	*	1	2%	-	-	1	17%	-	-	
55-64	*	4	9%	3	8%	-	-	1	100%	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b										
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
Black/African American	*	4	9%	4	11%	-	-	-	-	
Hispanic/Latino	30.5	7	16%	7	19%	-	-	-	-	
White	4.3	29	66%	22	59%	6	100%	1	100%	
Multi-Race	*	4	9%	4	11%	-	-	-	-	
Race/Ethnicity ^b and Sex										
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
Black/African American Males	*	4	9%	4	11%	-	-	-	-	
Black/African American Females	*	-	-	-	-	-	-	-	-	
Hispanic/Latino Males	40.9	5	11%	5	14%	-	-	-	-	
Hispanic/Latina Females	*	2	5%	2	5%	-	-	-	-	
White Males	7.7	26	59%	20	54%	5	83%	1	100%	
White Females	*	3	7%	2	5%	1	17%	-	-	
Multi-Race Males	*	3	7%	3	8%	-	-	-	-	
Multi-Race Females	*	1	2%	1	3%	-	-	-	-	
Total	5.7	44		37		6		1		

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by disease status and transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	22	58%	18	56%	4	80%	-	-
Injection drug use (IDU)	2	5%	2	6%	-	-	-	-
Male-to-male sexual contact and IDU	5	13%	5	16%	-	-	-	-
Heterosexual contact	1	3%	1	3%	-	-	-	-
Other/unknown	8	21%	6	19%	1	20%	1	100%
Subtotal	38	100%	32	100%	5	100%	1	100%
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	6	100%	5	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	6	100%	5	100%	1	100%	-	-
Child (<13 years at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	44		37		6		1	

Reported new diagnoses of HIV infection in 2024 by disease status and exposure category, HIV Prevention Planning Region 5

Exposure Category ^b	2024 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	13	30%	10	27%	3	50%	-	-
Injection drug use (IDU) only	1	2%	1	3%	-	-	-	-
Heterosexual contact only	12	27%	9	24%	2	33%	1	100%
Male-to-male sexual contact & IDU	1	2%	1	3%	-	-	-	-
IDU & Heterosexual contact	1	2%	1	3%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	9	20%	8	22%	1	17%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	4	9%	4	11%	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	3	7%	3	8%	-	-	-	-
Total	44		37		6		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 5

2024 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
0-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	1	25%	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	-	-	*	3	43%	*	3	10%	*	-	-
25-29	*	-	-	*	-	-	*	-	-	*	-	-	15.4	6	21%	*	2	50%
30-34	*	-	-	*	-	-	*	2	50%	*	1	14%	*	4	14%	*	-	-
35-39	*	-	-	*	-	-	*	1	25%	*	2	29%	*	3	10%	*	2	50%
40-44	*	-	-	*	-	-	*	-	-	*	1	14%	12.3	5	17%	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	3	10%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	1	3%	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	4	14%	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	*	4	-	30.5	7	-	4.3	29	-	*	4	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 5

Transmission Category ^c	2024 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	3	75%	4	80%	13	50%	2	67%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	2	8%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	25%	1	20%	3	12%	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	1	4%	-	-
Other/unknown	-	-	-	-	-	-	-	-	7	27%	1	33%
Subtotal	-	-	-	-	4	100%	5	100%	26	100%	3	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	2	100%	3	100%	1	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	2	100%	3	100%	1	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	4	-	7	-	29	-	4	-

Reported new diagnoses of HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 5

2024 diagnosis of HIV infection												
Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	2	50%	2	29%	9	31%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	1	3%	-	-
Heterosexual contact only	-	-	-	-	-	-	2	29%	8	28%	2	50%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	1	3%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	1	3%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	1	25%	2	29%	4	14%	2	50%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	1	25%	1	14%	2	7%	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	3	10%	-	-
Total	-	-	-	-	4	-	7	-	29	-	4	-

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
Asterisk (*) indicates rate not calculated, for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 5

Characteristic	Diagnoses of HIV Infection by Year															Cumulative Diagnoses	
	2020			2021			2022			2023			2024				
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex																	
Males	7.7	29	83%	6.0	23	77%	8.2	31	84%	5.8	22	85%	10.0	38	86%	1,054	83%
Females	1.5	6	17%	1.8	7	23%	1.5	6	16%	*	4	15%	1.5	6	14%	223	17%
Age at diagnosis (years)																	
0-9	*	1	3%	*	-	-	*	-	-	*	-	-	*	-	-	14	1%
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	4	11%	*	1	3%	*	-	-	*	-	-	*	1	2%	46	4%
20-24	17.3	8	23%	10.8	5	17%	12.3	6	16%	*	3	12%	12.8	6	14%	192	15%
25-29	10.7	5	14%	13.2	6	20%	11.1	5	14%	*	2	8%	17.7	8	18%	206	16%
30-34	*	3	9%	17.3	8	27%	10.7	5	14%	14.7	7	27%	14.7	7	16%	215	17%
35-39	*	4	11%	*	-	-	18.2	8	22%	*	4	15%	17.7	8	18%	193	15%
40-44	*	1	3%	*	2	7%	*	3	8%	*	4	15%	13.2	6	14%	145	11%
45-49	*	4	11%	*	3	10%	14.3	6	16%	*	2	8%	*	3	7%	122	10%
50-54	*	2	6%	*	1	3%	*	1	3%	*	1	4%	*	1	2%	58	5%
55-64	*	2	6%	*	3	10%	*	2	5%	*	3	12%	*	4	9%	70	5%
65+	*	1	3%	*	1	3%	*	1	3%	*	-	-	*	-	-	16	1%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	<1%
Black/African American	33.3	12	34%	19.1	7	23%	32.8	12	32%	19.1	7	27%	*	4	9%	288	23%
Hispanic/Latino	*	2	6%	*	1	3%	*	4	11%	*	-	-	30.5	7	16%	42	3%
White	3.1	21	60%	3.0	21	70%	2.9	20	54%	2.8	19	73%	4.3	29	66%	902	71%
Multi-Race	*	-	-	*	1	3%	*	1	3%	*	-	-	*	4	9%	39	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	<1%
Race/Ethnicity ^b and Sex																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Black/African American Males	50.3	9	26%	*	4	13%	49.6	9	24%	38.6	7	27%	*	4	9%	218	17%
Black/African American Females	*	3	9%	*	3	10%	*	3	8%	*	-	-	*	-	-	70	5%
Hispanic/Latino Males	*	2	6%	*	1	3%	*	4	11%	*	-	-	40.9	5	11%	37	3%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	2	5%	5	<1%
White Males	5.3	18	51%	5.3	18	60%	5.0	17	46%	4.5	15	58%	7.7	26	59%	762	60%
White Females	*	3	9%	*	3	10%	*	3	8%	*	4	15%	*	3	7%	140	11%
Multi-Race Males	*	-	-	*	-	-	*	1	3%	*	-	-	*	3	7%	33	3%
Multi-Race Females	*	-	-	*	1	3%	*	-	-	*	-	-	*	1	2%	6	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	<1%
Total	4.6	35		3.9	30		4.8	37		3.4	26		5.7	44		1,277	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 5

Diagnoses of HIV Infection by Year												
Transmission Category ^a	2020		2021		2022		2023		2024		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	21	75%	12	52%	14	45%	14	64%	22	58%	717	68%
Injection drug use (IDU)	-	-	-	-	2	6%	2	9%	2	5%	38	4%
Male-to-male sexual contact and IDU	-	-	2	9%	3	10%	3	14%	5	13%	55	5%
Heterosexual contact	2	7%	2	9%	5	16%	1	5%	1	3%	56	5%
Other/unknown	5	18%	7	30%	7	23%	2	9%	8	21%	184	18%
Subtotal	28	100%	23	100%	31	100%	22	100%	38	100%	1,050	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	23	11%
Heterosexual contact	6	100%	7	100%	6	100%	4	100%	6	100%	171	80%
Other/unknown	-	-	-	-	-	-	-	-	-	-	20	9%
Subtotal	6	100%	7	100%	6	100%	4	100%	6	100%	214	100%
Child (<13 years at diagnosis)												
Perinatal	1	100%	-	-	-	-	-	-	-	-	13	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-	13	100%
Total	35		30		37		26		44		1,277	

Trends in reported diagnoses of HIV infection by year of diagnosis (2020-2024) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 5

Diagnoses of HIV Infection by Year												
Exposure Category ^b	2020		2021		2022		2023		2024		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	13	37%	11	37%	10	27%	7	27%	13	30%	446	35%
Injection drug use (IDU) only	-	-	-	-	1	3%	-	-	1	2%	10	1%
Heterosexual contact only	10	29%	11	37%	14	38%	7	27%	12	27%	296	23%
Male-to-male sexual contact & IDU	-	-	1	3%	3	8%	1	4%	1	2%	23	2%
IDU & Heterosexual contact	-	-	-	-	1	3%	2	8%	1	2%	51	4%
Male-to-male sexual contact & Heterosexual contact	8	23%	1	3%	4	11%	7	27%	9	20%	270	21%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	3%	-	-	2	8%	4	9%	32	3%
Perinatal exposure	1	3%	-	-	-	-	-	-	-	-	13	1%
Other/unknown	3	9%	5	17%	4	11%	-	-	3	7%	136	11%
Total	35		30		37		26		44		1,277	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2020-2024) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2024.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 5

Characteristic	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex						
Males	10	71%	3	60%	7	78%
Females	4	29%	2	40%	2	22%
Age at AIDS diagnosis (years)						
0-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	1	7%	-	-	1	11%
25-29	2	14%	1	20%	1	11%
30-34	1	7%	-	-	1	11%
35-39	1	7%	1	20%	-	-
40-44	1	7%	-	-	1	11%
45-49	3	21%	1	20%	2	22%
50-54	2	14%	-	-	2	22%
55-64	3	21%	2	40%	1	11%
65+	-	-	-	-	-	-
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African American	2	14%	1	20%	1	11%
Hispanic/Latino	2	14%	2	40%	-	-
White	10	71%	2	40%	8	89%
Multi-Race	-	-	-	-	-	-
Race/Ethnicity^b and Sex						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African American Males	2	14%	1	20%	1	11%
Black/African American Females	-	-	-	-	-	-
Hispanic/Latino Males	1	7%	1	20%	-	-
Hispanic/Latina Females	1	7%	1	20%	-	-
White Males	7	50%	1	20%	6	67%
White Females	3	21%	1	20%	2	22%
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
Total	14		5		9	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 5

Transmission Category ^b	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	7	70%	2	67%	5	71%
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	1	10%	1	33%	-	-
Other/unknown	2	20%	-	-	2	29%
Subtotal	10	100%	3	100%	7	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	4	100%	2	100%	2	100%
Other/unknown	-	-	-	-	-	-
Subtotal	4	100%	2	100%	2	100%
Child (<13 years at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	14		5		9	

Reported diagnoses of AIDS in 2024 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 5

Exposure Category ^c	AIDS Diagnoses					
	2024 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	6	43%	2	40%	4	44%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	7	50%	3	60%	4	44%
Male-to-male sexual contact & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	1	7%	-	-	1	11%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Total	14		5		9	

Notes:

Data reflects all persons with an AIDS diagnosis in 2024 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 5

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex												
Males	7	78%	14	88%	15	100%	11	79%	10	71%	600	84%
Females	2	22%	2	13%	-	-	3	21%	4	29%	115	16%
Age at AIDS diagnosis (years)												
0-9	1	11%	-	-	-	-	-	-	-	-	10	1%
10-14	-	-	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-	3	<1%
20-24	1	11%	1	6%	2	13%	-	-	1	7%	34	5%
25-29	1	11%	4	25%	2	13%	-	-	2	14%	90	13%
30-34	1	11%	2	13%	1	7%	5	36%	1	7%	122	17%
35-39	-	-	-	-	1	7%	1	7%	1	7%	120	17%
40-44	-	-	1	6%	-	-	1	7%	1	7%	113	16%
45-49	1	11%	2	13%	3	20%	-	-	3	21%	83	12%
50-54	3	33%	2	13%	-	-	3	21%	2	14%	60	8%
55-64	1	11%	3	19%	5	33%	4	29%	3	21%	64	9%
65+	-	-	1	6%	1	7%	-	-	-	-	16	2%
Race/Ethnicity ^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	1	7%	-	-	-	-	1	<1%
Black/African American	2	22%	2	13%	2	13%	4	29%	2	14%	149	21%
Hispanic/Latino	1	11%	-	-	2	13%	-	-	2	14%	21	3%
White	6	67%	13	81%	10	67%	10	71%	10	71%	526	74%
Multi-Race	-	-	1	6%	-	-	-	-	-	-	18	3%
Race/Ethnicity ^a and Sex												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	1	7%	-	-	-	-	1	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
Black/African American Males	1	11%	2	13%	2	13%	3	21%	2	14%	112	16%
Black/African American Females	1	11%	-	-	-	-	1	7%	-	-	37	5%
Hispanic/Latino Males	1	11%	-	-	2	13%	-	-	1	7%	18	3%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	1	7%	3	<1%
White Males	5	56%	11	69%	10	67%	8	57%	7	50%	453	63%
White Females	1	11%	2	13%	-	-	2	14%	3	21%	73	10%
Multi-Race Males	-	-	1	6%	-	-	-	-	-	-	16	2%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Total	9		16		15		14		14		715	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic.

Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	5	83%	5	36%	9	60%	4	36%	7	70%	409	69%
Injection drug use (IDU)	-	-	1	7%	2	13%	1	9%	-	-	25	4%
Male-to-male sexual contact and IDU	-	-	2	14%	-	-	3	27%	-	-	33	6%
Heterosexual contact	-	-	1	7%	2	13%	1	9%	1	10%	35	6%
Other/unknown	1	17%	5	36%	2	13%	2	18%	2	20%	93	16%
Subtotal	6	100%	14	100%	15	100%	11	100%	10	100%	595	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	14	13%
Heterosexual contact	2	100%	2	100%	-	-	3	100%	4	100%	89	82%
Other/unknown	-	-	-	-	-	-	-	-	-	-	5	5%
Subtotal	2	100%	2	100%	-	-	3	100%	4	100%	108	100%
Child (<13 years at diagnosis)												
Perinatal	1	100%	-	-	-	-	-	-	-	-	12	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-	12	100%
Total	9		16		15		14		14		715	

Trends in reported diagnoses of AIDS by year of diagnosis (2020-2024) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 5

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2020		2021		2022		2023		2024			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	33%	4	25%	7	47%	1	7%	6	43%	244	34%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	3	<1%
Heterosexual contact only	3	33%	3	19%	3	20%	5	36%	7	50%	162	23%
Male-to-male sexual contact & IDU	-	-	1	6%	-	-	-	-	-	-	11	2%
IDU & Heterosexual contact	-	-	1	6%	2	13%	1	7%	-	-	36	5%
Male-to-male sexual contact & Heterosexual contact	2	22%	1	6%	2	13%	3	21%	1	7%	164	23%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	6%	-	-	3	21%	-	-	22	3%
Perinatal exposure	1	11%	-	-	-	-	-	-	-	-	12	2%
Other/unknown	-	-	5	31%	1	7%	1	7%	-	-	61	9%
Total	9		16		15		14		14		715	

Notes:

AIDS diagnoses by year (2020-2024) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of AIDS diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed in Ohio since the beginning of the epidemic through 2024.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.



Department of
Health

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 5

Counties: Carroll, Coshocton, Harrison, Holmes, Jefferson, Stark,
Tuscarawas, Wayne

Reported persons living with diagnosed HIV infection in 2024 by current disease status and selected characteristics, HIV Prevention Planning Region 5

Characteristic	Living with diagnosed HIV infection in 2024			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	170.5	648	79%	371	79%	277	80%
Females	43.5	169	21%	98	21%	71	20%
Age at end of year (years)							
0-9	*	3	<1%	1	<1%	2	1%
10-14	*	-	-	-	-	-	-
15-19	12.0	6	1%	6	1%	-	-
20-24	47.1	22	3%	20	4%	2	1%
25-29	110.6	50	6%	43	9%	7	2%
30-34	180.7	86	11%	61	13%	25	7%
35-39	196.5	89	11%	60	13%	29	8%
40-44	194.2	88	11%	59	13%	29	8%
45-49	152.9	65	8%	36	8%	29	8%
50-54	207.6	95	12%	48	10%	47	14%
55-64	194.2	193	24%	80	17%	113	32%
65+	74.3	120	15%	55	12%	65	19%
Race/Ethnicity ^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	87.4	6	1%	5	1%	1	<1%
Black/African American	499.3	185	23%	110	23%	75	22%
Hispanic/Latino	169.7	39	5%	22	5%	17	5%
White	79.6	543	66%	304	65%	239	69%
Multi-Race	232.1	43	5%	27	6%	16	5%
Unknown	*	1	<1%	1	<1%	-	-
Race/Ethnicity ^b and Sex							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	4	<1%	3	1%	1	<1%
Asian/Pacific Islander Females	*	2	<1%	2	<1%	-	-
Black/African American Males	763.9	140	17%	83	18%	57	16%
Black/African American Females	240.4	45	6%	27	6%	18	5%
Hispanic/Latino Males	253.8	31	4%	20	4%	11	3%
Hispanic/Latina Females	74.3	8	1%	2	<1%	6	2%
White Males	130.8	440	54%	243	52%	197	57%
White Females	29.8	103	13%	61	13%	42	12%
Multi-Race Males	342.5	32	4%	21	4%	11	3%
Multi-Race Females	119.8	11	1%	6	1%	5	1%
Unknown	*	1	<1%	1	<1%	-	-
Total	106.3	817		469		348	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2024 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by current disease status and transmission category, HIV Prevention Planning Region 5

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	447	70%	247	67%	200	73%
Injection drug use (IDU)	19	3%	10	3%	9	3%
Male-to-male sexual contact and IDU	52	8%	33	9%	19	7%
Heterosexual contact	32	5%	19	5%	13	5%
Other/unknown	91	14%	57	16%	34	12%
Subtotal	641	100%	366	100%	275	100%
Female adult or adolescent						
Injection drug use	15	9%	8	9%	7	10%
Heterosexual contact	140	86%	79	84%	61	88%
Other/unknown	8	5%	7	7%	1	1%
Subtotal	163	100%	94	100%	69	100%
Child (<13 years at diagnosis)						
Perinatal	11	85%	7	78%	4	100%
Other/unknown	2	15%	2	22%	-	-
Subtotal	13	100%	9	100%	4	100%
Total	817		469		348	

Reported persons living with diagnosed HIV infection in 2024 by current disease status and exposure category, HIV Prevention Planning Region 5

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2024		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	294	36%	169	36%	125	36%
Injection drug use (IDU) only	7	1%	5	1%	2	1%
Heterosexual contact only	213	26%	118	25%	95	27%
Male-to-male sexual contact & IDU	25	3%	19	4%	6	2%
IDU & Heterosexual contact	27	3%	13	3%	14	4%
Male-to-male sexual contact & Heterosexual contact	152	19%	78	17%	74	21%
Male-to-male sexual contact & IDU & Heterosexual contact	27	3%	14	3%	13	4%
Perinatal exposure	11	1%	7	1%	4	1%
Other/unknown	61	7%	46	10%	15	4%
Total	817		469		348	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and age, HIV Prevention Planning Region 5

Living with diagnosed HIV infection in 2024																				
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
0-9	*	-	-	*	1	17%	*	-	-	*	1	3%	*	1	<1%	*	-	-	-	-
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	2	1%	*	-	-	*	4	1%	*	-	-	-	-
20-24	*	-	-	*	1	17%	336.7	9	5%	*	-	-	30.4	12	2%	*	-	-	-	-
25-29	*	-	-	*	-	-	816.5	21	11%	417.9	7	18%	46.3	18	3%	*	4	9%	-	-
30-34	*	-	-	*	1	17%	923.7	27	15%	*	2	5%	118.6	49	9%	619.5	7	16%	-	-
35-39	*	-	-	*	-	-	788.4	18	10%	363.0	6	15%	140.0	56	10%	1,258.7	9	21%	-	-
40-44	*	-	-	*	2	33%	750.5	16	9%	*	3	8%	152.9	62	11%	697.4	5	12%	-	-
45-49	*	-	-	*	-	-	711.4	14	8%	*	2	5%	122.7	47	9%	*	2	5%	-	-
50-54	*	-	-	*	-	-	692.0	14	8%	636.9	6	15%	170.0	71	13%	*	4	9%	-	-
55-64	*	-	-	*	1	17%	1,046.0	40	22%	456.0	7	18%	146.3	135	25%	1,485.9	10	23%	-	-
65+	*	-	-	*	-	-	429.3	24	13%	303.6	5	13%	57.9	88	16%	*	2	5%	1	100%
Total	*	-	-	87.4	6	-	499.3	185	-	169.7	39	-	79.6	543	-	232.1	43	-	1	-

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and transmission category, HIV Prevention Planning Region 5

Living with diagnosed HIV infection in 2024														
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	2	67%	91	66%	24	80%	310	71%	20	63%	-	-
Injection drug use (IDU)	-	-	-	-	5	4%	-	-	12	3%	2	6%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	5	4%	4	13%	38	9%	5	16%	-	-
Heterosexual contact	-	-	-	-	13	9%	-	-	17	4%	2	6%	-	-
Other/unknown	-	-	1	33%	23	17%	2	7%	61	14%	3	9%	1	100%
Subtotal	-	-	3	100%	137	100%	30	100%	438	100%	32	100%	1	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	2	5%	1	13%	10	10%	2	18%	-	-
Heterosexual contact	-	-	2	100%	41	93%	7	88%	81	83%	9	82%	-	-
Other/unknown	-	-	-	-	1	2%	-	-	7	7%	-	-	-	-
Subtotal	-	-	2	100%	44	100%	8	100%	98	100%	11	100%	-	-
Child (<13 years at diagnosis)														
Perinatal	-	-	1	100%	4	100%	1	100%	5	71%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	29%	-	-	-	-
Subtotal	-	-	1	100%	4	100%	1	100%	7	100%	-	-	-	-
Total	-	-	6	-	185	-	39	-	543	-	43	-	1	-

Reported persons living with diagnosed HIV infection in 2024 by race/ethnicity and exposure category, HIV Prevention Planning Region 5

Living with diagnosed HIV infection in 2024														
Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	2	33%	56	30%	19	49%	206	38%	11	26%	-	-
Injection drug use (IDU) only	-	-	-	-	3	2%	-	-	4	1%	-	-	-	-
Heterosexual contact only	-	-	3	50%	65	35%	8	21%	125	23%	12	28%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	2	1%	-	-	19	3%	4	9%	-	-
IDU & Heterosexual contact	-	-	-	-	4	2%	1	3%	18	3%	4	9%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	35	19%	5	13%	103	19%	9	21%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	3	2%	4	10%	19	3%	1	2%	-	-
Perinatal exposure	-	-	1	17%	4	2%	1	3%	5	1%	-	-	-	-
Other/unknown	-	-	-	-	13	7%	1	3%	44	8%	2	5%	1	100%
Total	-	-	6	-	185	-	39	-	543	-	43	-	1	-

Notes:
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2024. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2024, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 5, 2020-2024

Living with Diagnosed HIV Infection															
Characteristic	2020			2021			2022			2023			2024		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex															
Males	157.7	592	81%	155.3	591	79%	154.8	588	78%	156.4	593	78%	170.5	648	79%
Females	36.7	143	19%	40.1	156	21%	41.8	162	22%	42.0	163	22%	43.5	169	21%
Age at end of year (years)															
0-9	*	2	<1%	*	2	<1%	*	3	<1%	*	3	<1%	*	3	<1%
10-14	*	4	1%	*	3	<1%	*	3	<1%	*	-	-	*	-	-
15-19	12.3	6	1%	12.0	6	1%	*	3	<1%	*	4	1%	12.0	6	1%
20-24	65.0	30	4%	54.0	25	3%	51.4	25	3%	57.9	27	4%	47.1	22	3%
25-29	125.9	59	8%	129.5	59	8%	117.2	53	7%	102.6	46	6%	110.6	50	6%
30-34	170.3	76	10%	177.8	82	11%	157.7	74	10%	162.0	77	10%	180.7	86	11%
35-39	150.9	66	9%	148.8	66	9%	170.8	75	10%	160.2	71	9%	196.5	89	11%
40-44	134.8	57	8%	117.8	52	7%	127.7	57	8%	157.1	71	9%	194.2	88	11%
45-49	167.4	74	10%	157.8	68	9%	162.3	68	9%	147.8	62	8%	152.9	65	8%
50-54	237.2	111	15%	228.5	110	15%	222.6	106	14%	232.1	109	14%	207.6	95	12%
55-64	159.0	171	23%	170.3	183	24%	175.4	182	24%	178.4	181	24%	194.2	193	24%
65+	51.6	79	11%	59.6	91	12%	64.9	101	13%	66.4	105	14%	74.3	120	15%
Race/Ethnicity ^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	4	1%	79.7	5	1%	93.5	6	1%	77.8	5	1%	87.4	6	1%
Black/African American	443.8	160	22%	438.9	161	22%	456.4	167	22%	471.9	173	23%	499.3	185	23%
Hispanic/Latino	167.6	28	4%	147.0	26	3%	173.5	32	4%	143.4	31	4%	169.7	39	5%
White	74.0	509	69%	75.8	524	70%	74.8	514	69%	75.2	514	68%	79.6	543	66%
Multi-Race	205.9	33	4%	180.7	30	4%	174.4	30	4%	180.4	32	4%	232.1	43	5%
Unknown	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Race/Ethnicity ^b and Sex															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	2	<1%	*	2	<1%	*	3	<1%	*	3	<1%	*	4	<1%
Asian/Pacific Islander Females	*	2	<1%	*	3	<1%	*	3	<1%	*	2	<1%	*	2	<1%
Black/African American Males	676.5	121	16%	640.1	117	16%	672.2	122	16%	683.7	124	16%	763.9	140	17%
Black/African American Females	214.7	39	5%	239.1	44	6%	244.0	45	6%	264.5	49	6%	240.4	45	6%
Hispanic/Latino Males	250.1	22	3%	203.5	19	3%	256.7	25	3%	210.0	24	3%	253.8	31	4%
Hispanic/Latina Females	75.8	6	1%	83.8	7	1%	80.4	7	1%	68.7	7	1%	74.3	8	1%
White Males	124.6	420	57%	126.1	430	58%	122.5	416	55%	123.8	417	55%	130.8	440	54%
White Females	25.4	89	12%	26.8	94	13%	28.2	98	13%	28.0	97	13%	29.8	103	13%
Multi-Race Males	320.7	26	4%	261.9	22	3%	241.8	21	3%	268.5	24	3%	342.5	32	4%
Multi-Race Females	88.4	7	1%	97.6	8	1%	105.6	9	1%	91.0	8	1%	119.8	11	1%
Unknown	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Total	96.1	735		97.0	747		97.7	750		98.5	756		106.3	817	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 5, 2020-2024

Living with Diagnosed HIV Infection										
Transmission Category ^a	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	427	72%	429	73%	415	71%	417	71%	447	70%
Injection drug use (IDU)	19	3%	16	3%	15	3%	15	3%	19	3%
Male-to-male sexual contact and IDU	35	6%	39	7%	45	8%	46	8%	52	8%
Heterosexual contact	30	5%	25	4%	30	5%	30	5%	32	5%
Other/unknown	78	13%	79	13%	79	14%	80	14%	91	14%
Subtotal	589	100%	588	100%	584	100%	588	100%	641	100%
Female adult or adolescent										
Injection drug use	11	8%	14	10%	16	10%	13	8%	15	9%
Heterosexual contact	116	87%	125	85%	131	85%	137	87%	140	86%
Other/unknown	7	5%	8	5%	7	5%	7	4%	8	5%
Subtotal	134	100%	147	100%	154	100%	157	100%	163	100%
Child (<13 years at diagnosis)										
Perinatal	11	92%	11	92%	11	92%	10	91%	11	85%
Other/unknown	1	8%	1	8%	1	8%	1	9%	2	15%
Subtotal	12	100%	12	100%	12	100%	11	100%	13	100%
Total	735		747		750		756		817	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 5, 2020-2024

Living with Diagnosed HIV Infection										
Exposure Category ^b	2020		2021		2022		2023		2024	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	273	37%	277	37%	271	36%	278	37%	294	36%
Injection drug use (IDU) only	7	1%	7	1%	7	1%	5	1%	7	1%
Heterosexual contact only	179	24%	183	24%	192	26%	200	26%	213	26%
Male-to-male sexual contact & IDU	15	2%	17	2%	25	3%	23	3%	25	3%
IDU & Heterosexual contact	23	3%	23	3%	24	3%	23	3%	27	3%
Male-to-male sexual contact & Heterosexual contact	153	21%	151	20%	143	19%	138	18%	152	19%
Male-to-male sexual contact & IDU & Heterosexual contact	20	3%	22	3%	20	3%	23	3%	27	3%
Perinatal exposure	12	2%	12	2%	11	1%	10	1%	11	1%
Other/unknown	53	7%	55	7%	57	8%	56	7%	61	7%
Total	735		747		750		756		817	

Notes:
Living with diagnosed HIV infection by year (2020-2024) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using Sex and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.
^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2025.