

Persons Living with Diagnosed HIV Infection Reported in Licking County

As of June 30, 2021

Reported persons living with diagnosed HIV infection in 2020 by current disease status and selected characteristics, Licking County

Characteristic	Living with diagnosed HIV infection in 2020			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex at birth							
Males	181.5	159	80%	79	84%	80	75%
Females	45.3	41	21%	15	16%	26	25%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	2	1%	1	1%	1	1%
20-24	*	3	2%	3	3%	-	-
25-29	153.7	16	8%	12	13%	4	4%
30-34	146.2	16	8%	10	11%	6	6%
35-39	87.6	10	5%	6	6%	4	4%
40-44	150.0	16	8%	4	4%	12	11%
45-49	236.2	26	13%	14	15%	12	11%
50-54	269.1	32	16%	14	15%	18	17%
55-64	248.6	62	31%	24	26%	38	36%
65+	55.5	17	9%	6	6%	11	10%
Race/Ethnicity^b							
American Indian/Alaska Native	*	1	1%	1	1%	-	-
Asian/Pacific Islander	*	3	2%	2	2%	1	1%
Black/African-American	460.5	34	17%	19	20%	15	14%
Hispanic/Latinx	201.4	8	4%	4	4%	4	4%
White	89.1	140	70%	62	66%	78	74%
Multi-Race	312.7	13	7%	5	5%	8	8%
Unknown	*	1	1%	1	1%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	1	1%	1	1%	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	2	1%	2	2%	-	-
Asian/Pacific Islander Females	*	1	1%	-	-	1	1%
Black/African-American Males	596.0	23	12%	15	16%	8	8%
Black/African-American Females	312.1	11	6%	4	4%	7	7%
Hispanic/Latino Males	*	4	2%	2	2%	2	2%
Hispanic/Latina Females	*	4	2%	2	2%	2	2%
White Males	154.4	119	60%	55	59%	64	60%
White Females	26.2	21	11%	7	7%	14	13%
Multi-Race Males	486.9	10	5%	4	4%	6	6%
Multi-Race Females	*	3	2%	1	1%	2	2%
Unknown	*	1	1%	1	1%	-	-
Total	112.3	200		94		106	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2020 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are non-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by current disease status and transmission category, Licking County

Transmission Category ^a	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	109	70%	52	68%	57	72%
Injection drug use (IDU)	7	4%	2	3%	5	6%
Male-to-male sexual contact and IDU	9	6%	2	3%	7	9%
Heterosexual contact	9	6%	4	5%	5	6%
Other/unknown	22	14%	17	22%	5	6%
Subtotal	156	100%	77	100%	79	100%
Female adult or adolescent						
Injection drug use	3	8%	1	7%	2	8%
Heterosexual contact	32	82%	9	64%	23	92%
Other/unknown	4	10%	4	29%	-	-
Subtotal	39	100%	14	100%	25	100%
Child (<13 yrs at diagnosis)						
Perinatal	5	100%	3	100%	2	100%
Other/unknown	-	-	-	-	-	-
Subtotal	5	100%	3	100%	2	100%
Total	200		94		106	

Reported persons living with diagnosed HIV infection in 2020 by current disease status and exposure category, Licking County

Exposure Category ^b	Living with diagnosed HIV infection		Current Disease Status			
	in 2020		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	104	52%	49	52%	55	52%
Injection drug use (IDU) only	4	2%	-	-	4	4%
Heterosexual contact only	41	21%	13	14%	28	26%
Male-to-male sexual contact & IDU	8	4%	2	2%	6	6%
IDU & Heterosexual contact	6	3%	3	3%	3	3%
Male-to-male sexual contact & Heterosexual contact	5	3%	3	3%	2	2%
Male-to-male sexual contact & IDU & Heterosexual contact	1	1%	-	-	1	1%
Perinatal exposure	5	3%	3	3%	2	2%
Other/unknown	26	13%	21	22%	5	5%
Total	200		94		106	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and age, Licking County

Living with diagnosed HIV infection in 2020

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	<5	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
13-14	*	<5	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	<5	-	*	-	-	*	1	3%	*	1	13%	*	-	-	*	-	-	-	-
20-24	*	<5	-	*	-	-	*	1	3%	*	-	-	*	1	1%	*	1	8%	-	-
25-29	*	<5	-	*	-	-	*	4	12%	*	1	13%	113.1	10	7%	*	1	8%	-	-
30-34	*	<5	-	*	-	-	*	2	6%	*	-	-	107.3	10	7%	*	4	31%	-	-
35-39	*	<5	-	*	-	-	*	2	6%	*	-	-	80.0	8	6%	*	-	-	-	-
40-44	*	<5	-	*	-	-	*	2	6%	*	-	-	148.7	14	10%	*	-	-	-	-
45-49	*	<5	-	*	3	100%	*	4	12%	*	-	-	194.7	19	14%	*	-	-	-	-
50-54	*	<5	-	*	-	-	*	1	3%	*	-	-	250.0	27	19%	*	3	23%	-	-
55-64	*	<5	-	*	-	-	1,484.6	14	41%	1,908.4	5	63%	173.1	40	29%	*	3	23%	-	-
65+	*	<5	-	*	-	-	*	3	9%	*	1	13%	37.8	11	8%	*	1	8%	1	100%
Total	*	1		*	3		460.5	34		201.4	8		89.1	140		312.7	13		1	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and transmission category, Licking County

Living with diagnosed HIV infection in 2020

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	1	100%	1	50%	9	41%	2	50%	89	75%	7	78%	-	-
Injection drug use (IDU)	-	-	-	-	2	9%	1	25%	4	3%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	5%	-	-	7	6%	1	11%	-	-
Heterosexual contact	-	-	1	50%	3	14%	1	25%	3	3%	1	11%	-	-
Other/unknown	-	-	-	-	7	32%	-	-	15	13%	-	-	-	-
Subtotal	1	100%	2	100%	22	100%	4	100%	118	100%	9	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	-	-	-	-	3	14%	-	-	-	-
Heterosexual contact	-	-	1	100%	10	100%	3	100%	16	76%	2	67%	-	-
Other/unknown	-	-	-	-	-	-	-	-	2	10%	1	33%	1	100%
Subtotal	-	-	1	100%	10	100%	3	100%	21	100%	3	100%	1	100%
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	2	100%	1	100%	1	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	2	100%	1	100%	1	100%	1	100%	-	-
Total	1		3		34		8		140		13		1	

Reported persons living with diagnosed HIV infection in 2020 by race/ethnicity and exposure category, Licking County

Living with diagnosed HIV infection in 2020

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	100%	1	33%	9	26%	2	25%	84	60%	7	54%	-	-
Injection drug use (IDU) only	-	-	-	-	1	3%	1	13%	2	1%	-	-	-	-
Heterosexual contact only	-	-	2	67%	13	38%	4	50%	19	14%	3	23%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	1	3%	-	-	7	5%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	1	3%	-	-	5	4%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	5	4%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	1	8%	-	-
Perinatal exposure	-	-	-	-	2	6%	1	13%	1	1%	1	8%	-	-
Other/unknown	-	-	-	-	7	21%	-	-	17	12%	1	8%	1	100%
Total	1		3		34		8		140		13		1	

Notes:
 Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2020. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2020, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
 Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
 Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Licking County, 2016-2020

Characteristic	Living with Diagnosed HIV Infection														
	2016			2017			2018			2019			2020		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	170.3	144	80%	174.9	149	81%	177.4	153	81%	172.8	150	78%	181.5	159	80%
Females	39.9	35	20%	39.7	35	19%	40.2	36	19%	47.8	43	22%	45.3	41	21%
Age at end of year															
<13	*	2	1%	*	1	1%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	1	1%	*	1	1%	*	-	-	*	-	-
15-19	*	3	2%	*	-	-	*	1	1%	*	1	1%	*	2	1%
20-24	46.9	5	3%	55.4	6	3%	54.6	6	3%	*	4	2%	*	3	2%
25-29	153.7	15	8%	99.4	10	5%	126.5	13	7%	115.8	12	6%	153.7	16	8%
30-34	77.2	8	4%	106.5	11	6%	94.6	10	5%	92.5	10	5%	146.2	16	8%
35-39	105.9	11	6%	112.9	12	7%	137.0	15	8%	143.4	16	8%	87.6	10	5%
40-44	164.8	17	9%	178.1	18	10%	166.6	17	9%	202.2	21	11%	150.0	16	8%
45-49	236.6	28	16%	262.6	31	17%	265.1	31	16%	253.2	29	15%	236.2	26	13%
50-54	277.9	35	20%	268.8	33	18%	316.4	38	20%	330.2	39	20%	269.1	32	16%
55-64	180.2	43	24%	214.1	52	28%	186.5	46	24%	184.3	46	24%	248.6	62	31%
65+	44.1	12	7%	32.1	9	5%	38.1	11	6%	50.6	15	8%	55.5	17	9%
Race/Ethnicity^b															
American Indian/Alaska Native	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
Asian/Pacific Islander	*	1	1%	*	1	1%	*	3	2%	*	3	2%	*	3	2%
Black/African-American	318.8	21	12%	353.1	24	13%	416.6	29	15%	515.7	37	19%	460.5	34	17%
Hispanic/Latinx	*	4	2%	148.9	5	3%	*	4	2%	136.1	5	3%	201.4	8	4%
White	87.5	137	77%	89.2	140	76%	86.2	136	72%	85.1	134	69%	89.1	140	70%
Multi-Race	369.9	13	7%	309.6	11	6%	376.4	14	7%	302.1	12	6%	312.7	13	7%
Unknown	*	2	1%	*	2	1%	*	2	1%	*	1	1%	*	1	1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	2	1%	*	2	1%	*	2	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	1%	*	1	1%	*	1	1%
Black/African-American Males	438.5	15	8%	508.0	18	10%	552.8	20	11%	699.9	26	13%	596.0	23	12%
Black/African-American Females	189.5	6	3%	184.4	6	3%	269.2	9	5%	317.9	11	6%	312.1	11	6%
Hispanic/Latino Males	*	2	1%	*	3	2%	*	2	1%	*	2	1%	*	4	2%
Hispanic/Latina Females	*	2	1%	*	2	1%	*	2	1%	*	3	2%	*	4	2%
White Males	148.4	114	64%	152.1	117	64%	149.9	116	61%	142.5	110	57%	154.4	119	60%
White Females	28.8	23	13%	28.8	23	13%	24.9	20	11%	29.9	24	12%	26.2	21	11%
Multi-Race Males	572.4	10	6%	454.5	8	4%	607.1	11	6%	461.3	9	5%	486.9	10	5%
Multi-Race Females	*	3	2%	*	3	2%	*	3	2%	*	3	2%	*	3	2%
Unknown	*	2	1%	*	2	1%	*	2	1%	*	1	1%	*	1	1%
Total	104.0	179		106.1	184		107.5	189		109.1	193		112.3	200	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Licking County, 2016-2020

Living with Diagnosed HIV Infection										
Transmission Category ^a	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	96	68%	97	66%	101	67%	97	66%	109	70%
Injection drug use (IDU)	6	4%	6	4%	5	3%	6	4%	7	4%
Male-to-male sexual contact and IDU	15	11%	15	10%	10	7%	8	5%	9	6%
Heterosexual contact	5	4%	5	3%	11	7%	10	7%	9	6%
Other/unknown	19	13%	23	16%	23	15%	26	18%	22	14%
Subtotal	141	100%	146	100%	150	100%	147	100%	156	100%
Female adult or adolescent										
Injection drug use	3	9%	2	6%	2	6%	3	7%	3	8%
Heterosexual contact	27	82%	28	85%	28	82%	34	83%	32	82%
Other/unknown	3	9%	3	9%	4	12%	4	10%	4	10%
Subtotal	33	100%	33	100%	34	100%	41	100%	39	100%
Child (<13 yrs at diagnosis)										
Perinatal	5	100%	5	100%	4	80%	5	100%	5	100%
Other/unknown	-	-	-	-	1	20%	-	-	-	-
Subtotal	5	100%	5	100%	5	100%	5	100%	5	100%
Total	179		184		189		193		200	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Licking County, 2016-2020

Living with Diagnosed HIV Infection										
Exposure Category ^b	2016		2017		2018		2019		2020	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	91	51%	92	50%	97	51%	91	47%	104	52%
Injection drug use (IDU) only	4	2%	4	2%	3	2%	3	2%	4	2%
Heterosexual contact only	32	18%	33	18%	39	21%	44	23%	41	21%
Male-to-male sexual contact & IDU	14	8%	15	8%	9	5%	7	4%	8	4%
IDU & Heterosexual contact	5	3%	4	2%	4	2%	6	3%	6	3%
Male-to-male sexual contact & Heterosexual contact	5	3%	5	3%	4	2%	6	3%	5	3%
Male-to-male sexual contact & IDU & Heterosexual contact	1	1%	-	-	1	1%	1	1%	1	1%
Perinatal exposure	5	3%	5	3%	4	2%	5	3%	5	3%
Other/unknown	22	12%	26	14%	28	15%	30	16%	26	13%
Total	179		184		189		193		200	

Notes:

Living with diagnosed HIV infection by year (2016-2020) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2021.