

New Diagnoses of HIV Infection Reported in Licking County

*Please note a decrease in new reported diagnoses of HIV in 2020 may not represent a true decline, and may be a reporting artifact as a result of COVID-19.

As of June 30, 2024

Reported new diagnoses of HIV infection in 2023 by disease status and selected characteristics, Licking County

Characteristic	2023 diagnosis of HIV infection			Disease Status						
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS		
				No.	%	No.	%	No.	%	
Sex at birth										
Males	8.8	8	89%	6	86%	2	100%	-	-	
Females	*	1	11%	1	14%	-	-	-	-	
Age at diagnosis (years)										
0-9	*	-	-	-	-	-	-	-	-	
10-14	*	-	-	-	-	-	-	-	-	
15-19	*	-	-	-	-	-	-	-	-	
20-24	*	2	22%	2	29%	-	-	-	-	
25-29	*	2	22%	2	29%	-	-	-	-	
30-34	*	-	-	-	-	-	-	-	-	
35-39	*	3	33%	2	29%	1	50%	-	-	
40-44	*	1	11%	1	14%	-	-	-	-	
45-49	*	-	-	-	-	-	-	-	-	
50-54	*	1	11%	-	-	1	50%	-	-	
55-64	*	-	-	-	-	-	-	-	-	
65+	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b										
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-	
Black/African American	*	1	11%	1	14%	-	-	-	-	
Hispanic/Latino	*	-	-	-	-	-	-	-	-	
White	5.1	8	89%	6	86%	2	100%	-	-	
Multi-Race	*	-	-	-	-	-	-	-	-	
Race/Ethnicity ^b and Sex at birth										
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-	
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-	
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-	
Black/African American Males	*	1	11%	1	14%	-	-	-	-	
Black/African American Females	*	-	-	-	-	-	-	-	-	
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-	
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-	
White Males	9.0	7	78%	5	71%	2	100%	-	-	
White Females	*	1	11%	1	14%	-	-	-	-	
Multi-Race Males	*	-	-	-	-	-	-	-	-	
Multi-Race Females	*	-	-	-	-	-	-	-	-	
Total	4.9	9		7		2		-		

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported new diagnoses of HIV infection in 2023 by disease status and transmission category, Licking County

Transmission Category ^a	2023 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	4	50%	4	67%	-	-	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	1	13%	1	17%	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	3	38%	1	17%	2	100%	-	-
Subtotal	8	100%	6	100%	2	100%	-	-
Female adult or adolescent								
Injection drug use	-	-	-	-	-	-	-	-
Heterosexual contact	1	100%	1	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	1	100%	1	100%	-	-	-	-
Child (<13 years at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	9		7		2		-	

Reported new diagnoses of HIV infection in 2023 by disease status and exposure category, Licking County

Exposure Category ^b	2023 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2	22%	2	29%	-	-	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	3	33%	2	29%	1	50%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	2	22%	2	29%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	1	11%	1	14%	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	1	11%	-	-	1	50%	-	-
Total	9		7		2		-	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and age at diagnosis, Licking County

2023 diagnosis of HIV infection																		
Age at diagnosis (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino*			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	-	-	*	-	-	*	2	25%	*	-	-
25-29	*	-	-	*	-	-	*	1	100%	*	-	-	*	1	13%	*	-	-
30-34	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
35-39	*	-	-	*	-	-	*	-	-	*	-	-	*	3	38%	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	1	13%	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	1	13%	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	*	1	-	*	-	-	5.1	8	-	*	-	-

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and transmission category, Licking County

Transmission Category ^c	2023 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	-	-	-	-	4	57%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	1	100%	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	3	43%	-	-
Subtotal	-	-	-	-	1	100%	-	-	7	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	1	100%	-	-
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-		-		1		-		8		-	

Reported new diagnoses of HIV infection in 2023 by race/ethnicity and exposure category, Licking County

Exposure Category ^d	2023 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-	-	-	2	25%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	-	-	-	-	3	38%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	2	25%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	1	100%	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	13%	-	-
Total	-	-	-	-	1	-	-	-	8	-	-	-

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by selected characteristics, Licking County

Diagnoses of HIV Infection by Year																	
Characteristic	2019			2020			2021			2022			2023			Cumulative Diagnoses	
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex at birth																	
Males	*	3	50%	6.8	6	86%	5.6	5	100%	7.8	7	88%	8.8	8	89%	218	84%
Females	*	3	50%	*	1	14%	*	-	-	*	1	13%	*	1	11%	41	16%
Age at diagnosis (years)																	
0-9	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	1%
10-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	1	14%	*	-	-	*	1	13%	*	-	-	11	4%
20-24	*	-	-	*	-	-	*	1	20%	*	-	-	*	2	22%	38	15%
25-29	*	-	-	*	1	14%	*	2	40%	*	-	-	*	2	22%	33	13%
30-34	*	1	17%	*	2	29%	*	-	-	*	1	13%	*	-	-	40	15%
35-39	*	1	17%	*	1	14%	*	1	20%	*	4	50%	*	3	33%	51	20%
40-44	*	1	17%	*	1	14%	*	1	20%	*	-	-	*	1	11%	36	14%
45-49	*	2	33%	*	1	14%	*	-	-	*	1	13%	*	-	-	20	8%
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	1	11%	12	5%
55-64	*	1	17%	*	-	-	*	-	-	*	1	13%	*	-	-	12	5%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	1%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African American	*	2	33%	*	1	14%	*	2	40%	*	1	13%	*	1	11%	39	15%
Hispanic/Latino	*	1	17%	*	1	14%	*	-	-	*	-	-	*	-	-	9	3%
White	*	3	50%	3.2	5	71%	*	3	60%	4.5	7	88%	5.1	8	89%	200	77%
Multi-Race	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	10	4%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Race/Ethnicity ^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African American Males	*	2	33%	*	1	14%	*	2	40%	*	1	13%	*	1	11%	31	12%
Black/African American Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	8	3%
Hispanic/Latino Males	*	-	-	*	1	14%	*	-	-	*	-	-	*	-	-	7	3%
Hispanic/Latina Females	*	1	17%	*	-	-	*	-	-	*	-	-	*	-	-	2	1%
White Males	*	1	17%	*	4	57%	*	3	60%	7.7	6	75%	9.0	7	78%	174	67%
White Females	*	2	33%	*	1	14%	*	-	-	*	1	13%	*	1	11%	26	10%
Multi-Race Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	5	2%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	5	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Total	3.4	6		3.9	7		2.8	5		4.4	8		4.9	9		259	

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2019-2023) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2023.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by transmission category, Licking County

Transmission Category ^a	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2019		2020		2021		2022		2023			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	4	67%	3	60%	4	57%	4	50%	143	66%
Injection drug use (IDU)	1	33%	-	-	1	20%	-	-	-	-	10	5%
Male-to-male sexual contact and IDU	-	-	1	17%	-	-	2	29%	1	13%	15	7%
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	11	5%
Other/unknown	2	67%	1	17%	1	20%	1	14%	3	38%	38	18%
Subtotal	3	100%	6	100%	5	100%	7	100%	8	100%	217	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	-	-	3	8%
Heterosexual contact	3	100%	1	100%	-	-	-	-	1	100%	32	82%
Other/unknown	-	-	-	-	-	-	1	100%	-	-	4	10%
Subtotal	3	100%	1	100%	-	-	1	100%	1	100%	39	100%
Child (<13 years at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	3	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	3	100%
Total	6		7		5		8		9		259	

Trends in reported diagnoses of HIV infection by year of diagnosis (2019-2023) and cumulative diagnoses by exposure category, Licking County

Exposure Category ^b	Diagnoses of HIV Infection by Year										Cumulative Diagnoses	
	2019		2020		2021		2022		2023			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	4	57%	3	60%	3	38%	2	22%	109	42%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	3	1%
Heterosexual contact only	4	67%	2	29%	1	20%	-	-	3	33%	57	22%
Male-to-male sexual contact & IDU	-	-	1	14%	-	-	1	13%	-	-	9	3%
IDU & Heterosexual contact	1	17%	-	-	1	20%	-	-	-	-	10	4%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	1	13%	2	22%	34	13%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	1	13%	1	11%	6	2%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	3	1%
Other/unknown	1	17%	-	-	-	-	2	25%	1	11%	28	11%
Total	6		7		5		8		9		259	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2019-2023) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2023.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Persons Living with Diagnosed HIV Infection Reported in Licking County

As of June 30, 2024

Reported persons living with diagnosed HIV infection in 2023 by current disease status and selected characteristics, Licking County

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2023			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex at birth							
Males	193.0	175	81%	101	87%	74	75%
Females	43.2	40	19%	15	13%	25	25%
Age at end of year (years)							
0-9	*	2	1%	2	2%	-	-
10-14	*	2	1%	2	2%	-	-
15-19	*	1	<1%	1	1%	-	-
20-24	44.8	5	2%	5	4%	-	-
25-29	115.5	12	6%	12	10%	-	-
30-34	156.0	18	8%	11	9%	7	7%
35-39	175.3	21	10%	14	12%	7	7%
40-44	153.1	18	8%	11	9%	7	7%
45-49	222.6	24	11%	13	11%	11	11%
50-54	218.6	26	12%	14	12%	12	12%
55-64	235.7	58	27%	20	17%	38	38%
65+	85.4	28	13%	11	9%	17	17%
Race/Ethnicity^b							
American Indian/Alaska Native	*	1	<1%	1	1%	-	-
Asian/Pacific Islander	*	4	2%	3	3%	1	1%
Black/African American	502.8	43	20%	25	22%	18	18%
Hispanic/Latino	238.1	11	5%	6	5%	5	5%
White	91.7	144	67%	75	65%	69	70%
Multi-Race	266.8	12	6%	6	5%	6	6%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	1	<1%	1	1%	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	<1%	1	1%	-	-
Asian/Pacific Islander Females	*	3	1%	2	2%	1	1%
Black/African American Males	696.5	31	14%	21	18%	10	10%
Black/African American Females	292.6	12	6%	4	3%	8	8%
Hispanic/Latino Males	303.0	7	3%	4	3%	3	3%
Hispanic/Latina Females	*	4	2%	2	2%	2	2%
White Males	163.8	127	59%	70	60%	57	58%
White Females	21.4	17	8%	5	4%	12	12%
Multi-Race Males	359.9	8	4%	4	3%	4	4%
Multi-Race Females	*	4	2%	2	2%	2	2%
Unknown	*	-	-	-	-	-	-
Total	117.4	215		116		99	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using 2023 U.S. Census estimates.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported persons living with diagnosed HIV infection in 2023 by current disease status and transmission category, Licking County

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2023		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	119	70%	68	70%	51	70%
Injection drug use (IDU)	7	4%	2	2%	5	7%
Male-to-male sexual contact and IDU	13	8%	8	8%	5	7%
Heterosexual contact	9	5%	5	5%	4	5%
Other/unknown	22	13%	14	14%	8	11%
Subtotal	170	100%	97	100%	73	100%
Female adult or adolescent						
Injection drug use	2	5%	-	-	2	8%
Heterosexual contact	30	81%	9	75%	21	84%
Other/unknown	5	14%	3	25%	2	8%
Subtotal	37	100%	12	100%	25	100%
Child (<13 years at diagnosis)						
Perinatal	8	100%	7	100%	1	100%
Other/unknown	-	-	-	-	-	-
Subtotal	8	100%	7	100%	1	100%
Total	215		116		99	

Reported persons living with diagnosed HIV infection in 2023 by current disease status and exposure category, Licking County

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2023		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	87	40%	50	43%	37	37%
Injection drug use (IDU) only	2	1%	-	-	2	2%
Heterosexual contact only	48	22%	20	17%	28	28%
Male-to-male sexual contact & IDU	4	2%	1	1%	3	3%
IDU & Heterosexual contact	7	3%	2	2%	5	5%
Male-to-male sexual contact & Heterosexual contact	32	15%	18	16%	14	14%
Male-to-male sexual contact & IDU & Heterosexual contact	9	4%	7	6%	2	2%
Perinatal exposure	8	4%	7	6%	1	1%
Other/unknown	18	8%	11	9%	7	7%
Total	215		116		99	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and age, Licking County

Living with diagnosed HIV infection in 2023																		
Age at end of year (years)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African American			Hispanic/ Latino ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
0-9	*	<5	-	*	1	25%	*	1	2%	*	-	-	*	-	-	*	-	-
10-14	*	<5	-	*	1	25%	*	1	2%	*	-	-	*	-	-	*	-	-
15-19	*	<5	-	*	-	-	*	1	2%	*	-	-	*	-	-	*	-	-
20-24	*	<5	-	*	-	-	*	2	5%	*	-	-	*	2	1%	*	1	8%
25-29	*	<5	-	*	-	-	932.8	5	12%	*	2	18%	*	3	2%	*	2	17%
30-34	*	<5	-	*	-	-	819.7	5	12%	*	-	-	116.5	11	8%	*	2	17%
35-39	*	<5	-	*	-	-	*	2	5%	*	2	18%	171.4	17	12%	*	-	-
40-44	*	<5	-	*	-	-	*	2	5%	*	-	-	158.1	16	11%	*	-	-
45-49	*	<5	-	*	2	50%	*	3	7%	*	-	-	205.1	19	13%	*	-	-
50-54	*	<5	-	*	-	-	750.8	5	12%	*	1	9%	172.1	18	13%	*	2	17%
55-64	*	<5	-	*	-	-	900.9	10	23%	*	4	36%	174.6	39	27%	*	4	33%
65+	*	<5	-	*	-	-	630.9	6	14%	*	2	18%	61.7	19	13%	*	1	8%
Total	*	1	-	*	4	-	502.8	43	-	238.1	11	-	91.7	144	-	266.8	12	-

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and transmission category, Licking County

Living with diagnosed HIV infection in 2023														
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	1	100%	-	-	15	54%	4	57%	92	73%	7	100%	-	-
Injection drug use (IDU)	-	-	-	-	1	4%	1	14%	5	4%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	4	14%	1	14%	8	6%	-	-	-	-
Heterosexual contact	-	-	1	100%	4	14%	1	14%	3	2%	-	-	-	-
Other/unknown	-	-	-	-	4	14%	-	-	18	14%	-	-	-	-
Subtotal	1	100%	1	100%	28	100%	7	100%	126	100%	7	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	-	-	-	-	2	12%	-	-	-	-
Heterosexual contact	-	-	1	100%	11	92%	3	100%	12	71%	3	75%	-	-
Other/unknown	-	-	-	-	1	8%	-	-	3	18%	1	25%	-	-
Subtotal	-	-	1	100%	12	100%	3	100%	17	100%	4	100%	-	-
Child (<13 years at diagnosis)														
Perinatal	-	-	2	100%	3	100%	1	100%	1	100%	1	100%	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	2	100%	3	100%	1	100%	1	100%	1	100%	-	-
Total	1		4		43		11		144		12		-	

Reported persons living with diagnosed HIV infection in 2023 by race/ethnicity and exposure category, Licking County

Living with diagnosed HIV infection in 2023														
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African American		Hispanic/Latino ^a		White		Multi-Race		Unknown	
Exposure Category ^d	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	100%	-	-	9	21%	2	18%	69	48%	6	50%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	1	9%	1	1%	-	-	-	-
Heterosexual contact only	-	-	2	50%	18	42%	4	36%	21	15%	3	25%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	1	2%	1	9%	2	1%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	1	2%	-	-	6	4%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	6	14%	2	18%	23	16%	1	8%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	3	7%	-	-	6	4%	-	-	-	-
Perinatal exposure	-	-	2	50%	3	7%	1	9%	1	1%	1	8%	-	-
Other/unknown	-	-	-	-	2	5%	-	-	15	10%	1	8%	-	-
Total	1		4		43		11		144		12		-	

Notes:
Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2023. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2023, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.
Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Licking County, 2019-2023

Characteristic	Living with Diagnosed HIV Infection														
	2019			2020			2021			2022			2023		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	167.0	145	78%	171.2	150	79%	169.2	151	79%	174.3	157	81%	193.0	175	81%
Females	46.6	42	22%	44.2	40	21%	45.0	41	21%	41.6	38	19%	43.2	40	19%
Age at end of year (years)															
0-9	*	-	-	*	-	-	*	-	-	*	2	1%	*	2	1%
10-14	*	-	-	*	-	-	*	-	-	*	1	1%	*	2	1%
15-19	*	1	1%	*	2	1%	*	1	1%	*	1	1%	*	1	<1%
20-24	*	4	2%	*	3	2%	*	3	2%	*	2	1%	44.8	5	2%
25-29	115.8	12	6%	124.9	13	7%	116.6	12	6%	87.2	9	5%	115.5	12	6%
30-34	92.5	10	5%	146.2	16	8%	170.0	19	10%	166.6	19	10%	156.0	18	8%
35-39	143.4	16	9%	96.4	11	6%	85.3	10	5%	101.6	12	6%	175.3	21	10%
40-44	202.2	21	11%	168.8	18	9%	151.1	17	9%	182.2	21	11%	153.1	18	8%
45-49	244.5	28	15%	236.2	26	14%	197.2	21	11%	150.7	16	8%	222.6	24	11%
50-54	313.3	37	20%	243.9	29	15%	237.7	29	15%	231.2	28	14%	218.6	26	12%
55-64	180.3	45	24%	228.5	57	30%	245.8	62	32%	245.5	61	31%	235.7	58	27%
65+	43.9	13	7%	48.9	15	8%	58.3	18	9%	72.3	23	12%	85.4	28	13%
Race/Ethnicity^b															
American Indian/Alaska Native	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	<1%
Asian/Pacific Islander	*	2	1%	*	2	1%	*	2	1%	*	4	2%	*	4	2%
Black/African American	501.7	36	19%	460.5	34	18%	452.8	35	18%	520.8	42	22%	502.8	43	20%
Hispanic/Latino	136.1	5	3%	151.0	6	3%	215.4	9	5%	229.5	10	5%	238.1	11	5%
White	82.5	130	70%	84.6	133	70%	86.2	136	71%	81.5	128	66%	91.7	144	67%
Multi-Race	327.3	13	7%	336.8	14	7%	204.8	9	5%	226.1	10	5%	266.8	12	6%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	1%	*	1	1%	*	1	1%	*	1	1%	*	1	<1%
Asian/Pacific Islander Females	*	1	1%	*	1	1%	*	1	1%	*	3	2%	*	3	1%
Black/African American Males	672.9	25	13%	570.1	22	12%	519.4	21	11%	733.2	31	16%	696.5	31	14%
Black/African American Females	317.9	11	6%	340.5	12	6%	379.8	14	7%	286.7	11	6%	292.6	12	6%
Hispanic/Latino Males	*	2	1%	*	3	2%	236.2	5	3%	230.5	5	3%	303.0	7	3%
Hispanic/Latina Females	*	3	2%	*	3	2%	*	4	2%	228.5	5	3%	*	4	2%
White Males	137.3	106	57%	145.3	112	59%	150.4	117	61%	143.9	112	57%	163.8	127	59%
White Females	29.9	24	13%	26.2	21	11%	23.8	19	10%	20.2	16	8%	21.4	17	8%
Multi-Race Males	512.6	10	5%	535.5	11	6%	275.5	6	3%	319.2	7	4%	359.9	8	4%
Multi-Race Females	*	3	2%	*	3	2%	*	3	2%	*	3	2%	*	4	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	105.7	187		106.7	190		106.4	192		107.5	195		117.4	215	

Notes:

Living with diagnosed HIV infection by year (2019-2023) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinos may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Licking County, 2019-2023

Living with Diagnosed HIV Infection										
Transmission Category ^a	2019		2020		2021		2022		2023	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	96	68%	107	73%	108	73%	108	71%	119	70%
Injection drug use (IDU)	6	4%	7	5%	7	5%	7	5%	7	4%
Male-to-male sexual contact and IDU	9	6%	8	5%	9	6%	11	7%	13	8%
Heterosexual contact	10	7%	8	5%	9	6%	9	6%	9	5%
Other/unknown	21	15%	17	12%	15	10%	18	12%	22	13%
Subtotal	142	100%	147	100%	148	100%	153	100%	170	100%
Female adult or adolescent										
Injection drug use	3	8%	3	8%	3	8%	3	9%	2	5%
Heterosexual contact	34	85%	32	84%	33	85%	27	77%	30	81%
Other/unknown	3	8%	3	8%	3	8%	5	14%	5	14%
Subtotal	40	100%	38	100%	39	100%	35	100%	37	100%
Child (<13 years at diagnosis)										
Perinatal	5	100%	5	100%	5	100%	7	100%	8	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	5	100%	5	100%	5	100%	7	100%	8	100%
Total	187		190		192		195		215	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Licking County, 2019-2023

Living with Diagnosed HIV Infection										
Exposure Category ^b	2019		2020		2021		2022		2023	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	66	35%	78	41%	81	42%	81	42%	87	40%
Injection drug use (IDU) only	1	1%	2	1%	2	1%	2	1%	2	1%
Heterosexual contact only	54	29%	48	25%	48	25%	44	23%	48	22%
Male-to-male sexual contact & IDU	4	2%	4	2%	6	3%	5	3%	4	2%
IDU & Heterosexual contact	8	4%	8	4%	8	4%	8	4%	7	3%
Male-to-male sexual contact & Heterosexual contact	30	16%	29	15%	27	14%	27	14%	32	15%
Male-to-male sexual contact & IDU & Heterosexual contact	5	3%	4	2%	3	2%	6	3%	9	4%
Perinatal exposure	5	3%	5	3%	5	3%	7	4%	8	4%
Other/unknown	14	7%	12	6%	12	6%	15	8%	18	8%
Total	187		190		192		195		215	

Notes:

Living with diagnosed HIV infection by year (2019-2023) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV Surveillance Program. Data reported through June 30, 2024.